

REA's

Problem Solvers[®]

Registered Trademark

CHEMISTRY

A Complete Solution Guide to Any Textbook



- ◆ For Homework, Test Preparation, Exams
- ◆ For use with introductory and advanced texts and courses
- ◆ Includes every type of problem that may be assigned by your instructor or given on a test
- ◆ Each problem worked out in step-by-step detail, enabling you to understand the subject fully
- ◆ Will save you hours of time in finding solutions to problems



Research & Education Association

Step-by-Step Solutions to Problems in this Chapter, "Units of Measurement"

LENGTH

• PROBLEM 1

The Eiffel Tower is 984 feet high. Express this height in meters, in kilometers, in centimeters, and in millimeters.

Solution: A meter is equivalent to 39.370 inches. In this problem, the height of the tower in feet must be converted to inches and then the inches can be converted to meters. There are 12 inches in 1 foot. Therefore, feet can be converted to inches by using the factor 12 inches/1 foot.

$$984 \text{ feet} \times 12 \text{ inches/1 foot} = 118 \times 10^2 \text{ inches.}$$

Once the height is found in inches, this can be converted to meters by the factor 1 meter/39.370 inches.

$$11808 \text{ inches} \times 1 \text{ meter/39.370 inches} = 300 \text{ m.}$$

Therefore, the height in meters is 300 m.

There are 1,000 meters in one kilometer. Meters can be converted to kilometers by using the factor 1 km/1000 m.

$$300 \text{ m} \times 1 \text{ km/1000 m} = .300 \text{ km.}$$

As such, there are .300 kilometers in 300 m.

There are 100 centimeters in 1 meter, thus meters can be converted to centimeters by multiplying by the factor 100 cm/1 m.

$$300 \text{ m} \times 100 \text{ cm/1 m} = 300 \times 10^2 \text{ cm.}$$

There are 30,000 centimeters in 300 m.

There are 1,000 millimeters in 1 meter; therefore,

meters can be converted to millimeters by the factor
1000 mm./1 m.

$$300 \text{ m} \times 1,000 \text{ mm/1 m} = 300 \times 10^3 \text{ mm.}$$

There are 300,000 millimeters in 300 meters.

• PROBLEM 2

The unaided eye can perceive objects which have a diameter of 0.1 mm. What is the diameter in inches?

Solution: From a standard table of conversion factors, one can find that 1 inch = 2.54 cm. Thus, cm can be converted to inches by multiplying by 1 inch/2.54 cm. Here, one is given the diameter in mm, which is .1 cm. Millimeters are converted to cm by multiplying the number of mm by .1 cm/1 mm. Solving for cm, you obtain:

$$0.1 \text{ mm} \times .1 \text{ cm/1 mm} = .01 \text{ cm.}$$

Solving for inches:

$$.01 \text{ cm} \times \frac{1 \text{ inch}}{2.54 \text{ cm}} = 3.94 \times 10^{-3} \text{ inches.}$$

AREA

• PROBLEM 3

One cubic millimeter of oil is spread on the surface of water so that the oil film has an area of 1 square meter. What is the thickness of the oil film in angstrom units?

Solution: Since one is asked to give the final thickness of the film in angstroms, it is useful to convert the other dimensions given to angstroms first. $1\text{\AA} = 10^{-10}\text{m} = 10^{-7}\text{mm}$. Therefore, $1\text{mm} = 1\text{\AA}/10^{-7} = 10^7\text{\AA}$.

Cubing both sides of this equation gives the number of cubic angstroms in 1 cubic millimeter.

$$(1\text{mm})^3 = (10^7\text{\AA})^3$$

$$1\text{mm}^3 = 10^{21}\text{\AA}^3 = \text{volume of oil.}$$

The final area of the film is given as 1m^2 . One

knows that $1 \text{ m} = 10^{10} \text{ \AA}$; therefore, $1 \text{ m}^2 = (10^{10} \text{ \AA})^2 = 10^{20} \text{ \AA}^2$. The volume is equal to the area of the film multiplied by the thickness. Thus, one can find the thickness of the film by dividing the volume by the area.

$$\text{thickness} = \frac{10^{21} \text{ \AA}^3}{10^{20} \text{ \AA}^2} = 10 \text{ \AA}.$$

• PROBLEM 4

How much area, in square meters, will one liter of paint cover if it is brushed out to a uniform thickness of 100 microns?

Solution: Because one is asked to give the final area in square meters, one should first convert the volume and thickness to meter units. One liter is equal to 1,000 cc. Since $1 \text{ m} = 100 \text{ cm}$, one can convert centimeters to meter units by cubing both sides of the equality:

$$1 \text{ m}^3 = (100 \text{ cm})^3$$

$$1 \text{ m}^3 = 1.0 \times 10^6 \text{ cc}$$

$$\frac{1 \text{ m}^3}{1.0 \times 10^6} = 1 \text{ cc}$$

$$10^{-6} \text{ m}^3 = 1 \text{ cc}$$

Therefore, 1000 cc or 1 liter is equal to $10^{-6} \text{ m}^3 \times 1,000$ or 10^{-3} m^3 . There are 10^6 microns in 1 m. Thus, 1 micron = 10^{-6} m and 100 microns = 10^{-4} m . The area of the film is equal to the volume divided by the thickness.

$$\text{Therefore, area} = \frac{10^{-3} \text{ m}^3}{10^{-4} \text{ m}} = 10 \text{ m}^2.$$

VOLUME

• PROBLEM 5

Determine the number of cubic centimeters in one cubic inch.

Solution: One meter equals 39.37 inches and, since there are 100 centimeters in 1 meter, there are 39.37 inches in 100 cm. Thus, 1 inch is equal to $100/39.37 \text{ cm}$.

$$1 \text{ inch} = \frac{100}{39.37} \text{ cm} = 2.54 \text{ cm}.$$

By cubing both sides of this equation, one can solve for the number of cubic centimeters in 1 cubic inch.

$$(1 \text{ inch})^3 = (2.54 \text{ cm})^3$$

$$1 \text{ inch}^3 = 16.4 \text{ cc}.$$

• PROBLEM 6

Calculate the number of liters in one cubic meter.

Solution: There are 1,000 milliliters (ml) or cubic centimeters (cc) in one liter. Thus, if one wishes to convert one cubic meter to liters, the cubic meter must be converted to cubic centimeters.

$$\begin{aligned} 1 \text{ meter} &= 100 \text{ centimeters} \\ (1 \text{ meter})^3 &= (100 \text{ centimeters})^3 \\ &= 1,000,000 \text{ centimeters}^3 \\ &= 1 \times 10^6 \text{ cubic centimeters} \end{aligned}$$

Cubic centimeters can be converted to liters by multiplying the number of cubic centimeters by the factor 1 liter/1,000 cubic centimeters.

$$\begin{aligned} 1 \times 10^6 \text{ cubic centimeters} \times 1 \text{ liter}/1,000 \text{ cubic centimeters} \\ = 1,000 \text{ liters.} \end{aligned}$$

There are 1,000 liters in one cubic meter.

• PROBLEM 7

What is the volume, in cubic centimeters, of a cube which is 150.0 mm along each edge?

Solution: There are 10 mm in 1 cm; therefore, millimeters can be converted to centimeters by multiplying the number of millimeters by 1 cm/10 mm.

$$\text{length of edge in cm} = 150 \text{ mm} \times 1 \text{ cm}/10 \text{ mm} = 15 \text{ cm}.$$

The volume of a cube is equal to the length of the side cubed.

$$\text{volume} = (15 \text{ cm})^3 = 3375 \text{ cc}.$$

• PROBLEM 8

What volume (in cc) is occupied by a block of wood of dimensions 25.0 m \times 10.0 cm \times 300 mm. All edges are 90° to one another.

Solution: Since all of the edges are 90° to one another, one knows that the block is a rectangular solid. The volume of a rectangle is equal to the length times the width times the height. If one wishes to find the volume in cubic centimeters, the lengths of all of the sides must be first expressed in centimeters.

There are 100 cm in 1 m; thus, to convert meters to centimeters, the number of meters must be multiplied by 100 cm/1 m.

$$25.0 \text{ m} \times 100 \text{ cm/1 m} = 2500 \text{ cm.}$$

There are 10 mm in 1 cm; thus, to convert millimeters to centimeters, multiply the number of millimeters by 1 cm/10 mm.

$$300 \text{ mm} \times 1 \text{ cm/10 mm} = 30 \text{ cm}$$

Solving for the volume:

$$\text{volume} = 2500 \text{ cm} \times 10.0 \text{ cm} \times 30 \text{ cm} = 7.50 \times 10^5 \text{ cc.}$$

• PROBLEM 9

What is the volume in liters of a rectangular tank which measures 2.0 m by 50 cm by 200 mm?

Solution: One liter is equal to 1000 cc; therefore, one should find the volume of the tank in cubic centimeters first and then convert to liters. This method is best for this problem because the sides of the tank are given in units which can quickly be converted to centimeters.

There are 100 cm in 1 m. Thus, 2 m is equal to 2.0 m \times 100 cm/1 m or 200 cm. There are 10 mm in 1 cm, therefore 200 mm is equal to 200 mm \times 1 cm/10 mm or 20 cm.

Solving for the volume of the tank in cubic centimeters:

$$\text{volume} = 200 \text{ cm} \times 50 \text{ cm} \times 20 \text{ cm} = 2.0 \times 10^5 \text{ cc.}$$

To convert from cubic centimeters to liters multiply by 1 liter/1000 cc.

$$\begin{aligned} \text{volume in liters} &= 2.0 \times 10^5 \text{ cc} \times 1 \text{ liter/1000 cc} \\ &= 200 \text{ liters.} \end{aligned}$$

• PROBLEM 10

A rectangular box is 5.00 in. wide, 8.00 in. long, and 6.0 in. deep. Calculate the volume in both cubic centimeters and in liters.

Solution: The volume of a solid is found by multiplying the height times the length times the width.

$$\text{volume} = (6.0 \text{ in}) \times (8.0 \text{ in}) \times (5.0 \text{ in}) = 240 \text{ in}^3.$$

From a standard conversion table, one finds that 1 inch = 2.54 cm. One finds the volume of cubic inches in cubic centimeters by cubing both sides of this equality.

$$1 \text{ inch} = 2.54 \text{ cm}$$

$$(1 \text{ inch})^3 = (2.54 \text{ cm})^3$$

$$1 \text{ inch}^3 = 16.4 \text{ cc.}$$

Thus, one can convert the volume of the rectangle from cubic inches to cubic centimeters by multiplying the number of cubic inches by the conversion factor, 16.4 cc/1 inch³.

$$\begin{aligned} \text{volume of rectangle} &= 240 \text{ in}^3 \times 16.4 \text{ cc/1 in}^3 \\ &= 3936 \text{ cc.} \end{aligned}$$

There are 1000 cc in 1 liter. Therefore, to convert from cubic centimeters to liters, multiply the number of cubic centimeters by 1 liter/1000 cc.

$$\begin{aligned} \text{volume in liters} &= 3936 \text{ cc} \times 1 \text{ liter/1000 cc} \\ &= 3.936 \text{ liters.} \end{aligned}$$

MASS

• PROBLEM 11

A student made three successive weighings of an object as follows: 9.17 g, 9.15 g, and 9.20 g. What is the average weight of the object in milligrams?

Solution: The average of a set of weights is found by adding together all of the weights and then dividing by the number of weighings used.

$$\text{avg. weight} = \frac{(9.17 \text{ g} + 9.15 \text{ g} + 9.20 \text{ g})}{3} = \frac{27.52 \text{ g}}{3} = 9.17 \text{ g}.$$

Now that the average weight in grams has been determined, convert it to milligrams using the conversion factor of 1,000 mg/g.

$$9.17 \text{ g} \times \frac{1000 \text{ mg}}{\text{g}} = 9170 \text{ mg}.$$

• PROBLEM 12

A silver dollar weighs about 0.943 ounces. Express this weight in grams, in kilograms, and in milligrams.

Solution: One ounce is equal to 28.35 g; thus, to convert from ounces to grams, one multiplies the number of ounces by the conversion factor, 28.35 g/1 ounce.

$$\begin{aligned} \text{no. of grams} &= 0.943 \text{ ounces} \times 28.35 \text{ g/1 ounce} \\ &= 26.73 \text{ g}. \end{aligned}$$

There are 1,000 g in 1 kg; therefore, to convert from grams to kilograms, one multiplies the number of grams by 1 kg/1,000 g.

$$\text{no. of kg} = 26.73 \text{ g} \times 1 \text{ kg/1000 g} = .02673 \text{ kg}.$$

There are 1,000 mg in one gram; thus, to convert from grams to milligrams, multiply the number of grams by the conversion factor, 1000 mg/1 g.

$$\text{no. of mg} = 26.73 \text{ g} \times 1000 \text{ mg/1 g} = 26,730 \text{ mg}.$$

• PROBLEM 13

It is estimated that 3×10^5 tons of sulfur dioxide, SO_2 , enters the atmosphere daily owing to the burning of coal and petroleum products. Assuming an even distribution of the sulfur dioxide throughout the earth's atmosphere (which is not the case), calculate in parts per million by weight the concentration of SO_2 added daily to the atmosphere. The weight of the atmosphere is 4.5×10^{15} tons. (On the average, about 40 days are required for the removal of the SO_2 by rain).

Solution: Here, one is asked to find the number of tons of SO_2 per 10^6 tons, i.e. per million, of atmosphere. This is done by using the following ratio: Let x = no. of tons of SO_2 per 10^6 tons of atmosphere.

$$\frac{3.0 \times 10^5 \text{ tons SO}_2}{4.5 \times 10^{15} \text{ tons atm}} = \frac{x}{10^6 \text{ tons atm}}$$

$$x = \frac{3.0 \times 10^5 \text{ tons SO}_2 \times 10^6 \text{ tons atm}}{4.5 \times 10^{15} \text{ tons atm}}$$

$$x = 6.67 \times 10^{-5} \text{ tons SO}_2 \text{ or } 6.67 \times 10^{-5} \text{ ppm SO}_2.$$

DENSITY

• PROBLEM 14

The density of alcohol is 0.8 g/ml. What is the weight of 50 ml. of alcohol?

Solution: Density is defined as weight per unit volume.

$$\text{density} = \frac{\text{weight}}{\text{volume}} = \frac{\text{g}}{\text{ml.}}$$

Thus, one can solve for the weight of the alcohol by multiplying the density by the volume.

$$\text{weight} = \text{density} \times \text{volume}$$

$$\text{weight} = 0.8 \text{ g/ml} \times 50 \text{ ml} = 40 \text{ g.}$$

• PROBLEM 15

Calculate the density of a block of wood which weighs 750 kg and has the dimensions 25 cm × 0.10 m × 50.0 m.

Solution: The density is a measure of weight per unit volume and is usually expressed in g/cc. Therefore, one must find the weight of this block in grams and the volume in cubic centimeters. The density is then found by dividing the weight by the volume.

$$1 \text{ kg} = 1,000 \text{ g; therefore, } 750 \text{ kg} = 750 \times 1,000 \text{ g}$$

$= 7.5 \times 10^5 \text{ g.}$ To find the volume in cubic centimeters, all of the dimensions must be converted to centimeters first.

$$1 \text{ m} = 100 \text{ cm; thus, } .10 \text{ m} = 10 \text{ cm and}$$

$$50.0 \text{ m} = 5,000 \text{ cm.}$$

$$\text{Volume} = 25 \text{ cm} \times 10 \text{ cm} \times 5,000 \text{ cm} = 1.25 \times 10^6 \text{ cc.}$$

Solving for the density:

$$\text{density} = \frac{\text{weight}}{\text{volume}} = \frac{7.5 \times 10^5 \text{ g}}{1.25 \times 10^6} = .60 \text{ g/cc.}$$

• PROBLEM 16

The density of concentrated sulfuric acid is 1.85 g/ml. What volume of the acid would weigh 74.0 g?

Solution: Density is defined as weight per unit volume.

$$\text{density} = \frac{\text{weight}}{\text{volume}} = \frac{\text{g}}{\text{ml.}}$$

$$\text{Therefore: volume} = \frac{\text{weight}}{\text{density}} .$$

Solving for the volume:

$$\text{volume} = \frac{74.0 \text{ g}}{1.85 \text{ g/ml}} = 40.0 \text{ ml.}$$

• PROBLEM 17

If 2.02 g of hydrogen gas occupies 22.4 liters, calculate the density of hydrogen gas in grams per milliliter.

Solution: Density = $\frac{\text{weight}}{\text{volume}} = \frac{\text{g}}{\text{ml.}}$

One is given the weight as 2.02 g but the volume is given in liters. Therefore, before calculating the density in g/ml., one must convert liters to milliliters. 1 liter = 1,000 ml; therefore, 22.4 l = 22,400 ml. Solving for the density:

$$\text{density} = \frac{2.02 \text{ g}}{22,400 \text{ ml}} = 9.0 \times 10^{-5} \text{ g/ml.}$$

• PROBLEM 18

One kilogram of metallic osmium, the "heaviest" substance known, occupies a volume of 44.5 cm³. Calculate the density of osmium in grams per cm³.

Solution: One is told that one kilogram of osmium occupies 44.5 cm³ and is then asked how many grams of osmium occupy one cm³. To find the density in grams per cm³, one kilo-

gram must be first converted to grams after which this number of grams is divided by 44.5 cm^3 , the volume that they occupy. There are 1000 grams in one kilogram. As such, kilograms are converted to grams by multiplying the number of kilograms present by the factor $1,000 \text{ g/1 kg}$.

$$1 \text{ kg} \times 1,000/1 \text{ kg} = 1,000 \text{ g}.$$

Therefore, $1,000 \text{ g}$ occupy 44.5 cm^3 . To find the number of grams present in 1 cm^3 , $1,000 \text{ g}$ is divided by 44.5 cm^3 .

$$1000 \text{ g}/44.5 \text{ cm}^3 = 22.5 \text{ g/cm}^3$$

The density of osmium is 22.5 g/cm^3 .

TEMPERATURE

• PROBLEM 19

Body temperature on the average is 98.6°F . What is this on (a) the Celsius scale and (b) the Kelvin scale?

Solution: (a) One converts $^\circ\text{F}$ to $^\circ\text{C}$ by using the following equation.

$$^\circ\text{C} = 5/9 (^\circ\text{F} - 32)$$

$$^\circ\text{C} = 5/9 (98.6 - 32)$$

$$= 5/9 (66.6) = 37.00^\circ\text{C}.$$

(b) $^\circ\text{C}$ can be converted to $^\circ\text{K}$ by adding 273.15 to the Celsius temperature.

$$^\circ\text{K} = ^\circ\text{C} + 273.15$$

$$^\circ\text{K} = 37.0 + 273.15 = 310.15^\circ\text{K}.$$

• PROBLEM 20

Liquid helium boils at 4°K . What is the boiling temperature on the Fahrenheit scale?

Solution: The temperature in Kelvin is the temperature in degrees Centigrade added to 273. In this problem, the boiling point is given in $^\circ\text{K}$. Hence, the temperature should be converted to $^\circ\text{C}$ and, then, to Fahrenheit using the relation

$$^\circ\text{F} = 9/5 ^\circ\text{C} + 32$$

The boiling point of helium can be converted to °C by subtracting 273 from the boiling point in °K.

$$^{\circ}\text{C} = ^{\circ}\text{K} - 273$$

$$^{\circ}\text{C} = 4^{\circ}\text{K} - 273 = -269^{\circ}\text{C}.$$

After the temperature is converted to the Centigrade scale, the temperature on the Fahrenheit scale can be determined.


$$^{\circ}\text{F} = 9/5 ^{\circ}\text{C} + 32$$

$$^{\circ}\text{F} = 9/5 (-269^{\circ}\text{C}) + 32 = -452^{\circ}\text{F}$$

The boiling point of helium on the Fahrenheit scale is -452°F.

• PROBLEM 21

The freezing point of silver is 960.8°C and the freezing point of gold is 1063.0°C. Convert these two readings to Kelvin (°K), Fahrenheit (°F), and Rankine (°R).



Solution: Kelvin: Temperatures measured in Celsius (°C) are converted to °K by adding 273.15 to the original measurement.

$$\text{freezing point of silver} = 960.8^{\circ}\text{C} + 273.15 = 1234^{\circ}\text{K}$$

$$\text{freezing point of gold} = 1063.0^{\circ}\text{C} + 273.15 = 1336.2^{\circ}\text{K}$$

Fahrenheit: °C are converted to °F by using the equation $^{\circ}\text{F} = 9/5 (^{\circ}\text{C}) + 32$.

$$\text{freezing point of silver} = 9/5 (960.8^{\circ}\text{C}) + 32 = 1761^{\circ}\text{F}$$

$$\text{freezing point of gold} = 9/5 (1063.0^{\circ}\text{C}) + 32 = 1945^{\circ}\text{F}$$

Rankine: The Rankine scale is an absolute scale

used by engineers. Its unit is the Fahrenheit degree. Absolute zero is equal to zero degrees Rankine. Convert $^{\circ}\text{K}$ to $^{\circ}\text{R}$ by using the equation

$$^{\circ}\text{R} = 9/5 (^{\circ}\text{K})$$

$$\text{freezing point of silver} = 9/5 (1234^{\circ}\text{K}) = 2221^{\circ}\text{R}$$

$$\text{freezing point of gold} = 9/5 (1336.2^{\circ}\text{K}) = 2405^{\circ}\text{R}.$$

CHAPTER 2



GASES

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 13 to 60 for step-by-step solutions to problems.

Gases, which represent one of the three important states of matter, differ remarkably from solids and liquids. Gases expand to uniformly fill a container and, unlike either solids or liquids, the volume is strongly dependent on the temperature and pressure. The pressure, volume, and temperature—called the state variables—are simply related for gases at low to moderate pressures by the ideal gas law or the ideal equation of state.

$$PV = nRT$$

2-1

The terms in this equation are defined as follows:

P = absolute pressure ($P = 0$ is a perfect vacuum)

V = volume

n = number of moles of gas*

T = absolute temperature ($T = 0$ where all molecular motion ceases) in kelvins

R = universal gas constant; it has the same value for all gases

In this chapter, the principles governing the pressure exerted by a column of static fluid are illustrated, and the Laws of Boyle, Charles, Gay-Lussac, and Avogadro (all early scientists who studied gas properties) are presented. These laws are combined into the ideal gas law, and a basic procedure for using the ideal gas law to calculate gas properties is outlined. It should be

*A mole is the mass numerically equal in grams to the molecular weight. It contains Avogadro's number, 6.02×10^{23} , of molecules which is the identical amount for any species. A lb. mole is also defined; it is the mass numerically equal in lbs. to the molecular weight. Note that a lb. mole does not contain Avogadro's number of molecules.

remembered that ideal gases, i.e., gases which obey the ideal gas law or ideal equation of state exactly, do not exist. But the behavior of real gases is often so closely approximated by the ideal gas law that it is very useful. A useful rule of thumb is that at pressures below 1 atmosphere and temperatures above the critical temperature (the temperature above which a gas cannot be liquified by an increase in pressure), real gases obey the ideal gas law extremely well.

Pressure

The pressure exerted by a column of fluid is equal to the weight of the fluid divided by the cross sectional area of the column. One simple equation, called the barometric equation, can be used to calculate this pressure.

$$P = \rho gh \text{ (barometric equation)} \quad 2-2$$

where ρ is the fluid density, g is the gravitational constant, and h is the height of the column. The principles or rules used to attack static pressure problems are as follows:

1. The pressure contribution of different static fluids in the same column, whether calculated from the barometric equation or obtained another way, can simply be added together to obtain the total pressure.

2. In a static fluid, all points at the same elevation have the same pressure.

In Problem 24, for example, the pressure at level h , is the same in every column. Therefore:

$$P_1 = P_2 + \rho gh_1 = P_3 + \rho g(h_3 + h_1) \quad 2-3$$

Since the level in column 2 is the same as the level open to the atmosphere, $P_2 = P_{\text{atm}}$. From the values of h_1 and h_3 , P_1 and P_3 are easily calculated and vice versa.

Gas Laws

The Laws of Boyle, Charles, Gay-Lussac, and Avogadro were developed from experimental observations.

Boyle's Law: The pressure of a quantity of a gas is inversely proportional to volume at constant temperature; $P \propto 1/V$ at constant T , n .

Charles' Law: The volume of a quantity of a gas is proportional to the absolute temperature at constant pressure; $V \propto T$ at constant P, n .

Gay-Lussac's Law: The pressure of a quantity of a gas is proportional to the temperature at constant volume; $P \propto T$ at constant V, n .

Avogadro's Law: At the same temperature and pressure, equal volumes of all gases contain the same number of molecules or moles; $V \propto n$ at constant T and P .

The four laws above can be combined into a single law — the ideal gas law or ideal equation of state.

Boyle's Law:	$P = k_1/V$	($T = \text{constant}$)
Charles' Law:	$V = k_2 T$	($P = \text{constant}$)
Gay-Lussac's Law:	$P = k_3 T$	($V = \text{constant}$)
Avogadro's Law:	$V = k_4 n$	(P and T constant)
Combined Law:		
$\frac{PV}{nT} = \text{constant}$ or $PV = nRT$		2-4

The value of R , the universal gas constant, depends on the units of pressure, volume, and temperature and is given in Table 1 for several common sets of units.

Table 1 - Value and Units of Universal Gas Constant (R)		
82.057	(atm cc)/(mol °K)	0.7302 (atm cubic feet)/(lb. mol R)
0.082	(atm liter)/(mol °K)	10.73 (psi cubic feet)/(lb. mol R)
62.361	(torr liter)/(mol °K)	1545 (lbf/ft ² cubic feet)/(lb. mol R)
8.31	(Pascal m ³)/(mol °K) or joule/(mol °K)	or (ft lbf)/(lb. mol R)

In solving ideal gas law problems, it is not necessary to remember the separate Laws of Boyle, Charles, Gay-Lussac, and Avogadro. They are all special cases of the ideal gas law which can be expressed in the following forms:

$$\frac{P_1 V_1}{n_1 T_1} = \frac{P_2 V_2}{n_2 T_2} \text{ or } PV = nRT \quad 2-5$$

The best procedure for attacking gas law problems is to tabulate, in one column, what is known and, in a second, what is to be calculated. Then select the form of the gas law that is the most convenient. Observe that the form on the left above permits variables which do not change (e.g., T_1 and T_2 in a constant temperature problem) to be canceled from both sides of the equation. Let us examine Problem 54 in the text from this perspective.

Known	To be calculated
-------	------------------

P_1, V_1, T_1	V_2
-----------------	-------

P_2, T_2	
------------	--

$n_1 = n_2 = \text{constant}$	
-------------------------------	--

Remember to convert Celsius to Kelvin.

$$V_2 = \frac{P_1 V_1 n_2 T_2}{P_2 n_1 T_1} = \frac{(600 \text{ torr})(350 \text{ ml})(298 \text{ K})}{(25 \text{ torr})(873 \text{ K})} = 24,600 \text{ ml}$$

Note that the unit for each quantity is written, and the unit of the answer is calculated along with its numerical value as discussed in Chapter 1.

Let us illustrate this procedure again using Problem 56 in the text.

Known	To be calculated
-------	------------------

P_{O_2}, V_{O_2}	$n = n_{O_2} + n_{N_2}$
--------------------	-------------------------

P_{N_2}, V_{N_2}	$n_{N_2} = P_{N_2} V_{N_2} / RT$
--------------------	----------------------------------

$T = \text{constant}$	$n_{O_2} = P_{O_2} V_{O_2} / RT$
-----------------------	----------------------------------

	$V_f = V_{O_2} + V_{N_2} = 3 \text{ liters} + 2 \text{ liters} = 5 \text{ liters}$
--	--

	P_f
--	-------

From the ideal gas law:

$$P_f = n_f(RT) / V_f = (n_{O_2} + n_{N_2})(RT) / V_f$$

$$= (P_{O_2} V_{O_2} / RT + P_{N_2} V_{N_2} / RT) RT / 5 \text{ liters}$$


$$= [(195 \text{ torr})(3 \text{ liters}) + (530 \text{ torr})(2 \text{ liters})] / 5 \text{ liters} = 329 \text{ torr}$$

Step-by-Step Solutions to Problems in this Chapter, "Gases"

PRESSURE

• PROBLEM 22

Given the setup in the figure, what would be the pressure of the gas (in atm) if P_{atm} is 745 Torr and P_{liq} is the equivalent of a mercury column 3.0 cm high?



Solution: A manometer is used to measure the pressure of a trapped sample of gas. If the right hand tube is open to the atmosphere, the pressure which is exerted in the right-hand surface is atmospheric pressure, P_{atm} . If the liquid level is the same in both arms of the tube, the pressures must be equal; otherwise, there would be a flow of liquid from one arm to the other. At the level indicated by the dashed line in the figure, the pressure in the left arm

is equal to the pressure of the trapped gas, P_{gas} , plus the pressure of the column of liquid above the dashed line, P_{liq} . One can write

$$P_{\text{gas}} = P_{\text{atm}} - P_{\text{liq}}$$

Here, one is given that P_{atm} is 745 Torr and that P_{liq} is equivalent to a mercury column 3.0 cm high. One wishes to find P_{gas} in atm, thus, one must convert 745 Torr to atm and find the P_{liq} . There are 760 Torr in 1 atm, which means

$$\text{no. of atm in } P_{\text{atm}} = \frac{745 \text{ Torr}}{760 \text{ Torr/1 atm}} = .98 \text{ atm}$$

One atmosphere pressure supports 76 cm of mercury, thus 3 cm of mercury supports $3/76$ atm. Therefore,
 $P_{\text{liq}} = 3 \text{ cm}/76 \text{ cm/atm} = .039 \text{ atm}$.

Solving for P_{gas}

$$P_{\text{gas}} = .980 \text{ atm} - .039 \text{ atm} = .941 \text{ atm}.$$


• PROBLEM 23

Consider the manometer, illustrated below, first constructed by Robert Boyle. When $h = 40 \text{ mm}$, what is the pressure of the gas trapped in the volume, V_{gas} . The temperature is constant, and atmospheric pressure is $P_{\text{atm}} = 1 \text{ atm}$.

Solution: We do not need to know any gas law to solve this problem. All we must realize is that the pressure exerted on the gas, P_{total} , is equal to the sum of the pressure exerted by the mercury, P_{Hg} , and the pressure exerted by the air, P_{atm} . Since $1 \text{ mm Hg} = 1 \text{ torr}$ and $1 \text{ atm} = 760 \text{ torr}$,


$$P_{\text{Hg}} = 40 \text{ mm Hg} = 40 \text{ torr and } P_{\text{atm}} = 1 \text{ atm} = 760 \text{ torr}.$$

$$\text{Then } P_{\text{total}} = P_{\text{Hg}} + P_{\text{atm}} = 40 \text{ torr} + 760 \text{ torr} = 800 \text{ torr}.$$



• PROBLEM 24

Consider gases confined by a liquid, as shown in the diagram below. Find an expression for the pressures P_1 , P_2 , and P_3 in terms of the density of the liquid, ρ (g/ml), the heights h_1 and h_3 (mm), and the barometric pressure P_{atm} (mm Hg).



Solution: The device shown above is called a eudiometer. It is used to compare the pressures of several gases. The pressure of the confined gas is equal to the barometric pressure plus the pressure needed to depress the column of liquid (for P_1) or to the barometric pressure minus the pressure needed to support the column of liquid (for P_3). The pressure of the liquid column is given by

$$\text{pressure of liquid} = \frac{\text{height or depth of column (mm)}}{\text{column (mm)}} \times \rho \text{ (g/ml)} (\text{ml/cm}^3) (\text{cm/10mm})$$

Hence:

P_1 = barometric pressure + pressure needed to depress liquid

$$= P_{\text{atm}} + h_1 \times \rho \left(\frac{\text{ml}}{\text{cm}^3} \right) \text{ cm/10mm}$$

$$= P_{\text{atm}} + 0.1 h_1 \left(\frac{\text{g}}{\text{cm}^2} \right) \rho$$

P_2 = barometric pressure + pressure needed to depress liquid (or - pressure needed to elevate liquid)

$$= P_{\text{atm}} + 0 \times \rho \times \text{ml/cm}^3 \times \text{cm/10mm}$$

$$= P_{\text{atm}}$$

P_3 = barometric pressure - pressure needed to elevate liquid

$$= P_{\text{atm}} - h_3 \times \rho \times \text{ml/cm}^3 \times \text{cm/10mm}$$


$$= P_{\text{atm}} - 0.1 h_3 \left(\frac{\text{g}}{\text{cm}^2} \right) \rho$$

• PROBLEM 25

(a) A diver descends to a depth of 15.0 m in pure water (density 1.00 g/cm^3). The barometric pressure is 1.02 standard atmospheres. What is the total pressure on the diver, expressed in atmospheres? (b) If, at the same barometric pressure, the water were the Dead Sea (1.20 g/cm^3), what would the total pressure be?

Solution: Pressure is defined as force per unit area. Atmospheric is measured by using a barometer (usually a mercury barometer; see figure). It is constructed by inverting a tube longer than 76 cm filled with mercury into a dish of mercury. The atmosphere will support only that height of mercury which exerts an equivalent pressure; any excess mercury will fall into the reservoir and leave a space with zero air pressure above it.

The pressure exerted on the diver from above is equal to the sum of the pressure exerted by the sea water and the pressure exerted by the atmosphere. One standard atmosphere equals the pressure exerted by exactly 76 cm (= exactly 760 mm) of mercury at 0°C



Mercury Barometer

(density Hg = 13.5951 g/cm^3) and at standard gravity, 980.665 cm/s^2 . Thus, 1 standard atm equals $13.5951 \text{ g/cm}^3 \times 76 \text{ cm (exactly)} \times 980.665 \text{ cm/s}^2 = 1.01325 \times 10^6 \text{ dynes/cm}^2$. The pressure exerted by the water is found similarly:

pressure of water = density \times height \times standard gravity

$$= 1.00 \text{ g/cm}^3 \times 1.5 \times 10^3 \text{ cm} \times 980.665 \text{ cm/s}^2$$

$$= 1.47 \times 10^6 \text{ dynes/cm}^2.$$

However, the problem states that atmospheric pressure is 1.02 standard atmospheres and, therefore, equals $(1.02)(1.01325 \times 10^6 \text{ dynes/cm}^2) = 1.03 \times 10^6 \text{ dynes/cm}^2$. To this atmospheric pressure, the pressure of the water is added to yield a total pressure of

$$\begin{aligned} (1.03 \times 10^6 \text{ dynes/cm}^2) + (1.47 \times 10^6 \text{ dynes/cm}^2) \\ = 2.50 \times 10^6 \text{ dynes/cm}^2. \end{aligned}$$

This answer expressed in atmospheres gives

$$\frac{2.50 \times 10^6 \text{ dynes/cm}^2}{1.01 \times 10^6 \text{ dynes/cm}^2/\text{atm}} = 2.48 \text{ atm.}$$

(b) This part is very similar to part (a). The total pressure exerted is the pressure of the water plus the pressure of the atmosphere. The pressure of the atmosphere from part (a) is $1.03 \times 10^6 \text{ dynes/cm}^2$. The pressure of the water must be calculated:

pressure of water = density \times height \times standard gravity

$$= 1.20 \text{ g/cm}^3 \times 1.5 \times 10^3 \text{ cm} \times 980.665 \text{ cm/s}^2$$

$$= 1.77 \times 10^6 \text{ dynes/cm}^2.$$

The total pressure is $1.77 \times 10^6 \text{ dynes/cm}^2 + 1.03 \times 10^6 \text{ dynes/cm}^2$ equals $2.80 \times 10^6 \text{ dynes/cm}^2$.

This answer expressed in atmospheres is

$$\frac{2.80 \times 10^6 \text{ dynes/cm}^2}{1.01 \times 10^6 (\text{dynes/cm}^2)/\text{atm}} = 2.77 \text{ atm.}$$

• PROBLEM 26

How many full strokes of a bicycle pump (chamber 4.0 cm diameter and 40.0 cm long) would you need to make in order to pump up an automobile tire from a gauge pressure of zero to 24 pounds per square inch (psi) (1.63 atm.)? Assume temperature stays constant at 25°C and atmospheric pressure is one atmosphere. Note, that gauge pressure measures only the excess over atmospheric pressure. A typical tire volume is about 25 liters.

Solution: One atmosphere equals 14.7 psi, therefore, the amount of pressure needed to fill the tire is $24 + 14.7$ psi or 38.7 psi. Converting back to atm:

$$\text{atm. contained in inflated tire} = 38.7 \text{ psi} \times 1 \text{ atm}/14.7 \text{ psi} = 2.63 \text{ atm.}$$

When the tire is deflated the pressure is 1 atm and the volume is 25 l. Each atm of pressure occupies 25 l. Therefore, the volume of the tire when inflated is $25 \text{ l} \times 2.63 \text{ atm}$.

$$\text{Volume of inflated tire} = 25 \text{ l} \times 2.63 = 65.75 \text{ l}$$

Since there is 25 l present in the tire before inflation, the volume that the pump must contribute is $65.75 - 25 \text{ l}$ or 40.75 l.

The volume of air forced into the tire at each stroke of the pump is equal to the volume of the pump.

$$\text{Volume of pump} = \pi r^2 h = \pi \times (2 \text{ cm})^2 \times 40 \text{ cm} = 503 \text{ cm}^3/\text{stroke}$$

There are 1000 cm³ in 1 liter, therefore 40750 cm³ of air must be pumped into the tire. If 503 cm³ is pumped in per stroke the number of strokes necessary to fill the tire is $\frac{40750 \text{ cm}^3}{503 \text{ cm}^3/\text{stroke}}$.


$$\text{No. of strokes} = \frac{40750 \text{ cm}^3}{503 \text{ cm}^3/\text{stroke}} = 81 \text{ strokes.}$$

BOYLE'S LAW, CHARLES' LAW, LAW OF GAY-LUSSAC

• PROBLEM 27

100 ml. of gas are enclosed in a cylinder under a pressure of 760 Torr. What would the volume of the same gas be at a pressure of 1520 Torr?

P = Pressure



If the pressure on a gas is doubled (at a constant temperature), the volume is halved (Boyle's Law).

Solution: Since this problem deals with the pressure and volume of a gas at a constant temperature, Boyle's law can be used. Boyle's law states that the volume, V , of a given mass of gas, at constant temperature, varies inversely with the pressure, P . It can be stated as

$$V = k \times 1/P,$$

where k is a constant.

Hence, $k = PV$

For a particular system, at constant temperature, k is constant. Therefore, if either the pressure or the volume is changed, the other must adjust accordingly.

Here, $P = 760$ Torr and $V = 100$ ml so

$$k = 760 \times 100 = 76000 \text{ Torr-ml.}$$

If P is doubled to 1520 Torr, then

$$k = 76000 \text{ Torr-ml} = 1520 \text{ Torr} \times V$$

$$V = \frac{76000 \text{ Torr-ml}}{1520 \text{ Torr}} = 50 \text{ ml}$$

Since k is a constant for a given system, another form of Boyle's Law can be expressed as

$$P_1 V_1 = P_2 V_2$$

This says that the pressure of the original system times the volume of the original system is equal to the new pressure times the new volume. Here,

$$760 \text{ Torr} \times 100 \text{ ml} = 1520 \text{ Torr} \times V_2$$

$$50 \text{ ml} = V_2.$$

• PROBLEM 28

What pressure is required to compress 5 liters of gas at 1 atm. pressure to 1 liter at a constant temperature?

Solution: In solving this problem, one uses Boyle's Law: The volume of a given mass of gas at constant temperature varies inversely with the pressure. This means that, for a given gas, the pressure and the volume are proportional; at a constant temperature, and their product equals a constant.

$$P \times V = K$$

where P is the pressure, V is the volume and K is a constant. From this one can propose the following equation

$$P_1 V_1 = P_2 V_2,$$

where P_1 is the original pressure, V_1 is the original volume, P_2 is the new pressure and V_2 is the new volume.

In this problem, one is asked to find the new pressure and is given the original pressure and volume and the new volume.

$$P_1 V_1 = P_2 V_2$$

$$1 \text{ atm} \times 5 \text{ liters} = P_2 \times 1 \text{ liter}$$

$$\frac{1 \text{ atm} \times 5 \text{ liters}}{1 \text{ liter}} = P_2$$

$$5 \text{ atm.} = P_2.$$

$$P_1 = 1 \text{ atm.}$$

$$V_1 = 5 \text{ liters}$$

$$P_2 = ?$$

$$V_2 = 1 \text{ liter}$$

• PROBLEM 29

A mass of gas is under a pressure 760 mm Hg and occupies a volume of 525 ml. If the pressure were doubled, what volume would the gas now occupy? Assume the temperature is constant.

Solution: This question can be answered by using Boyle's Law. This law states that, under constant temperature, the pressure is inversely proportional to the volume of a gas. In other words, $PV = K$, where P = pressure, V = volume and K = constant. Therefore, for a given gas at constant temperature, $P_1 V_1 = P_2 V_2$, where P_1 and V_1 are

the initial pressure and volume, and P_2 and V_2 are the final values.

In this problem $P_1 = 760$ mm Hg, $V_1 = .525$ liter (525 ml) and $P_2 = 1520$ mmHg. The unknown, V_2 , can be determined by substituting these values into $P_1V_1 = P_2V_2$.

$(760)(.525) = 1520 (V_2)$ Solving for
 V_2 , you obtain $V_2 = .2625 \text{ l} = 262.5 \text{ ml}$.

• PROBLEM 30

A gaseous sample of neon, maintained at constant temperature, occupies 500 ml at 2.00 atm. Calculate the volume when the pressure is changed to each of the following: (a) 4.00 atm; (b) 760 torr; (c) 1.8×10^{-3} torr.

Solution: We need a relationship between volume and pressure. Such a relationship is provided by Boyle's Law, which states that the product of pressure P and volume V of an ideal gas is a constant, k , or $PV = k$.

We must first determine k for the neon sample. We are given a value of P corresponding to a given value of V .

Therefore, $k = PV = 2.00 \text{ atm} \times 500 \text{ ml} = 2.00 \text{ atm} \times 0.500 \text{ l} = 1.00 \text{ l-atm}$. Now that k is determined, we can obtain the value of V corresponding to the given values of P by using the formula $V = k/P$.

$$(a) \quad P = 4.00 \text{ atm:} \quad V = \frac{k}{P} = \frac{1.00 \text{ l-atm.}}{4.00 \text{ atm}} = 0.25 \text{ l}$$

$$(b) \quad P = 760 \text{ torr} = 760 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 1 \text{ atm} :$$

$$V = \frac{k}{P} = \frac{1.00 \text{ l-atm.}}{1 \text{ atm}} = 1.00 \text{ l}$$

$$(c) \quad P = 1.8 \times 10^{-3} \text{ torr} = 1.8 \times 10^{-3} \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} \\ = 2.3 \times 10^{-6} \text{ atm:}$$

$$V = \frac{k}{P} = \frac{1.00 \text{ l-atm.}}{2.3 \times 10^{-6} \text{ atm}} = 4.35 \times 10^5 \text{ l.}$$

• PROBLEM 31

A sample of gaseous krypton, maintained at constant pressure, is found to have a volume of 10.5 l at 25°C. If the system is heated to 250°C, what is the resulting volume?

Solution: We need a relationship between volume and temperature. Such a relationship is provided by Charles' Law, which states that volume V and absolute temperature T are proportional, or, as an equality, $V = kT$, where k is a constant.

We can determine k for our system from the initial volume and temperature. Thus,

$$k = \frac{V}{T} = \frac{10.5 \text{ l}}{298.15^\circ\text{K}} = \frac{10.5 \text{ l}}{298.15^\circ\text{K}} = 0.0352 \text{ l}^\circ\text{K}^{-1}$$

The volume corresponding to a temperature of 250°C, which is 523.15°K, is $V = kT = 0.0352 \text{ l}^\circ\text{K}^{-1} \times 523.15^\circ\text{K} = 18.41 \text{ l}$.

• PROBLEM 32

A certain gas occupies a volume of 100 ml at a temperature of 20°C. What will its volume be at 10°C, if the pressure remains constant?

Solution: In a gaseous system, when the volume is changed by increasing the temperature, keeping the pressure constant, Charles' Law can be used to determine the new volume. Charles' Law states that, at a constant pressure, the volume of a given mass of gas is directly proportional to the absolute temperature. Charles' Law may also be written

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

where V_1 is the volume at the original temperature T_1 and V_2 is the volume at the new temperature T_2 .

To use Charles' Law, the temperature must be expressed on the absolute scale. The absolute temperature is calculated by adding 273 to the temperature in degrees Centigrade. In this problem, the centigrade temperatures are given and one must convert them to the absolute scale.

$$T_1 = 20^\circ\text{C} + 273 = 293^\circ\text{K}$$

$$T_2 = 10^\circ\text{C} + 273 = 283^\circ\text{K}$$

Using Charles' Law,

$$V_1 = 100 \text{ ml}$$

$$T_1 = 293^\circ\text{K}$$

$$T_2 = 283^\circ\text{K}$$

$$V_2 = ?$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \qquad V_2 = \frac{V_1 T_2}{T_1}$$

$$V_2 = \frac{(100 \text{ ml})(283^\circ\text{K})}{293^\circ\text{K}}$$

$$V_2 = 96.6 \text{ ml.}$$

• PROBLEM 33

Assume that one cubic foot of air near a thermonuclear explosion is heated from 0°C to $546,000^\circ\text{C}$. To what volume does the air expand?

Solution: Charles' Law ($V_1/T_1 = k$, where V_1 is the initial volume, T_1 the initial absolute temperature, and k is a constant) states that the volume is directly proportional to the temperature.

$$V \propto T$$

\propto means 'is proportional to.'

The absolute temperature can be found by adding 273 to the temperature in $^\circ\text{C}$.

$$T_1 = 0 + 273 = 273$$

$$T_2 = 546,000 + 273 = 546,273$$

Using the Charles' Law, one can set up the following ratio

$$\frac{V_1}{T_1} = \frac{V_2}{T_2},$$

where V_1 is the original volume, V_2 is the final volume, T_1 is the original temperature, and T_2 is the final temperature. Using this ratio, one can determine the final volume.

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

Substituting, one obtains

$$\frac{1 \text{ ft}^3}{273} = \frac{V_2}{546,273}$$

$$V_2 = \frac{546,273}{273} (1 \text{ ft}^3) = 2001 \text{ ft}^3.$$

• PROBLEM 34

The volume of a sample of gaseous argon maintained at constant pressure was studied as a function of temperature and the following data were obtained:

Temperature, T ($^{\circ}\text{K}$)	Volume, V (l)
250	0.005
300	0.006
350	0.007

Calculate and confirm the Charles' Law constant for this system. Determine the temperature corresponding to a volume of 22.4 l.

Solution: Charles' law states that the volume and absolute temperature of an ideal gas are directly proportional, i.e. $V = kT$, k is a constant. To determine the Charles' law constant k , we can use any one of the three sets of temperature-volume measurement. Choosing the first of these, we obtain

$$k = \frac{V}{T} = \frac{0.005 \text{ l}}{250^{\circ}\text{K}} = 2.00 \times 10^{-5} \text{ l}^{\circ}\text{K}^{-1}. \quad \text{This value}$$

is confirmed by showing that the other two sets of data give the same value for k :

$$k = \frac{V}{T} = \frac{0.006 \text{ l}}{300^{\circ}\text{K}} = 2.00 \times 10^{-5} \text{ l}^{\circ}\text{K}^{-1} \quad \text{and}$$

$$k = \frac{V}{T} = \frac{0.007 \text{ l}}{350^{\circ}\text{K}} = 2.00 \times 10^{-5} \text{ l}^{\circ}\text{K}^{-1}.$$


To determine the temperature corresponding to 22.4 l, we solve Charles' law for T , obtaining

$$T = \frac{V}{k} = \frac{22.4 \text{ l}}{2.00 \times 10^{-5} \text{ l}^{\circ}\text{K}^{-1}} = 1.12 \times 10^6 \text{ }^{\circ}\text{K}$$

or about a million degrees Kelvin.

• PROBLEM 35

Consider the gas thermometer illustrated below. At 0°C , the volume of the gas is 1.25 liters. Assuming that the cross-sectional area of the graduated arm is 1 cm^2 , what is the distance (in cm.) from the 0°C reading and a reading at 35°C ?



Solution: Since volume, V , and absolute temperature, T , are the only two variables being considered, we can approach this problem by means of Charles' Law, $V = kT$, where k is the Charles' law constant, to be determined for our system.

We can determine k by using the initial values of 0°C ($0^{\circ}\text{C} = 273.15^{\circ}\text{K}$) and 1.25 liters ($= 1250\text{ ml} = 1250\text{ cm}^3$). Then,

$$k = \frac{V}{T} = \frac{1250\text{ cm}^3}{273.15^{\circ}\text{K}} = 4.576\text{ cm}^3\text{-}^{\circ}\text{K}^{-1}$$

The volume at 35°C (308.15°K) can now be determined:
 $V = kT = 4.576\text{ cm}^3\text{-}^{\circ}\text{K}^{-1} \times 308.15^{\circ}\text{K} = 1410.1\text{ cm}^3 = 1.410\text{ l.}$
 The difference between this volume and the initial volume is $1410.1\text{ cm}^3 - 1250\text{ cm}^3 = 160.1\text{ cm}^3$. This volume of liquid will be displaced in the graduated arm.

The difference in height of the liquid in the graduated arm, can now be calculated from the volume displaced and the cross-sectional area:

$$\text{height} = \frac{\text{volume}}{\text{area}} = \frac{160.1\text{ cm}^3}{1\text{ cm}^2} = 160.1\text{ cm.}$$

The air in a tank has a pressure of 640 mm of Hg at 23°C. When placed in sunlight the temperature rose to 48°C. What was the pressure in the tank?

Solution: The law of Gay-Lussac deals with the relationship existing between pressure and the absolute temperature ($^{\circ}\text{C} + 273^{\circ}$), for a given mass of gas at constant volume. The relationship is expressed in the law of Gay-Lussac: volume constant, the pressure exerted by a given mass of gas varies directly with the absolute temperature. That is:

$P \propto T$ (volume and mass of gas constant).

The variation that exists between pressure and temperature at different states can be expressed as

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

where P_1 = pressure of original state, T_1 = absolute temperature of original state, P_2 = pressure of final state, and T_2 = absolute temperature of final state.

Thus this problem is solved by substituting the given values into Gay-Lussac's Law.

$$P_1 = 640 \text{ mm Hg} \qquad T_1 = 23^{\circ}\text{C} + 273^{\circ} = 296^{\circ}\text{K}$$

$$P_2 = ? \qquad T_2 = 48^{\circ}\text{C} + 273^{\circ} = 321^{\circ}\text{K}$$

Substituting and solving,

$$\frac{640 \text{ mm}}{296^{\circ}\text{K}} = \frac{P_2}{321^{\circ}\text{K}}$$

$$P_2 = 640 \text{ mm} \times \frac{321^{\circ}\text{K}}{296^{\circ}\text{K}}$$

$$= 694 \text{ mm of Hg.}$$

A sealed glass bulb contained helium at a pressure of 750 mm Hg and 27°C. The bulb was packed in dry ice at - 73°C. What was the resultant pressure of the helium?

Solution: The only parameters mentioned in this problem are pressure (P) and temperature (T). One is given the initial pressure and temperature and the final temperature. One is asked to determine the pressure at - 73°C. The

Law of Gay-Lussac relates temperature and pressure. It can be stated mathematically

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

where P_1 is the initial pressure, T_1 the initial absolute temperature, P_2 the final pressure and T_2 the final absolute temperature.

The temperature in $^{\circ}\text{C}$ is converted to $^{\circ}\text{K}$ by adding 273.

$$T_1 = 27^{\circ}\text{C} + 273 = 300^{\circ}\text{K}$$

$$T_2 = -73^{\circ}\text{C} + 273 = 200^{\circ}\text{K}$$

Solving for P_2

$$\frac{750 \text{ mm Hg}}{300^{\circ}\text{K}} = \frac{P_2}{200^{\circ}\text{K}}$$

$$P_2 = \frac{(750 \text{ mm Hg})(200^{\circ}\text{K})}{300^{\circ}\text{K}} = 500 \text{ mm Hg.}$$

COMBINED GAS LAWS

• PROBLEM 38

A gas occupies a volume of 1.0 liter at a temperature of 27°C and 500 Torr pressure. Calculate the volume of the gas if the temperature is changed to 60°C and the pressure to 700 Torr.

Solution: For a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. This combined law can also be written:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 is the original pressure, V_1 is the original volume, T_1 is the original absolute temperature, P_2 is the new pressure, V_2 is the new volume, and T_2 is the new absolute temperature.

In this problem, one is given the original pressure, temperature, and volume and the new temperature

and pressure. One is asked to calculate the new volume. The temperatures are given in °C; they must be converted to the absolute scale before using the combined law. This can be done by adding 273 to the temperature in °C.

Converting the temperature:

$$T_1 = 27^\circ + 273 = 300^\circ\text{K}$$

$$T_2 = 60^\circ + 273 = 333^\circ\text{K}$$

Using the combined law:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_1 = 500 \text{ Torr}$$

$$P_2 = 700 \text{ Torr}$$

$$V_1 = 1.0 \text{ liter}$$

$$V_2 = ?$$

$$T_1 = 300^\circ\text{K}$$

$$T_2 = 333^\circ\text{K}$$

$$\frac{(500 \text{ Torr})(1.0 \text{ liter})}{(300^\circ\text{K})} = \frac{(700 \text{ Torr})(V_2)}{(333^\circ\text{K})}$$

$$V_2 = \frac{(500 \text{ Torr})(1.0 \text{ liter})(333^\circ\text{K})}{(300^\circ\text{K})(700 \text{ Torr})}$$

$$V_2 = 0.79 \text{ liter.}$$

• PROBLEM 39

Calculate the pressure required to compress 2 liters of a gas at 700 mm pressure and 20°C into a container of 0.1 liter capacity at a temperature of - 150°C.

Solution: One is dealing with changing volumes, pressures and temperatures of a gas. Therefore, this problem can be solved using the combined gas law. It states that as the pressure increases, the volume decreases and that as the temperature increases, the volume increases. These factors are related by the equation

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 , V_1 and T_1 are the initial pressure, volume and temperature and P_2 , V_2 , and T_2 are the final values.

For any problem dealing with gases, the first step always involves converting all of the temperatures to the degree Kelvin scale by the equation

$$^\circ\text{K} = ^\circ\text{C} + 273$$

For this question

$$T_1 = 20^\circ\text{C} = 20 + 273 = 293^\circ\text{K}$$

$$T_2 = -150^\circ\text{C} = -150 + 273 = 123^\circ\text{K}.$$

This seems to indicate that the pressure would decrease. But one is also told that the volume decreases, which would have the effect of increasing the pressure. Therefore, one cannot predict the final change in volume.

For the sake of clarity, set up a table as given below.

$$P_1 = 700 \text{ mm}$$

$$P_2 = ?$$

$$V_1 = 2 \text{ liters}$$

$$V_2 = 0.1 \text{ liter}$$

$$T_1 = 293^\circ\text{K}$$

$$T_2 = 123^\circ\text{K}$$

Since one is given 5 of the 6 values, it is possible to use the combined gas law equation to determine P_2

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_2 = \frac{T_2 V_1 P_1}{T_1 V_2}$$

$$= \frac{123^\circ\text{K} (2 \text{ liters}) (700 \text{ mm})}{293^\circ\text{K} (0.1 \text{ liter})}$$

$$= 5877 \text{ mm}.$$

• PROBLEM 40

750 ml of gas at 300 torr pressure and 50°C is heated until the volume of gas is 2000 ml at a pressure of 700 torr. What is the final temperature of the gas?

Solution: Here, one is given a gaseous system involving pressure, volume and temperature, where two of the variables are changed in going from the original system to the final system. This indicates that the combined gas law should be used. It can be stated: For a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. This gas law can also be stated

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 is the original pressure, V_1 is the original volume, T_1 is the original absolute temperature, P_2 is the final

pressure, V_2 is the final volume, and T_2 is the final absolute temperature.

In this problem, you are given the original pressure, volume and temperature and the final pressure and volume. You are asked to find the final temperature. The temperature in $^{\circ}\text{C}$ must be converted to the absolute scale before using the combined law. This can be done by adding 273 to the temperature in $^{\circ}\text{C}$.

Converting the temperature:

$$T_1 = 50 + 273 = 323^{\circ}\text{K}$$

Using the combined law:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_1 = 300 \text{ torr}$$

$$P_2 = 700 \text{ torr}$$

$$V_1 = 750 \text{ ml}$$

$$V_2 = 2000 \text{ ml.}$$

$$T_1 = 323^{\circ}\text{K}$$

$$T_2 = ?$$

$$\frac{(300 \text{ torr})(750 \text{ ml})}{323^{\circ}\text{K}} = \frac{(700 \text{ torr})(2000 \text{ ml})}{T_2}$$

$$T_2 = \frac{(700 \text{ torr})(2000 \text{ ml})(323^{\circ}\text{K})}{(300 \text{ torr})(750 \text{ ml})}$$

$$T_2 = 2010^{\circ}\text{K}$$

Convert T_2 to centigrade by subtracting 273 from it.

$$2010 - 273 = 1737^{\circ}\text{C.}$$

• PROBLEM 41

500 liters of a gas at 27°C and 700 torr would occupy what volume at STP?

Solution: STP (Standard Temperature and Pressure) is defined as being 0°C and 760 torr, thus, in this problem, one is asked to find the new volume of a gas when the temperature and pressure are changed. One refers to the combined gas law in such a case. This law can be stated: For a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. Stated algebraically

$$\frac{PV}{T} = K$$

where P is the pressure, V is the volume, T is the absolute temperature, and K is a constant. This means that if two of the variables are changed, the third changes so that the relation $PV/T = K$ remains true. This means that one can now state that

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 is the original pressure, V_1 is the original volume, T_1 is the original temperature, P_2 is the new pressure, V_2 is the new volume, and T_2 is the new absolute temperature. In this problem, one is given the temperature on the Celsius scale. It must be converted to the absolute scale before using the combined gas law. This can be done by adding 273 to the temperature in $^{\circ}\text{C}$.

$$T_1 = 27 + 273 = 300^{\circ}\text{K}$$

$$T_2 = 0 + 273 = 273^{\circ}\text{K}$$

One knows P_1 , V_1 , T_1 , P_2 , and T_2 . One is asked to find V_2 .

Using the Combined Gas Laws:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_1 = 700 \text{ torr}$$

$$V_1 = 500 \text{ liters}$$

$$T_1 = 300^{\circ}\text{K}$$

$$P_2 = 760 \text{ torr}$$

$$V_2 = ?$$

$$T_2 = 273^{\circ}\text{K}$$

$$\frac{700 \text{ torr} \times 500 \text{ liters}}{300^{\circ}\text{K}} = \frac{760 \text{ torr} \times V_2}{273^{\circ}\text{K}}$$

$$V_2 = \frac{700 \text{ torr} \times 500 \text{ liters} \times 273^{\circ}\text{K}}{760 \text{ torr} \times 300^{\circ}\text{K}}$$

$$V_2 = 419 \text{ liters.}$$

• PROBLEM 42

A chemist has a certain amount of gas under a pressure of 33.3 atm; it occupies 30 l at 273°C . For his research, however, the gas must be at standard conditions. Under standard conditions what will the volume of the gas be?

Solution: Standard conditions are defined to be 0°C and

1 atm. Hence, the gas is cooled and the pressure on it is decreased. The combined gas law relates pressure P , volume V and absolute temperature T .

$$k = \frac{PV}{T}$$

where k is a constant that is characteristic to the system. Hence,

$$\frac{P_I V_I}{T_I} = \frac{P_F V_F}{T_F}$$

where the subscript I indicates the initial values and the subscript F indicates the final states.

In this problem one is asked to solve for V_F . The temperatures in $^{\circ}\text{C}$ are converted to $^{\circ}\text{K}$ by adding 273.

$$T_I = 273^{\circ}\text{C} + 273 = 546^{\circ}\text{K}$$

$$T_F = 0^{\circ}\text{C} + 273 = 273^{\circ}\text{K}$$

Solving for V_F :

$$\frac{(33.3 \text{ atm})(30 \text{ l})}{(546^{\circ}\text{K})} = \frac{(1 \text{ atm})(V_F)}{(273^{\circ}\text{K})}$$

$$V_F = 499.5 \text{ l.}$$

• PROBLEM 43

On a hot day, the pressure in an automobile tire increases. Assuming that the air in a tire at 59°F increases in pressure from 28.0 lbs/in^2 (1.9 atm.) to 30.0 lbs/in^2 (2.04 atm.), (a) what is the temperature of the air in the tire, assuming no change in volume? (b) What will the pressure be if the temperature rises to 106°F ?

Solution: (a) Because pressure, volume, and temperature are involved, one thinks of the combined gas law. This law can be stated: For a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. This law can be stated algebraically as:

$$\frac{PV}{T} = K,$$

where P is the pressure, V is the volume, T is the absolute temperature, and K is a constant. This means that if two of the parameters are changed, the third

will adjust itself so that $PV/T = K$ is still true. The following equation is thus true:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 is the original pressure, V_1 is the original volume, T_1 is the original absolute temperature, P_2 is the new pressure, V_2 is the new volume, and T_2 is the new absolute temperature. When $V_1 = V_2$, as in this problem, it follows from the formula that pressure is directly proportional to absolute temperature. In this case, the law can be written

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

In this problem, the temperature is given in $^{\circ}\text{F}$, and must be converted to $^{\circ}\text{K}$ before it is used in the combined gas law. This is done by first converting it to $^{\circ}\text{C}$ and then adding 273.15°C are converted to $^{\circ}\text{C}$ by use of the following equation:

$$^{\circ}\text{C} = 5/9 (^{\circ}\text{F} - 32)$$

$$T_1 \text{ in } ^{\circ}\text{C} = 5/9 (59^{\circ} - 32^{\circ}) = 15^{\circ}\text{C}$$

$$T_1 = 15 + 273 = 288^{\circ}\text{K}.$$

One is given P_1 , P_2 , and T_1 and asked to find T_2 .

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$P_1 = 28.0 \text{ lbs/in}^2$$

$$T_1 = 288^{\circ}\text{K}$$

$$P_2 = 30.0 \text{ lbs/in}^2$$

$$T_2 = ?$$

$$\frac{28.0 \text{ lbs/in}^2}{288^{\circ}\text{K}} = \frac{30.0 \text{ lbs/in}^2}{T_2}$$

$$T_2 = \frac{30.0 \text{ lbs/in}^2 \times 288^{\circ}\text{K}}{28.0 \text{ lbs/in}^2} = 309^{\circ}\text{K}.$$

The absolute temperature can now be converted to $^{\circ}\text{F}$ by first subtracting 273.15°C , and then using the equation for conversion from $^{\circ}\text{C}$ to $^{\circ}\text{F}$.

$$^{\circ}\text{F} = 9/5 ^{\circ}\text{C} + 32$$

$$T_2 \text{ in } ^{\circ}\text{C} = 309^{\circ} - 273 = 36^{\circ}\text{C}$$

$$T_2 \text{ in } ^{\circ}\text{F} = 9/5 (36^{\circ}) + 32 = 97^{\circ}\text{F}.$$

(b) One can again use the shortened form of the combined gas law, here:

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

Here, one is given T_1 , T_2 , and P_1 and asked to find P_2 . The temperatures must be converted to the absolute scale before use. The same method as used in part (a) will be applied.

$$T_1 \text{ in } ^\circ\text{C} = 5/9 (59 - 32) = 15^\circ\text{C}$$

$$T_1 \text{ in } ^\circ\text{K} = 15 + 273 = 288^\circ\text{K}$$

$$T_2 \text{ in } ^\circ\text{C} = 5/9 (106 - 32) = 41^\circ\text{C}$$

$$T_2 = 41 + 273 = 314^\circ\text{K}$$

The P_2 can now be found

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$P_1 = 28.0 \text{ lbs/in}^2$$

$$T_1 = 288^\circ\text{K}$$

$$P_2 = ?$$

$$T_2 = 314^\circ\text{K}$$

$$\frac{28.0 \text{ lbs/in}^2}{288^\circ\text{K}} = \frac{P_2}{314^\circ\text{K}}$$

$$P_2 = \frac{314^\circ\text{K} \times 28.0 \text{ lbs/in}^2}{288^\circ\text{K}} = 30.5 \text{ lbs/in}^2.$$

• PROBLEM 44

When J.F. Piccard made a stratosphere flight in a balloon, the balloon seemed to be only half filled as it left the ground near Detroit. The gas temperature was about 27°C , the pressure 700 mm, and the volume of gas in the balloon 80,000 cubic feet (2.26×10^6 liters). What was the gas volume at high altitude where the temperature was -3°C , and the pressure 400 mm?

Solution: The solution to this problem necessitates the use of the combined gas law. It is stated as follows: For a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. The general formula for the combined gas laws may be written

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \quad \text{or} \quad V_2 = V_1 \times \frac{P_1}{P_2} \times \frac{T_2}{T_1}$$

in which P_1 and T_1 are pressure and absolute temperature

of the gas at volume V_1 ; and P_2 and T_2 are pressure and absolute temperature at volume V_2 .

Thus, if

$$P_1 = 700 \text{ mm} \quad \text{and} \quad P_2 = 400 \text{ mm}$$

$$V_1 = 80,000 \text{ cu.ft.} \quad T_2 = 273^\circ - 3^\circ = 270^\circ\text{K}$$

$$T_1 = 27^\circ + 273^\circ = 300^\circ\text{K}$$

then V_2 can be determined by substitution in the general formula above.

$$V_2 = (80,000) \left(\frac{700 \text{ mm}}{400 \text{ mm}} \right) \left(\frac{270^\circ\text{K}}{300^\circ\text{K}} \right) = 125,000 \text{ cu. ft.} \\ (3.54 \times 10^6 \text{ liters})$$

AVOGADRO'S LAW-THE MOLE CONCEPT

• PROBLEM 45

How many moles are there in one atom?

Solution: A mole of atoms is defined as containing Avogadro's Number of atoms. Avogadro's number is 6.02×10^{23} . Therefore, the number of moles in one atom is equal to 1 atom divided by 6.02×10^{23} atoms/mole.

$$\begin{aligned} \text{No. of moles} &= \frac{1 \text{ atom}}{6.02 \times 10^{23} \text{ atoms/mole}} \\ &= 1.66 \times 10^{-24} \text{ moles.} \end{aligned}$$

• PROBLEM 46

During the course of World War I, 1.1×10^5 kg of poison gas was fired on Allied soldiers by German troops. If the gas is assumed to be phosgene (COCl_2), how many molecules of gas does this correspond to?

Solution: To solve this problem we must convert mass to number of moles and then multiply by Avogadro's number to obtain the corresponding number of molecules.

The number of moles of gas is given by

$$\text{moles} = \frac{\text{mass (grams)}}{\text{molecular weight of } \text{COCl}_2 \text{ (g/mole)}}$$

The mass of gas is $1.1 \times 10^8 \text{ kg} = 1.1 \times 10^8 \text{ kg} \times 10^3 \text{ g/kg} = 1.1 \times 10^{11} \text{ g}$. The molecular weight of COCl_2 is obtained by adding the atomic weights (atm. wgt.) of its constituents. Thus,

$$\begin{aligned} \text{molecular weight (COCl}_2\text{)} &= \text{atm wgt(C)} + \text{atm wgt(O)} \\ &\quad + 2 \times \text{atm wgt(Cl)} \\ &= 12.0 \text{ g/mole} + 16.0 \text{ g/mole} \\ &\quad + 2 \times 35.5 \text{ g/mole} \\ &= 99 \text{ g/mole.} \end{aligned}$$

Hence, the number of moles of gas is

$$\begin{aligned} \text{moles} &= \frac{\text{mass}}{\text{molecular weight}} = \frac{1.1 \times 10^{11} \text{ g}}{99 \text{ g/mole}} \\ &\approx 1.1 \times 10^9 \text{ moles.} \end{aligned}$$

Multiplying the number of moles by Avogadro's number, we obtain the number of molecules of gas:

$$\begin{aligned} \text{number of molecules} &= \text{moles} \times \text{Avogadro's number} \\ &= 1.1 \times 10^9 \text{ moles} \times 6 \times 10^{23} \text{ molecules/mole} \\ &= 6.6 \times 10^{32} \text{ molecules.} \end{aligned}$$

• PROBLEM 47

For a single year, the motor vehicles in a large city produced a total of $9.1 \times 10^6 \text{ kg}$ of the poisonous gas carbon monoxide (CO). How many moles of CO does this correspond to?

Solution: The number of moles of a substance is equal to the quotient of the mass (in grams) of that substance and its molecular weight (in g/mole), or

$$\text{moles} = \frac{\text{mass (g)}}{\text{molecular weight (g/mole)}}$$

The mass of CO is $9.1 \times 10^6 \text{ kg} = 9.1 \times 10^6 \text{ kg} \times 1000 \text{ g/kg} = 9.1 \times 10^9 \text{ g}$. The molecular weight of CO is the sum of the atomic weight of C and the atomic weight of O, or

$$\text{molecular weight of (CO)} = \text{atomic weight (C)} + \text{atomic weight (O)}$$

$$= 12 \text{ g/mole} + 16 \text{ g/mole}$$

$$= 28 \text{ g/mole.}$$

Hence,

$$\begin{aligned} \text{moles} &= \frac{\text{mass (g)}}{\text{molecular weight (g/mole)}} = \frac{9.1 \times 10^9 \text{ g}}{28 \text{ g/mole}} \\ &= 3.3 \times 10^8 \text{ moles.} \end{aligned}$$

• PROBLEM 48

An automobile travelling at 10 miles per hour produces 0.33 lb of CO gas per mile. How many moles of CO are produced per mile?

Solution: The number of moles of a substance is equal to the quotient of the mass (in grams) of that substance and its molecular weight (in g/mole), or

$$\text{moles} = \frac{\text{mass (g)}}{\text{molecular weight (g/mole)}}$$

The mass of CO is 0.33 lb = 0.33 lb × 454 g/lb = 150 g.
The molecular weight of CO is the sum of the atomic weight of C and the atomic weight of O, or

$$\begin{aligned} \text{molecular weight (CO)} &= \text{atomic weight (C)} + \text{atomic weight (O)} \\ &= 12 \text{ g/mole} + 16 \text{ g/mole} \\ &= 28 \text{ g/mole.} \end{aligned}$$

Hence,

$$\begin{aligned} \text{moles} &= \frac{\text{mass (g)}}{\text{molecular weight (g/mole)}} = \frac{150 \text{ g}}{28 \text{ g/mole}} \\ &= 5.4 \text{ moles per mile.} \end{aligned}$$

• PROBLEM 49

A flask containing H₂ at 0°C was sealed off at a pressure of 1 atm and the gas was found to weigh, 4512 g. Calculate the number of moles and the number of molecules of H₂ present.

Solution: A mole is defined as 6.023×10^{23} molecules of a substance. Hydrogen has a molecular weight of 1.008,

therefore, 1 mole of H_2 weighs 2.016 grams. The number of moles present is therefore

$$\text{moles} = \frac{\text{grams}}{\text{M.W.}} = \frac{.4512 \text{ g}}{2.016 \text{ g/mole}} = .2238 \text{ mole of } H_2$$

To calculate the number of molecules, recall that 1 mole of any gas at STP has 6.023×10^{23} molecules/mole. Therefore,

$$\begin{aligned} \text{no. of molecules} &= 6.023 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \times \text{no. of moles} \\ &= 6.023 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \times 0.2238 \text{ moles} \\ &= 1.348 \times 10^{23} \text{ molecules of } H_2. \end{aligned}$$

THE IDEAL GAS LAW

• PROBLEM 50

Three researchers studied 1 mole of an ideal gas at 273°K in order to determine the value of the gas constant, R. The first researcher found that at a pressure of 1 atm the gas occupies 22.4 l. The second researcher found that the gas occupies 22.4 l at a pressure of 760 torr. Finally, the third researcher reported the product of pressure and volume as 542 cal. What value for R did each researcher determine?

Solution: This problem is an application of the ideal gas equation, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. Specifically, we are trying to determine R from the relation $R = PV/nT$. All three researchers worked with one mole of gas ($n = 1$) at $T = 273^\circ\text{K}$. Their results are as follows:

First researcher: $P = 1 \text{ atm}$, $V = 22.4 \text{ l}$.

$$R = \frac{PV}{nT} = \frac{1 \text{ atm} \times 22.4 \text{ l}}{1 \text{ mole} \times 273^\circ\text{K}} = 0.0821 \frac{\text{l-atm}}{\text{K-mole}}$$

Second researcher: $P = 760 \text{ torr} = 760 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 1 \text{ atm}$, $V = 22.4 \text{ l}$.

$$R = \frac{PV}{nT} = \frac{1 \text{ atm} \times 22.4 \text{ l}}{1 \text{ mole} \times 273^\circ\text{K}} = 0.0821 \frac{\text{l-atm}}{\text{K-mole}}$$

Third researcher: $PV = 542 \text{ cal} = nRT$
 $= 1 \text{ mole } (R) (273^\circ\text{K})$

$$R = \frac{PV}{nT} = \frac{542 \text{ cal}}{(1 \text{ mole})(273^\circ\text{K})} = 1.99 \text{ cal/mole } ^\circ\text{K}.$$

• PROBLEM 51

How many moles of hydrogen gas are present in a 50 liter steel cylinder if the pressure is 10 atmospheres and the temperature is 27°C ? $R = .082 \text{ liter-atm/mole } ^\circ\text{K}$.

Solution: In this problem, one is asked to find the number of moles of hydrogen gas present where the volume, pressure and temperature are given. This would indicate that the Ideal Gas Law should be used because this law relates these quantities to each other. The Ideal Gas Law can be stated:

$$PV = nRT,$$

where P is the pressure, V is the volume, n is the number of moles, R is the gas constant ($.082 \text{ liter-atm/mole } ^\circ\text{K}$), and T is the absolute temperature. Here, one is given the temperature in $^\circ\text{C}$, which means it must be converted to the absolute scale. P , V , and R are also known. To convert a temperature in $^\circ\text{C}$ to the absolute scale, add 273 to the temperature in $^\circ\text{C}$.

$$T = 27 + 273 = 300^\circ\text{K}$$

Using the Ideal Gas Law:

$$PV = nRT \quad \text{or} \quad n = \frac{PV}{RT}$$

$$P = 10 \text{ atm}$$

$$V = 50 \text{ liters}$$

$$R = .082 \text{ liter-atm/mole } ^\circ\text{K}$$

$$T = 300^\circ\text{K}$$

$$n = \text{number of moles of } \text{H}_2 \text{ present}$$

$$n = \frac{(10 \text{ atm})(50 \text{ liters})}{(.082 \text{ liter-atm/mole } ^\circ\text{K})(300^\circ\text{K})}$$
$$= 20 \text{ moles.}$$

The barometric pressure on the lunar surface is about 10^{-10} torr. At a temperature of 100°K , what volume of lunar atmosphere contains (a) 10^6 molecules of gas, and (b) 1 millimole of gas?

Solution: This problem is an application of the ideal gas equation, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. Solving for V , $V = nRT/P$. In the first part of this problem, we use the definition

$$n = \frac{\text{number of molecules}}{\text{Avogadro's number}} = \frac{N}{A}. \text{ We must then substitute}$$

into the formula $V = nRT/P = (N/A)(RT/P)$ in order to obtain the volume corresponding to N molecules. In the second part of this problem we can use $V = nRT/P$ directly, remembering that 1 millimole = 10^{-3} mole.

Hence, for the first part of the problem,

$$\begin{aligned} V &= \frac{N}{A} \frac{RT}{P} = \frac{10^6 \text{ molecules}}{6.02 \times 10^{23} \text{ molecules/mole}} \\ &\times \frac{0.0821 \text{ l-atm/}^\circ\text{K-mole} \times 100^\circ\text{K}}{1.316 \times 10^{-13} \text{ atm}} \\ &= 1.04 \times 10^{-4} \text{ l, where we have used } P = 10^{-10} \text{ torr} = \\ &10^{-10} \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 1.316 \times 10^{-13} \text{ atm.} \end{aligned}$$

For the second part of the problem, $V = \frac{nRT}{P}$

$$\begin{aligned} &= \frac{10^{-3} \text{ mole} \times 0.0821 \text{ l-atm/}^\circ\text{K-mole} \times 100^\circ\text{K}}{1.316 \times 10^{-13} \text{ atm}} \\ &= 6.24 \times 10^{10} \text{ l.} \end{aligned}$$

Describe the curve one would obtain by plotting pressure versus volume for an ideal gas in which the temperature and number of moles of gas are held constant.


Solution: This problem requires a plot of the ideal gas equation. This equation reads

$$PV = nRT,$$

where P = pressure of gas, V = volume of gas, n = number of moles of gas, R = gas constant, and T = absolute temperature of gas. Since R is a constant and n and T are held constant, the product nRT may be combined into a single constant, call it k . Then, $PV = nRT = k$, or

$$PV = k$$

which is the equation of a hyperbola. Plotting P on the abscissa axis and V on the ordinate axis, we obtain the following curve:



Note: Even if we did not know $PV = k$ is the graph of a hyperbola, we could still obtain this graph. For example, if k is a constant, P and V must vary inversely with each other. In other words, if P is large, V must be small for k to remain constant. Likewise, if V is large, P must be small.

• PROBLEM 54

A sample of a gas exhibiting ideal behavior occupies a volume of 350 mL at 25.0°C and 600 torr. Determine its volume at 600°C and 25.0 torr.

Solution: We can solve this problem by determining the number of moles, n , of gas from the ideal gas equation, $PV = nRT$ (where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature) by using the first set of conditions and then substituting this value of n along with P and T from the second set of conditions into the ideal gas equation and solving for V . However, we can save useless calculation by denoting the first and second sets of conditions by subscripts "1" and "2", respectively, to obtain

$$P_1V_1 = nRT_1, \quad \text{and} \quad P_2V_2 = nRT_2,$$

where n is the same in both cases. Dividing the second of these equations by the first we obtain

$$\frac{P_2 V_2}{P_1 V_1} = \frac{nRT_2}{nRT_1} \quad \text{or} \quad \frac{P_2 V_2}{P_1 V_1} = \frac{T_2}{T_1},$$

where we have cancelled n and R (both constant). Solving for V_2 , we obtain

$$\begin{aligned} V_2 &= \frac{P_1 V_1}{P_2} \times \frac{T_2}{T_1} \\ &= \frac{600 \text{ torr} \times 350 \text{ ml}}{25.0 \text{ torr}} \times \frac{873.15^\circ\text{K}}{298.15^\circ\text{K}} \\ &= 24.6 \times 10^3 \text{ ml} \\ &= 24.6 \text{ l.} \end{aligned}$$

• PROBLEM 55

A sample of gas occupies 14.3 l at 19°C and 790 mm Hg. Determine the number of moles of gas present. What volume will this same amount of gas occupy at 190°C and 79.0 mm Hg?

Solution: This problem involves an application of the ideal gas equation, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, T = absolute temperature. We will solve this equation for n and then use this value of the number of moles, and the ideal gas equation, to obtain V at the new temperature and pressure.

For our gas, we initially have $V = 14.3 \text{ l}$, $T = 19^\circ\text{C} = 292.15^\circ\text{K}$, $P = 790 \text{ mmHg} \times 1 \text{ atm}/760 \text{ mmHg} = 1.039 \text{ atm}$ and $R = 0.0821 \text{ l-atm}/^\circ\text{K-mole}$. Solving the ideal gas equation for n , we obtain

$$\begin{aligned} n &= \frac{PV}{RT} = \frac{1.039 \text{ atm} \times 14.3 \text{ l}}{0.0821 \text{ l-atm}/^\circ\text{K-mole} \times 292.15^\circ\text{K}} \\ &= 0.6194 \text{ mole.} \end{aligned}$$

Under the new conditions, we have $T = 190^\circ\text{C} = 463.15^\circ\text{K}$ and $P = 79.0 \text{ mmHg} \times 1 \text{ atm}/760 \text{ mmHg} = 0.1039 \text{ atm}$. Solving the ideal gas equation for V , the new volume is

$$\begin{aligned} V &= \frac{nRT}{P} = \frac{0.6194 \text{ mole} \times 0.0821 \text{ l-atm}/^\circ\text{K-mole} \times 463.15^\circ\text{K}}{0.1039 \text{ atm}} \\ &= 226.68 \text{ l.} \end{aligned}$$

A 3-liter bulb containing oxygen at 195 torr is connected to a 2-liter bulb containing nitrogen at 530 torr by means of a closed valve. The valve is opened and the two gases are allowed to equilibrate at constant temperature, T . Calculate the equilibrium pressure.

Solution: We will solve for the final or equilibrium pressure, P_f , by using the ideal gas law,

$$P_f = \frac{nRT}{V_f}$$

where n is the number of moles of O_2 (n_{O_2}) plus the number of moles of N_2 (n_{N_2}), that is, the total number of moles of gas, R is the gas constant, T the absolute temperature, and V_f the final volume (3 liters + 2 liters = 5 liters). We cannot solve this equation directly, since we do not know T .

The quantities n_{O_2} and n_{N_2} are calculated using the initial conditions and the ideal gas law,

$$n_a = \frac{P_a V_a}{RT}$$

where P_a is the original partial pressure of gas a and V_a is the volume initially occupied by gas a . Thus,

$$n_{O_2} = \frac{P_{O_2} V_{O_2}}{RT} = \frac{195 \text{ torr} \times 3 \text{ liters}}{RT}, \quad \text{and}$$

$$n_{N_2} = \frac{P_{N_2} V_{N_2}}{RT} = \frac{530 \text{ torr} \times 2 \text{ liters}}{RT}$$

Multiplying both equations by RT and then adding,

$$\begin{aligned} n_{O_2} RT + n_{N_2} RT \\ = \left(\frac{195 \text{ torr} \times 3 \text{ liters}}{RT} \right) RT + \left(\frac{530 \text{ torr} \times 2 \text{ liters}}{RT} \right) RT \\ (n_{O_2} + n_{N_2}) RT = 195 \text{ torr} \times 3 \text{ liters} + 530 \text{ torr} \\ \times 2 \text{ liters}. \end{aligned}$$

Dividing both sides by V_f , we obtain

$$(n_{O_2} + n_{N_2}) \frac{RT}{V_f} = \frac{195 \text{ torr} \times 3 \text{ liters} + 530 \text{ torr} \times 2 \text{ liters}}{V_f}.$$

But $n_{O_2} + n_{N_2} = n$, and $P_f = nRT/V_f$. Hence,

$$\begin{aligned}P_f &= \frac{nRT}{V_f} = \left(n_{O_2} + n_{N_2} \right) \frac{RT}{V_f} \\&= \frac{195 \text{ torr} \times 3 \text{ liters} + 530 \text{ torr} \times 2 \text{ liters}}{V_f} \\&= \frac{195 \text{ torr} \times 3 \text{ liters} + 530 \text{ torr} \times 2 \text{ liters}}{5 \text{ liters}} \\&= 329 \text{ torr.}\end{aligned}$$

• PROBLEM 57

Determine the molecular weight of a gas if 4.50 g of it occupies 4.0 liters at 950 torr Hg and 182°C.
 $R = .082 \text{ liter} \cdot \text{atm/mole} \cdot ^\circ\text{K}.$

Solution: One can find the molecular weight of this gas, once one knows the number of moles present in 4.50 g. The molecular weight is equal to 4.50 g divided by the number of moles present

$$MW = \frac{4.50 \text{ g}}{\text{no. of moles}}$$

The number of moles of the gas can be found by using the Ideal Gas Law. This law is stated

$$PV = nRT,$$

where P is the pressure, V is the volume, n is the number of moles, R is the gas constant ($0.082 \text{ liter} \cdot \text{atm/mole} \cdot ^\circ\text{K}$) and T is the absolute temperature. One must convert the pressure in torr to atm to use the gas constant. There are 760 torr in 1 atm. Thus, torr are converted to atm by multiplying the number of torr by 1 atm/760 torr.

$$\begin{aligned}P \text{ (in atm.)} &= 950 \text{ torr} \times 1 \text{ atm}/760 \text{ torr} \\&= 1.25 \text{ atm.}\end{aligned}$$

To use the Ideal Gas Law, the temperature must be in $^\circ\text{K}$. Here, it is given in $^\circ\text{C}$. Temperature in $^\circ\text{C}$ can be converted to $^\circ\text{K}$ by adding 273 to the temperature in $^\circ\text{C}$.

$$T = 182 + 273 = 455^\circ\text{K}$$

One can now use the Ideal Gas Law to solve for n .

$$n = \frac{PV}{RT}$$

$$P = 1.25 \text{ atm.}$$

$$V = 4.0 \text{ liters}$$

$$R = 0.082 \text{ liter-atm/mole-}^\circ\text{K}$$

$$T = 455^\circ\text{K}$$

$$n = \frac{1.25 \text{ atm} \times 4.0 \text{ liters}}{0.082 \frac{\text{liter-atm}}{\text{mole-}^\circ\text{K}} \times 455^\circ\text{K}}$$

$$= 0.13 \text{ moles}$$

In 4.50 g of this gas, there are 0.13 moles. The molecular weight can now be found.

$$\text{MW} = \frac{4.50 \text{ g}}{0.13 \text{ moles}} = 34.7.$$

• PROBLEM 58

A research worker isolated an unknown compound from one of his reaction products. This compound was a volatile liquid with a boiling point of 65°C . A .134 g sample of the liquid was vaporized and the gas was collected. The temperature of the collecting chamber was kept at 98°C in order that liquefaction would not occur. After all the sample was vaporized, the volume of the gas was measured as .0532 liters at 98°C and 737 mm. Calculate the molecular weight.

Solution: We are dealing with the variables m = mass, V = volume, P = pressure, and T = temperature. Whenever this occurs, it indicates that one should employ the ideal gas equation, $PV = nRT$, where n = moles and R = universal gas constant. An ideal gas is one in which the gas molecules take up no space and do not attract one another. Real gases do not completely meet these conditions, but at values in the neighborhood of standard conditions (0°C and 1 atm), the difference in the real and ideal values is small enough such that the ideal gas law is accurate enough to carry out the calculations.

One further bit of information is required. Since M.W. (Molecular Weight) does not appear anywhere in the equation, a relation must be found between M.W. and one of the 4 variables. The relation is n = moles = mass/M.W. Substituting this into the ideal gas equation we obtain

$$PV = \frac{m}{\text{M.W.}} RT$$

Solving for the molecular weight, we obtain:

$$\text{MW} = \frac{mRT}{PV}$$

$m = \text{mass of the sample} = .134 \text{ g}$

$R = \text{gas constant} = .082 \text{ liter} \cdot \text{atm/mole} \cdot ^\circ\text{K}$

$T = 98^\circ\text{C} = 273 + 98 = 371^\circ\text{K}$

$V = 0.0532 \text{ l}$

$P = 737 \text{ mm}$. For the formula, P must be given in atmospheres. To do this, we multiply 737 mm by the conversion factor of $\frac{1 \text{ atm}}{760 \text{ mm}}$. Thus,

$$737 \text{ mm} = 737 \text{ mm} \cdot \frac{1 \text{ atm}}{760 \text{ mm}} = \frac{737}{760} \text{ atm}.$$

$$MW = \frac{(0.134)(0.082)(371)}{(\frac{737}{760})(0.0532)} = 79.018 \text{ g/mole}.$$

• PROBLEM 59

Assuming ideal gas behavior, what is the molecular weight of a gas with a density of 2.50 g/liter at 98°C and .974 atm?

Solution: You must employ the ideal gas law to answer this question. This law, also called the equation of state for an ideal gas, relates pressure, volume, temperature, and moles to each other quantitatively. It states $PV = nRT$, where R is called the universal gas constant, P = pressure, V = volume, T = temperature in Kelvin, and n = number of moles. You can also write this equation as $n/V = P/RT$ to determine moles per liter. By substituting,

$$\frac{.974 \text{ atm}}{(.0821 \text{ liter} \cdot \text{atm/mole} \cdot ^\circ\text{K})(371^\circ\text{K})} = .0320 \text{ moles/liter}.$$

Therefore, according to the above calculation, you know there are .0320 moles in one liter. However, the density is 2.50 g/liter. Hence, you can compare the two, and obtain the fact that .0320 moles weighs 2.50 g. From this, the molecule weight determination follows via a proportion. If 0.032 moles weighs 2.50 g, then 1 mole weighs x grams, which equals the molecular weight. In other words,

$$\frac{.032}{2.50} = \frac{1}{x}. \text{ Solving for } x, \text{ we obtain } x = 78.1 \text{ g}.$$

• PROBLEM 60

A cylinder contains oxygen at a pressure of 10 atm and a temperature of 300°K . The volume of the cylinder is 10 liters. What is the mass of the oxygen?

Solution: In this problem, we are dealing with 4 variables M = mass, V = volume, P = pressure, and T = temperature. Three of them are given; we must calculate the fourth. These variables are governed by the ideal gas law.

An ideal gas is one in which the molecules have no attraction for one another and the molecules themselves occupy no space (a situation contrary to fact). Real gases do not completely satisfy either of these conditions, but under conditions close to STP (0°C and 1 atm), the real gases come very close to meeting these conditions. Therefore, the difference between the real and ideal values is small enough such that the ideal gas law equation can be used. It is:

$$PV = nRT$$

where n is the number of moles. Moles is defined as grams/molecular weight. R is a gas constant equal to .082 (liter)·(atm)/(°K)·(mole).

We are asked to find the mass of oxygen, yet there is no value for mass in the ideal gas equation. Therefore, we must find an equation that involves mass and moles. That equation is moles = grams/M.W. The molecular weight of O_2 = 32 g/mole. Therefore, substitute the mole equation into the ideal gas equation

$$PV = nRT = \frac{M}{\text{M.W.}} RT$$


The only unknown value is mass and the solution is arrived at by substituting in the appropriate values

$$(10 \text{ atm})(10 \text{ liter}) = \frac{m}{32} (.082)(300^{\circ}\text{K})$$

$$m = \frac{32 \times 10 \times 10}{.082 \times 300} = 130.08 \text{ grams.}$$

• PROBLEM 61

Metallic sodium violently liberates H_2 gas according to the reaction



If you collect the gas at 25°C under 758 mm Hg of pressure, and it occupies 2.24 liters, (a) find the number of moles of hydrogen produced, (b) find the number of moles of water decomposed and (c) find the percent error in (a) if the gaseous hydrogen had been collected dry, assuming the vapor pressure of water is 23.8 mm at 25°C . ($R = .0821 \text{ l-atm/mole } ^{\circ}\text{K}$.)

Solution: (a) You want to determine the number of moles of hydrogen gas liberated given its pressure, volume and temperature. Therefore, you make use of the equation of state, which indicates that $PV = nRT$, where P = pressure, V = volume, R = universal gas constant, T = temperature in Kelvin (Celsius plus 273°), and n = moles. Thus, to find n , substitute into this equation and solve for n .

In 1 atm, there is 760 mm of pressure. You use atmospheres as the units of pressure, since R is expressed in atmospheres. Hence,


$$P = 758 \text{ mm} = \frac{758}{760} \text{ atms} = .997 \text{ atm.}$$

We are given R in the problem statement and that $T = 298^\circ\text{K}$ and $V = 2.24 \text{ l}$. Substituting,

$$n = \frac{(.997)(2.24)}{(.0821)(298^\circ\text{K})} = .0913 \text{ mole H}_2.$$

Now that you know how many moles of H_2 are liberated, you also know how much water decomposed via the stoichiometry of the reaction.

In (b), therefore, you make use of the equation



which tells you that for every mole of $\text{H}_2\text{(g)}$ produced you have 2 moles of H_2O decomposed. You calculated a release of .0913 moles of H_2 . Therefore, $2 \times .0913 = .183$ mole of water must have decomposed. For (c) you must realize that if the hydrogen gas were wet (not dried), the pressure given is not the true pressure of the H_2 . For in this pressure is included the vapor pressure of water. Thus, the actual pressure of H_2 is only $758 - 23.8 = 734.2$, where 23.8 was the given vapor pressure of water at 25°C . The % error is, then

$$\frac{23.8}{734.2} \times 100 = 3.24 \%$$

• PROBLEM 62

Compare the number of H_2 and N_2 molecules in two containers described as follows: (1) A 2-liter container of Hydrogen filled at 127°C and 5 atm. (2) A 5-liter container of nitrogen filled at 27°C and 3 atm.

Solution: Avogadro's Law states that equal numbers of molecules are contained in equal volumes of different gases if the pressure and temperature are the same. Therefore, if the conditions were the same, then there would be equal numbers of molecules of H_2 and N_2 .

For reasons of clarity, it is useful to set up a table like that shown below.

gas	V	T	P
H ₂	2 liters	400°K	5 atm
N ₂	5 liters	300°K	3 atm

By using the general gas equation

$$PV = nRT$$

where n = number of moles, and $R = 0.082 \text{ atm-liter/}^\circ\text{K-mole}$, one can find the number of moles of H₂ and N₂.

Thus, for H₂

$$n = \frac{PV}{RT} = \frac{(5 \text{ atm})(2 \text{ l})}{(0.082 \text{ atm-l/mole-}^\circ\text{K})(400^\circ\text{K})} = 0.305 \text{ moles}$$

for N₂

$$n = \frac{PV}{RT} = \frac{(3 \text{ atm})(5 \text{ l})}{(0.082 \text{ atm-l/mole-}^\circ\text{K})(300^\circ\text{K})} = 0.610 \text{ moles.}$$

There are twice as many moles of N₂ as there are of H₂. Since the number of molecules is directly proportional to the number of moles, then there are twice as many molecules of N₂ as molecules of H₂.

• PROBLEM 63

A 0.100 l container maintained at constant temperature contains 5.0×10^{10} molecules of an ideal gas. How many molecules remain if the volume is changed to 0.005 l? What volume is occupied by 10,000 molecules at the initial temperature and pressure?

Solution: Since only the volume changed and no molecules were added to or withdrawn from the system, the number of molecules at a volume of 0.005 l is the same as that at a volume of 0.100 l, or 5.0×10^{10} molecules.

In the second part of the problem we require a relationship between volume and number of molecules, taking into account the fact that the number of molecules can change. We will obtain such a relationship by modifying the ideal gas equation, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. Since the number of moles is equal to the number of molecules, N , divided by Avogadro's number, A , or $n = N/A$, we can write the ideal gas equation as $PV = nRT = N/A RT$, or

$$PVA = NRT.$$

The initial and final pressure and temperature are the same. Also, A and R are constants. Denoting the initial volume and number of molecules by V_i and N_i , respectively, and the final volume and number of molecules by V_f and N_f , respectively, we obtain

$$PV_i A = N_i RT \quad \text{and} \quad PV_f A = N_f RT.$$

Dividing the second of these by the first we obtain

$$\frac{PV_f A}{PV_i A} = \frac{N_f RT}{N_i RT} \quad \text{or} \quad \frac{V_f}{V_i} = \frac{N_f}{N_i},$$

where we have cancelled all the constants. Solving for the final volume we obtain

$$V_f = V_i \times \frac{N_f}{N_i} = 0.100 \text{ l} \times \frac{10,000}{5 \times 10^{10}} = 2 \times 10^{-8} \text{ l}.$$

• PROBLEM 64

At 100°C and 720 mm Hg or 720 torr, what is the density of carbon dioxide, CO_2 ?

Solution: According to the kinetic theory of gases, the parameters of pressure, volume, and temperature of a gas are mutually dependent - that is, a change in one influences the others. In this problem, you want the density of CO_2 . Density is defined as mass/volume. Thus, density = (mass of 1 mole CO_2) / (volume of 1 mole CO_2). The mass of one mole of carbon dioxide is 44 g. (This number represents the sum of the atomic weights of all atoms in the gas.) Consequently, this problem involves only the calculation of CO_2 's volume at 100°C and 720 torr.

At standard temperature and pressure, one mole of CO_2 or any gas occupies 22.4 liters. (Recall that standard temperature and pressure is 760 mm Hg and 0°C.) Because the temperature and pressure has been altered, the volume will no longer be 22.4 liters for one mole of CO_2 .

According to the combined gas law, $PV/T = \text{constant}$.

Thus at standard temperature and pressure,

$$\frac{(760 \text{ mm Hg})(22.4 \text{ liters})}{273^\circ} = \text{a constant}.$$

(You add 273° to the Celsius temperature, since T must be in Kelvin.) At 100°C and 720 mm Hg, you have

$$\frac{(720 \text{ mm Hg}) V}{373^\circ} = \text{a constant.}$$

According to this law, both expressions are equal, i.e.

$$\frac{(760 \text{ mm Hg})(22.4 \text{ liters})}{273^\circ} = \frac{(720 \text{ mm Hg})(V)}{373^\circ}$$

Solving for V in this equation, you obtain
 $V = 32.3 \text{ liters.}$

The density is, therefore, $44/32.3 = 1.32 \text{ g/liter.}$

• PROBLEM 65

Using the ideal gas law, show that for an ideal gas of density ρ , $P/\rho T$ is a constant.

Solution: The ideal gas law reads $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. The number of moles is equal to the mass m of a sample of gas divided by the molecular weight MW of that gas: $n = m/MW$. Then $PV = nRT = (m/MW)RT$. Dividing both sides by the volume, we obtain $P = \left(\frac{m}{MW}\right)\frac{RT}{V}$. But mass per unit volume is density, hence $\rho = m/V$ and this equation becomes $P = (\rho/MW)RT$ or $P/(\rho T) = R/MW$. Since R is a universal constant and MW , the molecular weight, is constant for a particular gas, $P/(\rho T) = R/MW = \text{constant.}$

• PROBLEM 66

The density of a gas is measured as 0.222 g/l at 20.0°C and 800 torr. What volume will 10.0 g of this gas occupy under standard conditions?

Solution: Since we know the density at a given temperature and pressure, we can apply the equation $P/(\rho T) = \text{constant}$, which holds for ideal gases of density ρ at pressure P and absolute temperature T . We must first determine the constant for our system and then use this value along with standard pressure (760 torr) and temperature (273.15°K) to determine the density of the gas under standard conditions. From this density, we obtain the volume that 10.0 g of the gas occupies under standard conditions.

The value of the constant is, as previously indicated, $P/(\rho T)$

$$= \frac{800 \text{ torr}}{0.222 \text{ g/l} \times 20.0^\circ\text{C}} = \frac{800 \text{ torr}}{0.222 \text{ g/l} \times 293.15^\circ\text{K}} = 12.28 \frac{\text{torr-l}}{\text{g-}^\circ\text{K}}$$

Hence, for our gas,

$$\frac{P}{\rho T} = 12.28 \frac{\text{torr-l}}{\text{g-}^\circ\text{K}} \quad \text{or} \quad \rho = \frac{P}{T} \times \frac{1}{12.28 \text{ torr-l/g-}^\circ\text{K}}$$

Under standard conditions, the density of our gas is

$$\begin{aligned} \text{then } \rho &= \frac{P}{T} \times \frac{1}{12.28 \text{ torr-l/g-}^\circ\text{K}} \\ &= \frac{760 \text{ torr}}{273.15^\circ\text{K}} \times \frac{1}{12.28 \text{ torr-l/g-}^\circ\text{K}} = 0.2265 \text{ g/l.} \end{aligned}$$

Density is defined as $\rho = \text{mass/volume}$. Hence, volume = mass/ ρ , and the volume occupied by 10.0 g of our gas under standard conditions is

$$\text{volume} = \frac{\text{mass}}{\rho} = \frac{10.0 \text{ g}}{0.2265 \text{ g/l}} = 44.15 \text{ l.}$$

• PROBLEM 67

At standard conditions, it is found that 28.0 g of carbon monoxide occupies 22.4 l. What is the density of carbon monoxide at 20°C and 600 torr?

Solution: The density of carbon monoxide under STP or standard conditions (pressure = 1 atm or 760 torr and temperature = 0°C or 273.15°K) is $\rho = \text{mass/volume}$. 28 g of CO, carbon monoxide, represents one mole and a mole of any gas occupies 22.4 liters of volume under standard conditions. Thus, $\rho = 28\text{g}/22.4 \text{ l} = 1.25 \text{ g/l}$.

We can apply the equation $P/(\rho T) = \text{constant}$, which holds for ideal gases of density ρ at pressure P and absolute temperature T . We will first determine the constant for our system and then use this value to calculate the density at 20°C = 293.15°K and 600 torr.

The value of the constant for our system is

$$\text{constant} = \frac{P}{\rho T} = \frac{760 \text{ torr}}{1.25 \text{ g/l} \times 273.15^\circ\text{K}} = 2.225 \frac{\text{torr-l}}{\text{g-}^\circ\text{K}}$$

$$\text{Hence, for our system, } \frac{P}{\rho T} = 2.225 \frac{\text{torr-l}}{\text{g-}^\circ\text{K}} \quad \text{or}$$

$$\rho = \frac{P}{T} \times \frac{1}{2.225 \text{ torr-l/g-}^\circ\text{K}} \quad \text{At } 20^\circ\text{C} = 293.15^\circ\text{K and}$$

600 torr, the density of CO is

$$\rho = \frac{P}{T} \times \frac{1}{2.225 \text{ torr} - 1/g - ^\circ K}$$

$$= \frac{600 \text{ torr}}{293.15^\circ K} \times \frac{1}{2.225 \text{ torr} - 1/g - ^\circ K} = .9198 \text{ g/l.}$$

• PROBLEM 68

One gram of an unknown gaseous compound of boron (B, atomic weight = 10.8 g/mole) and hydrogen (H, atomic weight = 1.0 g/mole) occupies 0.820 liter at 1.00 atm at 276 K. What is the molecular formula of this compound?

Solution: The general formula for the compound is $B_a H_b$, where a and b are to be determined. To find these, we must first find the molecular weight of the compound by using the ideal gas law,

$$PV = nRT,$$

where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. n equals the mass, m, divided by the molecular weight MW, $PV =$

$$nRT = \frac{m}{MW} RT, \text{ or } MW = \frac{mRT}{PV}.$$

$$MW = \frac{mRT}{PV} = \frac{1 \text{ g} \times 0.082 \text{ liter} \cdot \text{atm/mole} \cdot ^\circ K \times 276^\circ K}{1.00 \text{ atm} \times 0.820 \text{ liter}}$$

$$= 27.6 \text{ g/mole.}$$

To find a and b, we use the relation $MW_{B_a H_b} = a \times$ atomic weight of B plus $b \times$ atomic weight of H, or $27.6 \text{ g/mole} = a \times 10.8 \text{ g/mole} + b \times 1.0 \text{ g/mole}.$

By trial and error, we find that $a = 2$ and $b = 6$ ($2 \times 10.8 \text{ g/mole} + 6 \times 1.0 \text{ g/mole} = 27.6 \text{ g/mole}$), so that the formula of our compound is $B_2 H_6$ (diborane).

• PROBLEM 69

Of the several ways in which the supersonic transport (SST) acts as a pollutant, one is by the elimination of carbon dioxide (CO_2). During normal flight, CO_2 is released at a temperature of about $627^\circ C$ and at a pressure of 0.41 atm at an estimated rate of $6.6 \times 10^4 \text{ kg}$ per hour. To what volume does this amount of gas correspond?

Solution: This problem is an application of the ideal gas law, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. Solving for V ,

$$V = \frac{nRT}{P}$$

We know that $R = 0.082(\text{liter} - \text{atm}/\text{mole} - ^\circ\text{K})$, $T = 627^\circ\text{C} = 900^\circ\text{K}$, and $P = 0.41 \text{ atm}$. n is obtained by dividing the mass of CO_2 ($6.6 \times 10^4 \text{ kg} = 6.6 \times 10^4 \text{ kg} \times 10^3 \text{ g/kg} = 6.6 \times 10^7 \text{ g}$) by the molecular weight of CO_2 . The molecular weight of CO_2 is equal to

atomic mass_C + 2 atomic mass_O = 12 g/mole + 2(16 g/mole)
or 44 g/mole. Hence

$$n = \frac{\text{mass } \text{CO}_2}{\text{molecular weight } \text{CO}_2} = \frac{6.6 \times 10^7 \text{ g}}{44 \text{ g/mole}} = 1.5 \times 10^6 \text{ moles.}$$

Substituting the values of n , R , T , and P into the equation for V gives

$$V = \frac{nRT}{P} = \frac{1.5 \times 10^6 \text{ moles} \times 0.082 \text{ liter-atm/mole} - ^\circ\text{K} \times 900^\circ\text{K}}{0.41 \text{ atm}}$$

$$= 2.7 \times 10^8 \text{ liters.}$$

• PROBLEM 70

KMnO_4 is added to a 350 ml solution of an approximately 3% H_2O_2 solution. It is found that 3.94 cm³ of O_2 gas is released. It is collected over H_2O at a pressure (barometric) of 0.941 atm and 28°C . Assuming that the pressure due to water vapor is 0.0373 atm, what is the actual concentration of the H_2O_2 solution?

Solution: The concentration of a solution is given in terms of molarity, i.e., moles/liter. This means that the number of moles of H_2O_2 must be determined, since you already know the volume.

To determine the number of moles of H_2O_2 , you know that, when oxidized, 1 mole of H_2O_2 yields 1 mole of O_2 . Thus, if you know the number of moles of O_2 produced, you also know the moles of H_2O_2 originally present. To find the number of moles of O_2 , you must use the equation of state. This tells us $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = universal gas constant, ($.082057 \text{ l-atm mol}^{-1} \text{ deg}^{-1}$), and T = temperature in degrees Kelvin (degree Celsius plus 273°). Hence,

$$n_{\text{O}_2} = \frac{PV}{RT}$$

All the values needed to find n_{O_2} are known or can be calculated. The pressure of O_2 will not be .941 atm, since this figure also includes vapor pressure from water. Therefore, the actual pressure of O_2 is .941 - .0373. That is, the difference between barometric and vapor pressure. Substituting, you obtain

$$n_{O_2} = \frac{(.941 - .0373)(3.94 \times 10^{-3})}{(.082)(301)} = 1.44 \times 10^{-4}$$

moles of O_2 . This means there were originally 1.44×10^{-4} moles. Now, recalling the definition of molarity, which represents the concentration of a solution, you have

$$\frac{1.44 \times 10^{-4} \text{ moles}}{.350 \text{ liters}} = 4.12 \times 10^{-4} \text{ M}$$

which is the concentration of the H_2O_2 solution.

• PROBLEM 71

The van der Waal equation is a modification of the ideal gas equation. It reads

$$\left(P + \frac{an^2}{V^2} \right) (V - nb) = nRT$$

where P = pressure, V = volume, n = number of moles, R = gas constant, T = absolute temperature, and (a) and (b) are constants for a particular gas. The term an^2/V^2 corrects the pressure for intermolecular attraction and the term $-nb$ corrects the volume for molecular volume. Using this equation, determine whether a gas becomes more or less ideal when: (a.) the gas is compressed at constant temperature; (b.) more gas is added at constant volume and temperature; and (c.) The temperature of the gas is raised at constant volume.

Solution: As the behavior of a gas approaches ideality, PV/nRT approaches 1. Hence, we must manipulate the van der Waal equation into a form in which it can be determined whether PV/nRT approaches 1 as the variables are changed. Thus,

$$\left(P + \frac{an^2}{V^2} \right) (V - nb) = nRT$$

$$PV - Pnb + \frac{an^2}{V^2} V - \frac{an^2}{V^2} nb = nRT$$

$$PV - Pnb + \frac{an^2}{V} - \frac{an^3b}{V^2} = nRT$$

$$PV = nRT + Pnb - \frac{an^2}{V} + \frac{an^3b}{V^2} .$$

Dividing by nRT we obtain

$$\frac{PV}{nRT} = 1 + \frac{Pb}{RT} - \frac{an}{RTV} + \frac{an^2b}{RTV^2}$$

(a.) When the gas is compressed at constant temperature, volume decreases and pressure increases. Hence, the last three terms on the right side increase, and PV/nRT deviates from 1. The gas thus becomes less ideal.

(b.) If more gas is added at constant volume and temperature, both n and P increase, hence the last three terms on the right side increase and the gas becomes less ideal.

(c.) As the temperature of the gas is increased at constant volume, the pressure increases. The last two terms on the right side become smaller while the second term (Pb/RT) increases slightly. Hence, the gas comes closer to being ideal.

• PROBLEM 72

Using Van der Waal's equation, calculate the pressure exerted by 1 mole of carbon dioxide at 0°C in a volume of (a) 1.00 liter, (b) 0.05 liter.

Solution: Real gases are more compressible than ideal gases because their molecules attract each other. The intermolecular attraction is provided for by adding to the observed pressure a term of n^2a/V^2 in the ideal gas law ($PV = nRT$) where n is the number of moles, V is the volume, and a is a van der Waal's constant. At very high pressures, real gases occupy larger volumes than ideal gases; this effect is provided for by subtracting an excluded volume nb from the observed volume V to give the actual volume in which the molecules move (b is a constant). Thus, van der Waal's full equation is:

$$\left(P + \frac{n^2a}{V^2}\right)(V - nb) = nRT$$

To solve this problem one must 1) convert the temperature from units of $^\circ\text{C}$ to $^\circ\text{K}$ by adding 273° to the $^\circ\text{C}$ temperature. This is done because T is expressed in the absolute. (P = pressure and R = gas constant.) 2) Find the values for the constants (a) and (b) (these can be found in any reference text). For CO_2 ; $a = 3.592 \text{ liter}^2 \text{ atm/mole}^2$; $b = 0.04267 \text{ liter/mole}$. 3) Substitute the known values into the van der Waal's equation and solve for pressure.

Known values:

$$n = 1 \quad R = 0.082 \frac{\text{liter-atm}}{\text{mole-}^\circ\text{K}} \quad T = 273^\circ\text{K}$$

(a) $V = 1.00$ liter. Substituting,

$$\left[P + \frac{(1)^2 (3.592)}{(1.00)^2} \right] \left[1.00 - (1) (0.04267) \right] = (1) (0.082) (273)$$

$$P = \frac{(0.082) (273^\circ\text{K})}{(1.00 - 0.04267)} - \frac{3.592}{(1.00)^2}$$

$$= 23.38 - 3.592$$

$$= 19.79 \text{ atm}$$

(b) $V = 0.05$ liter. Substitute

$$\left[P + \frac{(1)^2 (3.592)}{(0.05)^2} \right] (0.05 - (1) 0.04267) = (1) (0.082) (273 \text{ K})$$

$$P = \frac{(0.082) (273^\circ\text{K})}{(0.05 - 0.04267)} - \frac{3.592}{(0.05)^2}$$

$$= 3054 - 1437$$

$$= 1617 \text{ atm.}$$

• PROBLEM 73

Using the data from the accompanying figure, calculate the pressure exerted by .250 moles of CO_2 in .275 liters at 100°C . Compare this value with that expected for an ideal gas.

van der Waals constants

Gas	a, liter ² atm/mol ²	b, liter/mol
Helium	0.0341	0.0237
Argon	1.35	0.0322
Nitrogen	1.39	0.0391
Carbon dioxide	3.59	0.0427
Acetylene	4.39	0.0514
Carbon tetrachloride	20.39	0.1383

Solution: To solve this problem you must understand the concept of an "ideal gas" and the formulas associated with it. Boyle's law states that $PV = \text{constant}$, if the temperature is fixed. In other words, if T is constant, then a fixed mass of gas occupies a volume inversely proportional to the pressure exerted on it. If a gas obeys this law, it is termed an "ideal gas." When a gas is ideal, $PV = nRT$, where P = pressure, V = volume, n = moles, R = universal gas constant, and T = temperature in Kelvin. When a gas is not ideal, it doesn't obey Boyle's Law;

and instead of the Ideal Gas law you use Van der Waal's equation,

$$\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT, \text{ where, for } CO_2, a =$$

3.59 liter²-atm/mole² and $b = .0427$ liter/mole. These values are called Van der Waal's constants. Substitute the values into these equations and solve for P. As such,

$$\left(P + \frac{(.250)^2 (3.59)}{(.275)^2} \right) [.275 - (.250)(.0427)] \\ = (.250)(.08206)(373).$$

Solving for P, you obtain $P = 26.0$ atm.

If you had considered the gas as ideal, then $PV = nRT$, and $P = 27.8$ atm.

• PROBLEM 74

Of the following two pairs, which member will more likely deviate from ideal gas behavior? (1) N_2 versus CO , (2) CH_4 versus C_2H_6 .

Solution: Characteristics of ideal gases include:

(1) Gases are composed of molecules such that the actual volume of the molecules is negligible compared with the empty space between them. (2) There are no attractive forces between molecules. (3) Molecules do not lose net kinetic energy in collisions. (4) The average kinetic energy is directly proportional to the absolute temperature.

From these assumptions came the ideal equation of state: $PV = nRT$, where P = pressure, V = volume, n = moles, R = universal gas constant, and T = temperature. No gas is ideal; they don't obey these assumptions absolutely. To reflect these limitations the ideal gas law can be written as:

$$\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT.$$

The term $V = nb$ can be thought of as representing the free volume minus the volume occupied by the gas molecules themselves. The magnitude of (b) is proportional to a gas molecule's size. Thus, the greater the size of a gas molecule, the greater the nb is, and, the greater the deviation, since the volume is assumed to equal exactly V in the Ideal Gas Law. $V - nb$ is smaller than V . The greater nb , the smaller it becomes.

The term n/V , a concentration term, when squared, gives probability of collisions. (a) gives a measure of the cohesive force between molecules. Real molecules do have an attraction for each other. The greater the value of (a), the larger n^2a/V^2 becomes, and the larger $(P + n^2a/V^2)$ gets. The ideal gas law assumes a value of exactly P . Thus, as (a) increases, the more deviation.

To determine which gas deviates the most, look at the size of the gas molecules and their dipole moment. The dipole moment is an indication of unlike charges separated by a given amount of distance. Unlike charges on separate molecules will be attracted to each other. Thus, a higher dipole moment indicates greater cohesive attraction among molecules and greater deviation. Proceed as follows:

(1) N_2 versus CO . The CO gas has a net dipole moment, while N_2 does not. This is because O is more electronegative than C . In N_2 both of the atoms are the same and thus their electronegativities are equal and no net dipole moment exists. Thus, CO deviates to a greater extent.

(2) CH_4 versus C_2H_6 . C_2H_6 is a much larger molecule and occupies more volume. Therefore, it is more likely to deviate from ideal gas behavior.

• PROBLEM 75

What do the terms critical temperature and pressure mean? If you want to liquefy a gas, what physical properties must you consider? If one gas has a higher critical temperature than another, what can be said about the relative forces of attraction between like molecules?

Solution: The critical temperature is the temperature above which a gas may not be liquefied regardless of the pressure. The critical pressure is the pressure necessary to liquefy a gas at its critical temperature.

Both temperature and pressure are factors in the condensation of a gas. Molecules of a particular compound have a lower kinetic energy in their liquid state than in their gaseous state.

$$\text{kinetic energy} = \text{mass} \times (\text{velocity})^2$$

Therefore, kinetic energy is proportional to the square of the velocities of the particles. In the gaseous state the velocities of the molecules are greater, when the temperature of a system is decreased the velocities of the molecules decrease and, hence, the kinetic energies decrease. If the pressure of the system is increased, the

particles are forced closer together. At a certain point, when the kinetic energy will be low and/or the particles will be close enough together, the gas will liquefy. If a gas has a high critical temperature, less energy is necessary to liquefy it. In the liquid state the molecules are much closer together than in the gas. When less energy is needed to force the molecules together, the inter-molecular forces of attraction are greater.

CHAPTER 3



GAS MIXTURES AND PHYSICAL PROPERTIES OF GASES

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 61 to 94 for step-by-step solutions to problems.

Mixtures of Ideal Gases

In mixtures of ideal gases there is negligible interaction between molecules — either between the same or different species. Therefore, each component in a mixture of ideal gases behaves as though it were present all by itself.

The pressure exerted by each component in a mixture of gases is called that component's partial pressure. Since the components in a gas mixture do not interact, the partial pressure of a species in a mixture is the total pressure that component would exert if it were the only one present in the system. The ideal gas law from Chapter 2 applies equally well for the individual components in gas mixtures if we replace the total pressure with the partial pressure and the total number of moles with the number of moles of the individual component.

$$P_i V = n_i RT \quad 3-1$$

This equation is identical to Equation 2-1 of Chapter 2 except that P_i is the partial pressure of species i , and n_i is the number of moles of species i .

P_i = partial (absolute) pressure of species i

V = volume

n_i = number of moles of species i

T = absolute temperature

R = universal gas constant

It should be obvious intuitively that the total number of moles in a system (n) will equal the sum of the numbers of moles of each of the individual components.

$$n = n_1 + n_2 + n_3 \dots + n_m = \sum n_i \quad 3-2$$

It follows, therefore, that at constant temperature and volume, the total pressure of a system containing several gaseous species (P) is simply the sum of the partial pressures.

$$P = \sum P_i = (\sum n_i)RT/V = nRT/V \quad 3-3$$

This result (that each species exerts a partial pressure as though it were present alone, and the sum of the partial pressures equals the total pressure) is known as Dalton's Law of Partial Pressures. An important corollary of Dalton's Law can be obtained by multiplying and dividing Equation 3-1 by n , the total number of moles in the system.

$$P_i = (n_i/n)nRT/V = (n_i/n)P = N_i P \quad 3-4$$

or

$$P_i/P = n_i/n = N_i$$

The quantity n_i/n (written N_i) is defined as the mole fraction of species i and is simply the fraction of the total number of moles or molecules in the system that are of component i . Note that the mole fraction represents a fraction of molecules; since unlike molecules normally have different masses, the mole fraction and the mass fraction are not the same. This concept is explained in greater detail in Chapter 4.

In this chapter, problems illustrating principles governing ideal gas mixtures are presented. These problems are best attacked by applying the ideal gas law to each individual species. While Dalton's Law of Partial Pressures is exact only for mixtures of ideal gases, it is often an excellent approximation for real gases at pressures and temperatures where deviations from ideal gas behavior are not significant.

Kinetic Theory of Gases / Graham's Law of Gaseous Diffusion

The kinetic theory of gases, first developed in the mid 1800s, explains many observed properties of gases. The basic postulate of the kinetic theory is that gas molecules move about at high velocities and collide elastically with other molecules and the vessel walls. Collisions with the walls of the

container exert a force we observe as the gas pressure. An important observation is that the kinetic energy associated with the motion of a molecule is proportional to the absolute temperature and is the same for all species.

$$E_k = 1/2 m v^2 = 3/2 k T \quad 3-5$$

where the terms in Equation 3-5 are defined as follows:

E_k = kinetic energy of the gas molecule

m = mass of the gas molecule

v = velocity of the molecule

k = Boltzmann constant ($1.38 \times 10^{-23} \text{ J/K}$)

T = absolute temperature

There is a distribution of gas velocities in any gas, and several different averages are defined in Problem 100. The important fact to remember is that the average energy is a function of temperature and not the mass of the molecule. If Equation 3-5 is multiplied by N_A , the number of molecules in a mole, the following equation is obtained:

$$E_k = 1/2 N_A m v^2 = 1/2 M v^2 = 3/2 R T \quad 3-6$$

where ($N_A \times m$) is the molecular weight, M , of the gas, and ($N_A \times k$) is the universal gas constant, R .

Graham's Law of Gaseous Diffusion follows logically from Equation 3-6. The rate at which a gas molecule will diffuse is proportional to its velocity. Since all gases in a mixture are at the same temperature and have, therefore, the same average kinetic energy per molecule, the molecular velocity is inversely proportional to the square root of the molecular weight.

$$v \propto D \propto \frac{1}{\sqrt{M}} \quad 3-7$$

In Equation 3-7, D is the diffusivity; the equation provides a means for comparison of the diffusivities of different gaseous species. Graham's Law of Diffusion states, however, that the rate of diffusion is inversely proportional to the gas density, ρ . From the ideal gas law the density can be calculated as follows and is proportional to the molecular weight for a given temperature and pressure.

$$\rho = \frac{m}{V} = \frac{n \cdot m}{V}$$

$$PV = nRT \therefore \frac{n}{V} = \frac{P}{RT}$$

$$\frac{n \cdot M}{V} = \frac{PM}{RT} \quad 3-8$$

It follows that, at constant temperature and pressure, the rate of diffusion is inversely proportional to the square root of the molecular weight of a gas.

$$r_1/r_2 = (\rho_2/\rho_1)^{1/2} = (M_2/M_1)^{1/2} \quad 3-9$$

Problems in this chapter which involve the molecular diffusion of different gases all require a comparison of rates of diffusion or diffusivity. These problems are most efficiently attacked by observing the ratios in Equation 3-9 and determining which of the quantities can be calculated from those given.

Step-by-Step Solutions to Problems in this Chapter, "Gas Mixtures and Other Physical Properties of Gases"

MOLE FRACTION

• PROBLEM 76

Calculate the mole fractions of ethyl alcohol, C_2H_5OH , and water in a solution made by dissolving 9.2 g of alcohol in 18 g of H_2O . M.W. of H_2O = 18, M.W. of C_2H_5OH = 46.

Solution: Mole fraction problems are similar to % composition problems. A mole fraction of a compound tells us what fraction of 1 mole of solution is due to that particular compound. Hence,

$$\text{mole fraction of solute} = \frac{\text{moles of solute}}{\text{moles of solute} + \text{moles of solvent}}$$

The solute is the substance being dissolved into or added to the solution. The solvent is the solution to which the solute is added.

The equation for finding mole fractions is:

$$\frac{\text{moles A}}{\text{moles A} + \text{moles B}} = \text{mole fraction A}$$

Moles are defined as grams/molecular weight (MW). Therefore, first find the number of moles of each compound present and then use the above equation.

$$\text{moles of } C_2H_5OH = \frac{9.2 \text{ g}}{46.0 \text{ g/mole}} = .2 \text{ mole}$$

$$\text{moles of } H_2O = \frac{18 \text{ g}}{18 \text{ g/mole}} = 1 \text{ mole}$$

$$\text{mole fraction of } C_2H_5OH = \frac{.2}{1 + .2} = .167$$

$$\text{mole fraction of H}_2\text{O} = \frac{1}{1 + .2} = .833.$$

Note, that the sum of the mole fractions is equal to 1.

• PROBLEM 77

Of the many compounds present in cigarette smoke, such as the carcinogen 3,4-benzo[a]pyrene, some of the more abundant are listed in the following table along with their mole fractions:

<u>Component</u>	<u>Mole Fraction</u>
H ₂	0.016
O ₂	0.12
CO	0.036
CO ₂	0.079

What is the mass of carbon monoxide in a 35.0 ml puff of smoke at standard temperature and pressure (STP)?

Solution: Assuming ideal gas behavior, we will calculate the number of moles of ideal gas in a 35.0 ml volume. Using the mole fraction of CO in smoke, we will then obtain the number of moles of CO in a 35.0 ml volume and finally convert this to a mass.

The molar volume of an ideal gas is 22.4 liter/mole when STP conditions exist, i.e. when temp. = 0°C and pressure = 1 atm. Hence, the number of moles of ideal gas in a 35.0 ml volume is obtained by dividing this volume by the molar volume of gas, or
 $35.0 \text{ ml} / 22.4 \text{ liter/mole} = 0.035 \text{ liter} / 22.4 \text{ liter/mole}$
 $= 1.56 \times 10^{-3} \text{ mole}.$

The mole fraction is defined by the equation

$$\text{mole fraction CO} = \frac{\text{moles CO}}{\text{total number of moles}}$$

solving for the number of moles of CO,

$$\begin{aligned} \text{moles CO} &= \text{mole fraction CO} \times \text{total number of moles} \\ &= 0.036 \times 1.56 \times 10^{-3} \text{ mole} \\ &= 5.6 \times 10^{-5} \text{ mole.} \end{aligned}$$

This is converted to a mass by multiplying by the molecular weight of CO (28 g/mole). Hence,

$$\begin{aligned} \text{mass of CO} &= \text{moles of CO} \times \text{molecular weight of CO} \\ &= 5.6 \times 10^{-5} \text{ mole} \times 28 \text{ g/mole} \\ &= 1.6 \times 10^{-3} \text{ g.} \end{aligned}$$

• PROBLEM 78

It has been estimated that each square meter of the earth's surface supports 1×10^7 g of air above it. If air is 20% oxygen (O_2 , molecular weight = 32 g/mole) by weight, approximately how many moles of O_2 are there above each square meter of the earth?

Solution: This problem is solved by first calculating what weight of oxygen is present in 1×10^7 g of air and then dividing by the molecular weight of oxygen to convert this mass to moles.

Using the definition of weight percent,

$$\text{weight \% of } O_2 = \frac{\text{weight of } O_2}{\text{total weight}} \times 100\%$$

$$20\% = \frac{\text{weight of } O_2}{1 \times 10^7 \text{ g}} \times 100\%$$

$$20 = \frac{\text{weight of } O_2}{1 \times 10^7 \text{ g}} \times 100$$

Solving for the weight of O_2 ,

$$\text{weight of } O_2 = \frac{20}{100} \times 1 \times 10^7 \text{ g} = 2 \times 10^6 \text{ g.}$$

The number of moles of O_2 is equal to the weight of O_2 divided by the molecular weight, or

$$\text{moles of } O_2 = \frac{\text{weight of } O_2}{\text{molecular weight}} = \frac{2 \times 10^6 \text{ g}}{32 \text{ g/mole}}$$

$$\approx 6 \times 10^4 \text{ moles.}$$

Therefore, each square meter of the earth's surface supports 6×10^4 moles of O_2 .

DALTON'S LAW OF PARTIAL PRESSURES

• PROBLEM 79

A mixture of nitrogen and oxygen gas is collected by displacement of water at 30°C and 700 torr Hg pressure. If the partial pressure of nitrogen is 550 torr Hg, what is the partial pressure of oxygen? (Vapor pressure of H_2O at 30°C = 32 torr Hg.)

Solution: Here, one uses Dalton's Law of partial pressures. This law can be stated: Each of the gases in a gaseous mixture behaves independently of the other gases and exerts its own pressure; the total pressure of the mixture being the sum of the partial pressures exerted by each gas present. Stated algebraically

$$P_{\text{total}} = p_1 + p_2 + \dots + p_n$$

where P_{total} is the total pressure and p_1, p_2, \dots, p_n are the partial pressures of the gases present. In this problem, one is told that oxygen and nitrogen are collected over H_2O , which means that there will also be water vapor present in the gaseous mixture. In this case, the equation for the total pressure can be written

$$P_{\text{total}} = P_{\text{O}_2} + P_{\text{N}_2} + P_{\text{H}_2\text{O}}$$

One is given P_{total} , P_{N_2} , and $P_{\text{H}_2\text{O}}$ and asked to find P_{O_2} . This can be done by using the law of partial pressures

$$P_{\text{total}} = P_{\text{O}_2} + P_{\text{N}_2} + P_{\text{H}_2\text{O}}$$

$$P_{\text{total}} = 700 \text{ torr}$$

$$P_{\text{O}_2} = ?$$

$$700 \text{ torr} = P_{\text{O}_2} + 550 \text{ torr} + 32 \text{ torr}$$

$$P_{\text{N}_2} = 550 \text{ torr}$$

$$P_{\text{O}_2} = (700 - 550 - 32) \text{ torr}$$

$$P_{\text{H}_2\text{O}} = 32 \text{ torr}$$

$$= 118 \text{ torr}$$

$$P_{\text{O}_2} = 118 \text{ torr.}$$

• PROBLEM 80

Methane is burnt in oxygen to produce carbon dioxide and water vapor according to the following reaction, $\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g})$. The final pressure of the gaseous products was 6.25 torr. Calculate the partial pressure of the water vapor.

Solution: The partial pressure of each of the components in a mixture of gaseous substances will be equal to the product of the mole fraction of that component in the gas and the total pressure of the system. Given the total vapor pressure of the gases, one can calculate the mole fraction of the water vapor.

Let P_A = partial pressure of the water vapor, N_A = total number of moles of gases produced, N_A = number of moles of water vapor, and P_T = total pressure. This law

can now be expressed in these terms:

$$P_A = \frac{N_A}{N_T} P_T \quad . \quad \text{From the stoichiometry of the}$$

equation, the total number of moles of products is 3. Out of these 2 moles are water vapor. Thus, substituting,

$$P_A = 2/3 (6.25 \text{ torr}) = 4.16 \text{ torr of water vapor.}$$

• PROBLEM 81

A mixture of gaseous oxygen and nitrogen is stored at atmospheric pressure in a 3.7 l iron container maintained at constant temperature. After all the oxygen has reacted with the iron walls of the container to form solid iron oxide of negligible volume, the pressure is measured at 450 torr. Determine the final volume of nitrogen and the initial and final partial pressures of nitrogen and oxygen.

Solution: The partial pressure of each component, which is independent of any other component in a gaseous mixture at a defined volume, is equal to the pressure each component would exert if it were the only gas in that volume. The total pressure of a gaseous mixture is the sum of the partial pressures of the components. These two definitions are sufficient to solve this problem.

Once all the oxygen has reacted, no oxygen is present in the gaseous phase, so that the final partial pressure of oxygen is zero. By the second definition, final total pressure = final partial pressure of N_2 + final partial pressure of O_2

$$450 \text{ torr} = \text{final partial pressure of } N_2 + 0 \text{ torr}$$

or, final partial pressure of N_2 = 450 torr.

But since, after all the oxygen has reacted, only nitrogen fills the entire volume, the final volume of nitrogen is 3.7 l. By the first definition, the initial partial pressure of nitrogen is the same as the final partial pressure of nitrogen, 450 torr. We now again employ the second definition to determine the last remaining unknown quantity, the initial partial pressure of oxygen. Proceeding, we obtain

$$\begin{aligned} \text{initial total pressure} &= \text{initial partial pressure of } O_2 \\ &\quad + \text{initial partial pressure of } N_2 \end{aligned}$$

$$1 \text{ atm} = \text{initial partial pressure of } O_2 + 450 \text{ torr}$$

or, initial partial pressure of O_2 = 1 atm - 450 torr

$$= 760 \text{ torr} - 450 \text{ torr}$$

$$= 310 \text{ torr.}$$

Summarizing these results in tabular form we have:

	<u>Initial</u>	<u>Final</u>
partial pressure of O ₂	310 torr	0 torr
partial pressure of N ₂	450 torr	450 torr
total pressure	760 torr	450 torr

• PROBLEM 82

A cylinder contains 40 g He, 56 g N₂, and 40 g Ar.
 (a) What is the mole fraction of each gas in the mixture? (b) If the total pressure of the mixture is 10 atm, what is the partial pressure of He?

Solution: (a) The mole fraction of a component in a system is defined as the number of moles of that component divided by the sum of all the moles present in the system. Here, one must first calculate the number of moles present of each component, then the mole fractions can be found. The number of moles of each gas can be found by dividing the number of grams present of the gas by the molecular weight. (MW He = 4, MW N₂ = 28, MW Ar = 40.)

$$\text{no. of moles} = \frac{\text{no. of grams}}{\text{MW}}$$

$$\text{no. of moles of He} = \frac{40 \text{ g}}{4 \text{ g/mole}} = 10 \text{ moles}$$

$$\text{no. of moles of N}_2 = \frac{56 \text{ g}}{28 \text{ g/mole}} = 2 \text{ moles}$$

$$\text{no. of moles of Ar} = \frac{40 \text{ g}}{40 \text{ g/mole}} = 1 \text{ mole}$$

The total number of moles of gas in the system is the sum of the number of moles of the three gases. Thus, there are 13 moles of gas in the system. The mole fraction can now be found for each gas by dividing the number of moles of each gas by 13, the total number of moles

$$\text{mole fraction} = \frac{\text{no. of moles}}{\text{total no. of moles in system}}$$

$$\text{mole fraction of He} = \frac{10 \text{ moles}}{13 \text{ moles}} = .77$$

$$\text{mole fraction of N}_2 = \frac{2 \text{ moles}}{13 \text{ moles}} = .15$$

$$\text{mole fraction of Ar} = \frac{1 \text{ mole}}{13 \text{ moles}} = .08$$

(b) In a system where various gases are present, the partial pressure of each gas is proportional to the mole fraction of the gas. The relationship between the partial pressure of a particular gas and the total pressure is

$$\text{partial pressure} = \text{total pressure} \times \text{mole fraction}$$

In this problem, one is given the total pressure of the system and one has found the mole fraction of He. One can now find the partial pressure of He

$$\text{partial pressure of He} = 10 \text{ atm} \times .77 = 7.7 \text{ atm.}$$

• PROBLEM 83

What is the partial pressure of each gas in a mixture which contains 40 g. He, 56 g. N₂, and 16 g. O₂, if the total pressure of the mixture is 5 atmospheres.

Solution: In a mixture of gases, the partial pressure of each gas is proportional to its mole fraction. Here, to calculate the partial pressures of the various gases contained in this system, one must first calculate the mole fraction of each component. This is done by calculating the number of moles present of each component and dividing that by the total number of moles present in the system. To calculate the partial pressure of each component, the mole fraction must then be multiplied by the total pressure of the system.

(a) To calculate the number of moles present of each component, divide the number of grams present by the molecular weight of the element.

$$\text{number of moles} = \frac{\text{number of grams present}}{\text{molecular weight}}$$

$$\text{Moles of He} = \frac{40 \text{ g}}{4 \text{ g/mole}} = 10 \text{ moles}$$

$$\text{Moles of N}_2 = \frac{56 \text{ g}}{28 \text{ g/mole}} = 2 \text{ moles}$$

$$\text{Moles of O}_2 = \frac{16 \text{ g}}{32 \text{ g/mole}} = 0.5 \text{ moles}$$

(b) To calculate the total number of moles present, add the number of moles of all components together

$$\begin{aligned} \text{total number of moles} &= \text{moles He} + \text{moles N}_2 + \text{moles O}_2 \\ &= 10 + 2 + 0.5 \end{aligned}$$

$$= 12.5 \text{ moles}$$

(c) To calculate the mole fraction of each component, divide the number of moles present of each component by the total number of moles in the system.

$$\text{mole fraction He} = \frac{10}{12.5} = .80$$

$$\text{mole fraction N}_2 = \frac{2}{12.5} = .16$$

$$\text{mole fraction O}_2 = \frac{0.5}{12.5} = .04$$

(d) To find the partial pressure of each component, multiply the mole fraction by the total pressure in the system.

$$\text{partial pressure He} = (.80) \times 5 \text{ atm} = 4 \text{ atm}$$

$$\text{partial pressure N}_2 = .16 \times 5 \text{ atm} = 0.8 \text{ atm}$$

$$\text{partial pressure O}_2 = .04 \times 5 \text{ atm} = 0.2 \text{ atm.}$$

• PROBLEM 84

The composition of dry air by volume is 78.1% N₂, 20.9% O₂ and 1% other gases. Calculate the partial pressures, in atmospheres, in a tank of dry air compressed to 10.0 atmospheres.

Solution: A partial pressure is the individual pressure caused by one gas in a mixture of several gases. The total pressure, P_{total} , according to Dalton's Laws, is the sum of these individual partial pressure, p_1 , p_2 and p_3 .
i.e. $P_{\text{total}} = p_1 + p_2 + p_3$.

The term dry air indicates that no water vapor is present.

The partial pressure is found by multiplying the percent of each gas in the volume by the total pressure.

$$\text{partial pressure} = \text{proportion by volume} \times \text{total pressure}$$

$$P_{\text{N}_2} = .781 \times 10 \text{ atm} = 7.81 \text{ atm}$$


$$P_{\text{O}_2} = .209 \times 10 \text{ atm} = 2.09 \text{ atm}$$

$$P_{\text{other gases}} = .010 \times 10 \text{ atm} = .1 \text{ atm.}$$

The total pressure, as required, is 10 atm.

• PROBLEM 85

A sample of air held in a graduated cylinder over water has a volume of 88.3 ml at a temperature of 18.5°C and a pressure of 741 mm (see figure). What would the volume of the air be if it were dry and at the same temperature and pressure?



Solution: This problem is an application of Dalton's Law which states that the total pressure of a mixture, P_{total} , of gases is the sum of the individual partial pressures, P_1 , P_2 . Stated mathematically

$$P_{\text{total}} = P_1 + P_2 + P_3 \dots$$

The mixture of gases in this problem is located in the cylinder and is made up of dry air and water vapor. At 18.5°C the vapor pressure of water is 16 mm. The pressures are related by the equation

$$P_{\text{total}} = P_{\text{dry air}} + P_{\text{water vapor}}$$

Using this equation, we can calculate $P_{\text{dry air}}$. Having

obtained this value, the problem becomes a direct pressure-volume relationship. As the volume increases, the pressure decreases and vice versa. These factors are related by the equation

$$P_1 V_1 = P_2 V_2$$

where P_1 and V_1 represent pressure and volume in the presence of water vapor, and P_2 and V_2 represent the same quantities for dry air. By Dalton's Law:

$$\begin{aligned} P_{\text{dry air}} &= P_{\text{total}} - P_{\text{water vapor}} \\ &= 741 - 16 \\ &= 725 \text{ mm when water vapor is present.} \end{aligned}$$

If there is no water vapor present and the dry air were to exert the entire pressure of 741 mm then, we predict that the dry air volume will be smaller. Next substitute the values into the pressure-volume equation and solve for unknown volume V_2

$$\begin{aligned} V_2 &= \frac{P_1 V_1}{P_2} \\ &= \frac{725(88.3)}{741} \\ &= 86.4 \text{ ml} \end{aligned}$$

Therefore, as predicted the volume has decreased.

• PROBLEM 86

What would be the final partial pressure of oxygen in the following experiment? A collapsed polyethylene bag of 30 liters capacity is partially blown up by the addition of 10 liters of nitrogen gas measured at 0.965 atm and 298°K. Subsequently, enough oxygen is pumped into the bag so that, at 298°K and external pressure of 0.990 atm, the bag contains a full 30 liters. (assume ideal behavior.)

Solution: One should first find the number of moles of gas that fill a volume of 30 liters at .990 atm. One then solves for the number of moles of N_2 that was pumped in and the number of moles of O_2 that are needed to fill the bag. The partial pressures of these gases are equal to the mole fraction of the gas times the total pressure.

To solve for the partial pressure of O_2 one should find:

- 1) the total number of moles of gas that will fill 30 liters at .990 atm.
- 2) the number of moles of N_2 present
- 3) the partial pressure of N_2
- 4) the partial pressure of O_2 .

Solving:

1) One uses the Ideal Gas Law to solve for the number of moles of gas needed to fill 30 liters at 298°K and .990 atm. The Ideal Gas Law can be stated as

$$PV = nRT \quad \text{or} \quad n = \frac{PV}{RT}$$

where n is the number of moles, P is the pressure, V is the volume, R is the gas constant (.082 liter-atm/mole-°K)

and T is the absolute temperature. Solving for n in this system

$$\begin{aligned}P &= .990 \text{ atm} \\V &= 30 \text{ liters} \\R &= .082 \text{ liter-atm/mole-}^{\circ}\text{K} \\T &= 298^{\circ}\text{K}\end{aligned}$$

$$\begin{aligned}n &= \frac{.990 \text{ atm} \times 30 \text{ liters}}{.082 \text{ liter-atm/mole-}^{\circ}\text{K} \times 298^{\circ}\text{K}} \\&= 1.22 \text{ moles}\end{aligned}$$

2) One also uses the Ideal Gas Law to find the number of moles of N_2 already present.

$$\begin{aligned}P &= .965 \text{ atm} \\V &= 10 \text{ liters} \\R &= .082 \text{ liter-atm/mole-}^{\circ}\text{K} \\T &= 298^{\circ}\text{K}\end{aligned}$$

$$\begin{aligned}n &= \frac{.965 \text{ atm} \times 10 \text{ liters}}{.082 \text{ liter-atm/mole-}^{\circ}\text{K} \times 298^{\circ}\text{K}} \\n &= 0.395 \text{ moles.}\end{aligned}$$

3) The partial pressure is related to the total pressure by the following

$$P_x = P_o \times X$$

where P_x is the partial pressure, P_o is the total original pressure and X is the mole fraction of the gas. The mole fraction is defined as the number of moles of the gas present divided by the total number of moles in the system. Solving for the partial pressure of N_2 , P_{N_2} ,

$$\begin{aligned}P_N + P_o \times X & & P_{N_2} &= ? \\P_o &= .990 \text{ atm} \\X &= .395/1.22\end{aligned}$$

$$P_{N_2} = .990 \text{ atm} \times \frac{.395}{1.22}$$

$$P_{N_2} = .321 \text{ atm.}$$

4) The total pressure of the system is equal to the sum of the partial pressures. Thus, .990 atm is the sum of P_{N_2} and P_{O_2} . One has already found P_{N_2} , therefore, one can now solve for P_{O_2}

$$P_{O_2} = P_{\text{total}} - P_{N_2} \qquad P_{\text{total}} = .990 \text{ atm}$$

$$P_{N_2} = .321 \text{ atm}$$

$$P_{O_2} = .990 \text{ atm} - .321 \text{ atm}$$

$$P_{O_2} = .669 \text{ atm.}$$

• PROBLEM 87

200 ml of oxygen is collected over water at 25°C and 750 torr. If the oxygen thus obtained is dried at a constant temperature of 25°C and 750 torr, what volume will it occupy? What volume will be occupied by the water vapor removed from the oxygen if it is maintained at 25°C and 750 torr? (The equilibrium vapor pressure of water at 25°C is 28.3 torr.)

Solution: This problem will be approached by determining the partial pressures of oxygen and of water vapor in the initial mixture and then using these partial pressures to determine the separate volumes of each component.

The partial pressures exerted by the oxygen and the water vapor must add up to the total pressure. Since the partial pressure of water vapor is equal to its equilibrium vapor pressure, we can determine the partial pressure of oxygen:

$$\text{total pressure} = \text{partial pressure of } O_2 + \text{partial pressure of water}$$

$$750 \text{ torr} = \text{partial pressure of } O_2 + 28.3 \text{ torr}$$

$$\text{or, } \text{partial pressure of } O_2 = 750 \text{ torr} - 28.3 \text{ torr} = 721.7 \text{ torr.}$$

By definition, this partial pressure is the pressure the oxygen would exert if it filled the entire 200 ml volume. We must determine the volume this amount of oxygen would fill at 750 torr. To do this we use the relationship $P_1V_1 = P_2V_2$, which is valid for constant temperature and constant number of moles of gas. Let $P_1 = 721.7 \text{ torr}$, $V_1 = 200 \text{ ml}$, and $P_2 = 750 \text{ torr}$. Then,

$$V_2 = \frac{P_1}{P_2} \times V_1 = \frac{721.7 \text{ torr}}{750 \text{ torr}} \times 200 \text{ ml} = 192.45 \text{ ml.}$$

Hence, the dried oxygen would occupy a volume of 192.45 ml.

We follow a similar procedure in calculating the volume that the water vapor would occupy at 750 torr. The vapor pressure of water, 28.3 torr, is the pressure the water vapor would exert if it filled the entire 200 ml volume. Again, applying the relationship $P_1V_1 = P_2V_2$, remembering that the only reason we can do so is that we

have constant temperature and constant number of moles of gas, we can calculate the volume that the water vapor occupies under a pressure of 750 torr. Let $P_1 = 28.3$ torr, $V_1 = 200$ ml, and $P_2 = 750$ torr. Then the volume at 750 torr is

$$V_2 = \frac{P_1}{P_2} \times V_1 = \frac{28.3 \text{ torr}}{750 \text{ torr}} \times 200 \text{ ml} = 7.55 \text{ ml}.$$

As a check, we ascertain that the individual volumes at 750 torr add up to the total volume at 750 torr:

$$\begin{aligned} \text{Volume of O}_2 + \text{volume of water} &= 192.45 \text{ ml} + 7.55 \text{ ml} \\ &= 200 \text{ ml}. \end{aligned}$$

• PROBLEM 88

Suppose 100 ml of oxygen were collected over water in the laboratory at a pressure of 700 torr and a temperature of 20°C. What would the volume of the dry oxygen gas be at STP?

Solution: STP means Standard Temperature and Pressure, which is 0°C and 760 torr. In this problem oxygen is gathered over water, therefore, Dalton's law of partial pressure (each of the gases in a gaseous mixture behaves independently of the other gases and exerts its own pressure, the total pressure of the mixture being the sum of the partial pressures exerted by each gas present; that is $P_{\text{total}} = p_1 + p_2 + p_3 \dots p_n$) is used to calculate the original pressure of the oxygen. There is both water vapor and oxygen gas present.

After you obtain the original pressure of the oxygen, you can use the combined gas law to calculate the final volume of the oxygen. The combined gas law stated that for a given mass of gas, the volume is inversely proportional to the pressure and directly proportional to the absolute temperature. It can be written as follows:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where P_1 is the original pressure, V_1 is the original volume, T_1 is the original absolute temperature, P_2 is the final pressure, V_2 is the final volume, and T_2 is the final absolute temperature. You are given the temperature in °C, so it must be converted to the absolute temperature by adding 273.

(a) Finding the original pressure of oxygen. The partial pressure of water is 17.5 torr.

$$P_{\text{total}} = 700 \text{ torr} = P_{\text{O}_2} + P_{\text{H}_2\text{O}}$$

$$P_{O_2} = 700 - 17.5$$

$$= 682.5 \text{ torr.}$$

(b) Converting the temperature to the absolute scale.

$$T_1 = 20 + 273 = 293^\circ\text{K}$$

$$T_2 = 0 + 273 = 273^\circ\text{K}$$

(c) Using the combined law

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P_1 = 682.5 \text{ torr}$$

$$P_2 = 760 \text{ torr}$$

$$V_1 = 100 \text{ ml}$$

$$V_2 = ?$$

$$T_1 = 293^\circ\text{K}$$

$$T_2 = 273^\circ\text{K}$$

$$\frac{(682.5 \text{ torr})(100 \text{ ml})}{293^\circ\text{K}} = \frac{(760 \text{ torr}) V_2}{273^\circ\text{K}}$$

$$V_2 = \frac{(682.5 \text{ torr})(100 \text{ ml})(273^\circ\text{K})}{(760 \text{ torr})(293^\circ\text{K})} = 83.7 \text{ ml.}$$

• PROBLEM 89

If 40 liters of N_2 gas are collected at 22°C over water at a pressure of .957 atm, what is the volume of dry nitrogen at STP (Standard temperature and pressure)?

Solution: The key to this problem is the recognition that $PV/T = k$, where P is the pressure, V the volume, T the absolute temperature and k a constant. If you let the STP conditions (1-atm. and 0°C), be represented by $P_1 V_1 / T_1$ and the conditions of the gas collected over the water by $P_2 V_2 / T_2$, then

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} ,$$

since both equal the same constant, k . Therefore, to answer the question, you have to substitute the known values into the equation and solve for the unknown, V_1 .


However, there is one precaution. Over water, you have vapor pressure. At 22°C , water has a vapor pressure of .026 atm. As such, this makes the initial pressure of N_2 .957 - .026, or .931 atm. Add 273° to the Celsius temperature to obtain the temperature in Kelvin. Therefore, $T_2 = 22^\circ\text{C} + 273^\circ = 295^\circ\text{K}$ and $T_1 = 0^\circ\text{C} + 273 = 273^\circ\text{K}$. Therefore, by substitution

$$\frac{(1)(V_1)}{273} = \frac{(.931)(40)}{295}$$

Now, solving for V_1 , the volume at STP, you obtain 34.4 liters.

• PROBLEM 90

A sample of hydrogen is collected in a bottle over water. By carefully raising and lowering the bottle, the height of the water outside is adjusted so that it is just even with the water level inside (see figure). When a sample of gas was collected the initial conditions were: volume = 425 ml, pressure = 753 mm and the temperature of the water (and thus, the gas also) = 34°C . Calculate the volume of the hydrogen if it were dry and at a pressure of 760 mm and a temperature of 0°C (STP).



Solution: First, apply Dalton's Law and then make use of the Combined Gas Law to obtain the desired answer.

Dalton's Law states that the total pressure of a mixture of gases, P_{total} , is equal to the sum of the individual partial pressures, P_{H_2} and $P_{\text{H}_2\text{O}}$. In this example the individual gases are hydrogen and water vapor. Therefore, the total pressure is the sum of these two gases. At 35°C , the partial pressure of H_2O vapor is 40 mm,

$$P_{\text{total}} = P_{\text{H}_2} + P_{\text{H}_2\text{O}}$$

$$P_{\text{H}_2} = P_{\text{total}} - P_{\text{H}_2\text{O}}$$

$$= 753 - 40 = 713 \text{ mm}$$

This means that if no water vapor was present the H_2 gas would fill the container at 713 mm of pressure.

Since pressure is indirectly proportional to the volume of H_2 gas, the effect of changing the pressure of

713 mm to STP pressure (760 mm) is to decrease the volume.

In addition, however, the temperature of the H_2 gas is directly proportional to the volume of the gas, so that the effect of changing the temperature of ($34^\circ C + 273^\circ =$) $307^\circ K$ to STP temperature ($273^\circ K$) is also to decrease the volume of the gas. Therefore, we can predict that the net effect of changing the pressure and temperature of the gas to STP conditions is to decrease the volume.

Using the combined gas law equation to solve for the dry H_2 gas volume, V_2 ,

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

where $P_1 = 713 \text{ mm}$

$P_2 = 760 \text{ mm}$

$V_1 = 425 \text{ ml}$

$V_2 = \text{volume at STP}$

$T_1 = 307^\circ K$

$T_2 = 273^\circ K$

Thus, substituting,

$$\begin{aligned} V_2 &= \frac{P_1 V_1 T_2}{P_2 T_1} \\ &= \frac{(713 \text{ mm})(425 \text{ ml})(273^\circ K)}{(760 \text{ mm})(307^\circ K)} = 355 \text{ ml.} \end{aligned}$$

We see that this answer is in total agreement with our prediction that the volume at STP, $V_2 = 355 \text{ ml}$, is less than the volume at the initial conditions, $V_1 = 425 \text{ ml}$.

• PROBLEM 91

A 20 g chunk of Dry Ice (CO_2) is placed in an "empty" 0.75 liter wine bottle and tightly corked. What would be the final pressure in the bottle after all the CO_2 has evaporated and the temperature has reached $25^\circ C$?

Solution: The final pressure in the bottle will equal the original pressure plus the pressure contributed by the CO_2 . Since the bottle was originally open in atmospheric pressure, the original pressure is 1 atm. The pressure contributed by the CO_2 is found by using the Ideal Gas Law. This law is stated

$$PV = nRT \quad \text{or} \quad P = \frac{nRT}{V}$$

where P is the pressure of the gas, n is the number of moles, R is the gas constant, 0.082 liter-atm/mole $^\circ K$, V is the volume, and T is the absolute temperature. One

is given R and V in the problem. n is found by dividing the number of grams of CO_2 present by the molecular weight (MW of $\text{CO}_2 = 44$).

$$\text{no. of moles} = \frac{20 \text{ g}}{44 \text{ g/mole}} = .45 \text{ moles}$$

One can convert $^{\circ}\text{C}$ to $^{\circ}\text{K}$ by adding 273° to the temperature in $^{\circ}\text{C}$.

$$^{\circ}\text{K} = ^{\circ}\text{C} + 273^{\circ}$$

$$T = 25 + 273^{\circ} = 298^{\circ}\text{K}$$

Solving for P_{CO_2} :

$$P_{\text{CO}_2} = \frac{.45 \text{ moles} \times .082 \text{ liter} \cdot \text{atm} / \text{mole} \cdot ^{\circ}\text{K} \times 298^{\circ}\text{K}}{.75 \text{ liter}}$$


$$= 14.81 \text{ atm.}$$

$$P_{\text{total}} = P_{\text{CO}_2} + P_{\text{original}}$$

$$P_{\text{total}} = 14.81 \text{ atm} + 1 \text{ atm} = 15.81 \text{ atm.}$$

• PROBLEM 92

The volume of hydrogen evolved during the course of a reaction is measured by the displacement of water as shown in the diagram below. Hydrogen is evolved in flask A and displaces water from flask B into beaker C. If, during a particular run of this experiment in which atmospheric pressure is 765 torr and the water temperature is 293.15°K , 65.0 ml of water is displaced, how much water would be displaced at 760 torr and 298.15°K ? (The equilibrium vapor pressure of water at 293.15°K is 17.5 torr and at 298.15°K is 23.8 torr.)



Solution: We can convert from the volume under non-standard conditions in the problem to the volume under standard conditions by means of the ideal gas equation, $PV = nRT$ (where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature). For our nonstandard conditions we may write

$$P_n V_n = nRT_n$$

and for final conditions (760 torr and 298.15°K) we may write

$$P_s V_s = nRT_s.$$

Note that n and R are the same in both cases. Dividing the second equation by the first we obtain

$$\frac{P_s V_s}{P_n V_n} = \frac{nRT_s}{nRT_n} = \frac{T_s}{T_n} \quad \text{or} \quad V_s = \frac{P_n V_n}{P_s} \times \frac{T_s}{T_n}.$$

The pressure experienced by the hydrogen under non-standard conditions is equal to the pressure exerted by the atmosphere (765 torr) plus that from water vapor pressure at 20°C (17.5 torr), or $P_n = 765 \text{ torr} + 17.5 \text{ torr} = 782.5 \text{ torr}$. Under the final conditions, the pressure experienced by hydrogen is equal to the sum of the atmospheric pressure (760 torr) and the vapor pressure of water at 298.15°K (23.8 torr), or

$$P_s = 760 \text{ torr} + 23.8 \text{ torr} = 783.8 \text{ torr}.$$

We can now calculate the volume of hydrogen under the final conditions. Thus,

$$V_s = \frac{P_n V_n}{P_s} \times \frac{T_s}{T_n} = \frac{765 \text{ torr} \times 65.0 \text{ ml}}{783.8 \text{ torr}} \times \frac{298.15^\circ\text{K}}{293.15^\circ\text{K}} = 64.5 \text{ ml}.$$

• PROBLEM 93

A technician is working with dry ice (solid CO_2) in a closed 6000 liter fume hood maintained at a constant pressure of 1.00 atm and a constant temperature of 27°C. If he has not been conditioned to tolerate CO_2 , he will succumb to CO_2 poisoning when the concentration rises to 10%, by volume, of the atmosphere. If the ventilation stops, what minimum weight of dry ice must sublime to constitute a hazard? Assume that there was no CO_2 present initially.

Solution: To solve this problem we will convert the concentration, 10% by volume, to a partial pressure and then determine what mass of CO_2 gives rise to this pressure.

At the point where CO_2 is harmful, it occupies 10% by volume of the total atmosphere. Its partial pressure p is therefore 10% of the total pressure, or $p = 10\% \times 1 \text{ atm} = 0.10 \text{ atm}$.

The ideal gas law reads

$$pV = nRT = \frac{m}{MW} RT$$

where V is the volume, n the number of moles, R the gas constant, T the absolute temperature, m the mass, and MW the molecular weight, and we have used $n = m/MW$. Solving for m

$$m = \frac{pV(MW)}{RT}$$

Now $p = 0.10$ atm, $V = 6000$ liters, $MW = 44$ g/mole of CO_2 , $R = 0.082$ liter-atm/mole-degree, and $T = 27^\circ\text{C} = 300^\circ\text{K}$. Hence,

$$m = \frac{pV(MW)}{RT} = \frac{0.10 \text{ atm} \times 6000 \text{ liters} \times 44 \text{ g/mole}}{0.082 \text{ liter-atm/mole-deg} \times 300^\circ\text{K}}$$

$$\approx 1100 \text{ g.}$$

Thus, 1100 g, or 1.10 kg, must sublime before there is sufficient CO_2 in the atmosphere to be harmful.

GRAHAM'S LAW OF GASEOUS DIFFUSION

• PROBLEM 94

Under standard temperature and pressure conditions, compare the relative rates at which inert gases, Ar, He, and Kr diffuse through a common orifice.

Solution: This problem involves the application of Graham's law of diffusion. It states that the relative rates at which gases will diffuse will be inversely proportional to the square roots of their respective densities or molecular weights. That is, rate $\propto 1/\sqrt{\text{mass}}$. Thus, to compare the rates of diffusion of Ar, He, and Kr, look up their weights in the Periodic Table of Elements and substitute this value into $1/\sqrt{M} \propto \mu_m$, where M = mass of that element and μ_m = rate. Therefore, $\mu_{\text{Ar}} : \mu_{\text{He}} : \mu_{\text{Kr}}$

$$= \frac{1}{\sqrt{M_{\text{Ar}}}} : \frac{1}{\sqrt{M_{\text{He}}}} : \frac{1}{\sqrt{M_{\text{Kr}}}}$$

$$= \frac{1}{\sqrt{39.95}} : \frac{1}{\sqrt{4.003}} : \frac{1}{\sqrt{83.80}}$$

$$= .1582 : .4998 : .1092.$$

Two gases, HBr and CH₄, have molecular weights 81 and 16, respectively. The HBr effuses through a certain small opening at the rate of 4 ml/sec. At what rate will the CH₄ effuse through the same opening?

Solution: The comparative rates or speeds of effusion of gases are inversely proportional to the square roots of their molecular weights. This is written

$$\frac{\text{rate}_1}{\text{rate}_2} = \frac{\sqrt{\text{MW}_2}}{\sqrt{\text{MW}_1}}$$

For this case $\frac{\text{rate}_{\text{HBr}}}{\text{rate}_{\text{CH}_4}} = \frac{\sqrt{\text{MW}_{\text{CH}_4}}}{\sqrt{\text{MW}_{\text{HBr}}}}$

One is given the rate_{HBr}, MW_{CH₄}, and MW_{HBr} and asked to find rate_{CH₄}.

Solving for rate_{CH₄}:

$$\frac{\text{rate}_{\text{HBr}}}{\text{rate}_{\text{CH}_4}} = \frac{\sqrt{\text{MW}_{\text{CH}_4}}}{\sqrt{\text{MW}_{\text{HBr}}}}$$

$$\text{rate}_{\text{HBr}} = 4 \text{ ml/sec.}$$

$$\text{rate}_{\text{CH}_4} = ?$$

$$\text{MW}_{\text{CH}_4} = 16$$

$$\text{MW}_{\text{HBr}} = 81$$

$$\frac{4 \text{ ml/sec}}{\text{rate}_{\text{CH}_4}} = \frac{\sqrt{16}}{\sqrt{81}}$$

$$\text{rate}_{\text{CH}_4} = \frac{4 \text{ ml/sec} \times \sqrt{81}}{\sqrt{16}}$$

$$= \frac{4 \text{ ml/sec} \times 9}{4} = 9 \text{ ml/sec.}$$

The time required for a volume of gas, X, to effuse through a small hole was 112.2 sec. The time required for the same volume of oxygen was 84.7 sec. Calculate the molecular weight of gas X.

Solution: This problem involves the rate of effusion through a small hole and the method to solve it involves a modified statement of Graham's Law. High density gases effuse more slowly than low density gases. The time required for effusion is inversely proportional to the rate of effusion. It is mathematically stated as

$$\frac{r_1}{r_2} = \frac{t_2}{t_1} = \frac{\sqrt{M.W._2}}{\sqrt{M.W._1}},$$

where, r_1 , r_2 are the rates of the two gases; t_2 , t_1 are the times; and $M.W._2$ and $M.W._1$, = molecular weights.

Knowing 3 of the 4 values, one can use this equation to find the 4th value. We can write

$$\frac{t_{O_2}}{t_X} = \frac{\sqrt{M.W._{O_2}}}{\sqrt{M.W._X}}$$

$$\frac{84.7 \text{ sec}}{112.2 \text{ sec}} = \frac{\sqrt{32}}{\sqrt{X}}$$

$$M.W._X = 56.2 \text{ amu.}$$

Molecular Weight is expressed in terms of a.m.u. (atomic mass units).

• PROBLEM 97

At standard conditions, 1 liter of oxygen gas weighs almost 1.44 g, whereas 1 liter of hydrogen weighs only .09 g. Which gas diffuses faster? Calculate how much faster.

Solution: This question deals with diffusion of gases, which is governed by Graham's law. A gas that has a high density diffuses more slowly than one with a lower density. The rates of diffusion of 2 gases are inversely proportional to the square roots of their densities, as shown by the equation

$$\frac{r_1}{r_2} = \frac{\sqrt{d_2}}{\sqrt{d_1}}, \text{ where } r_1, r_2 = \text{rates of diffusion of}$$

the gases, and d_1 , d_2 are their respective densities.

Therefore, by Graham's law, we know that H_2 diffuses faster because of its lower density. (H_2 's density = $\frac{M}{V} = \frac{.09 \text{ g}}{1 \text{ l}} = .09 \text{ g/l}$ while O_2 's density = $\frac{1.44}{1} = 1.44 \text{ g/liter.}$)

To find out how much faster H_2 diffuses, plug the necessary factors into the equation

$$\frac{r_{H_2}}{r_{O_2}} = \frac{\sqrt{d_{O_2}}}{\sqrt{d_{H_2}}} \quad \text{Rewriting,}$$

$$r_{H_2} = r_{O_2} \frac{\sqrt{d_{O_2}}}{\sqrt{d_{H_2}}} = r_{O_2} \frac{\sqrt{1.44}}{\sqrt{.09}}$$

$$r_{H_2} = 4r_{O_2}$$

Thus, the calculated comparison is in agreement with the predicted comparison that H_2 diffuses more quickly. To be precise, the rate of diffusion of H_2 is 4 times that of the rate of diffusion of O_2 .

• PROBLEM 98

The relative rates of diffusion of $NH_3(g)$ and $HX(g)$ can be determined experimentally by simultaneously injecting $NH_3(g)$ and $HX(g)$, respectively, into the opposite ends of a glass tube and noting where a deposit of $NH_4X(s)$ is formed. Given an 1m tube, how far from the NH_3 injection end would you expect the $NH_4X(s)$ to be formed when HX is a) HF , b) HCl ?

Solution: The rate of diffusion of a gas is observed to be inversely proportional to the square root of its molecular weight. This is called Graham's law.

$$\text{Rate} = \frac{\text{constant}}{\sqrt{MW}}$$

For HF : According to Graham's Law:

the speed of NH_3 is proportional to $\frac{1}{\sqrt{17}} = .243$
(MW of NH_3 = 17)

the speed of HF is proportional to $\frac{1}{\sqrt{20}} = .224$
(MW of HF = 20)

Thus, for an arbitrary amount of time, let us say 1 min., NH_3 moves .243 cm and HF moves .224 cm. Because the gases are ejected from either end of the tube, after 1 minute the gases move (.243 + .224) cm or .467 cm/min. closer together. When this distance is equal to 100 cm the solid forms. One can determine the time needed by dividing 100 cm by .467 cm/min

$$\text{time after which solid forms} = \frac{100 \text{ cm}}{.467 \text{ cm/min}} = 214.13 \text{ min.}$$

Thus, after 214.13 min. the HF meets the NH_3 .

The distance the NH_3 would travel is found by multiplying the distance it travels in 1 minute, .243 cm, by the time elapsed till the NH_4F deposits.

$$\text{distance } NH_3 \text{ travels} = .243 \text{ cm/min} \times 214.13 \text{ min}$$

$$= 52.03 \text{ cm.}$$

b) One solves for HCl in a similar manner

$$\text{MW of HCl} = 36.5$$

$$\text{the speed of HCl is proportional to } \frac{1}{\sqrt{36.5}} = .165$$

$$\text{As previously found, the speed of NH}_3 \text{ is proportional to } \frac{1}{\sqrt{17}} = .243.$$

Speed is a measure of distance covered over a unit time. Let us designate the speeds here as cm./min. Thus, HCl moves .165 cm/min and NH₃ moves .243 cm/min. The solid forms after the sum of the distances travelled by the two gases is 100 cm. In 1 min. the gases travel a distance of (.165 + .243)cm or .408 cm. The time it takes for the gases to travel 100 cm. is found by dividing 100 cm. by .408 cm/min.

time after which the solid will deposit =

$$= \frac{100 \text{ cm}}{.408 \text{ cm/min}} = 245.10 \text{ min.}$$

The distance the NH₃ travels is then found by multiplying 245.10 min by the distance the gas travels in 1 minute.

$$\begin{aligned} \text{distance travelled by NH}_3 &= .243 \text{ cm/min} \times 245.10 \text{ min} \\ &= 59.56 \text{ cm.} \end{aligned}$$

• PROBLEM 99

Two balloons at the same temperature of equal volume and porosity are each filled to a pressure of 4 atmospheres, one with 16 kg of oxygen, the other with 1 kg of hydrogen. The oxygen balloon leaks to a pressure of $\frac{1}{4}$ atmosphere (atm) in 1 hour. How long will it take for the hydrogen balloon to reach a pressure of $\frac{1}{4}$ atm?

Solution: This problem deals with effusion, the escape of molecules through a hole, and is therefore an application of Graham's Law: Rate of effusion of a gas is inversely proportional to the square root of its density

$$\frac{\text{rate}_1}{\text{rate}_2} = \frac{\sqrt{d_2}}{\sqrt{d_1}}$$

From this we know that large molecules effuse less rapidly than small molecules, since they move more slowly.

This allows us to predict that hydrogen effuses more rapidly than oxygen. By applying Graham's Law, we can determine how much faster.

$$\frac{\text{rate}_{\text{hydrogen}}}{\text{rate}_{\text{oxygen}}} = \frac{\sqrt{16 \text{ kg per unit volume}}}{\sqrt{1 \text{ kg per unit volume}}} = 4.$$


There, hydrogen effuses 4 times more quickly, which is in agreement with our prediction.

Since it takes 1 hour for oxygen to leak to a pressure of $\frac{1}{4}$ atm, it requires $\frac{1}{4}$ hr. = 15 min for hydrogen to decrease to the same pressure inside the balloon.

KINETIC THEORY OF GASES

• PROBLEM 100

Calculate the most probable speed v_p , the arithmetic mean speed \bar{v} , and the root mean square speed v_{rms} for hydrogen molecules at 0°C .



Probability density of various speeds v for oxygen at 100, 300, 500, and 1000 K.

Solution: Before beginning this problem, we must first understand the basic differences in the three types of speeds. The arithmetic mean speed \bar{v} is obtained by summing all the speeds and dividing by the total number of molecules N .

$$\bar{v} = \frac{1}{N} \sum_{i=1}^N v_i$$

The symbol $\sum v_i$ indicates the sum of the speeds $v_1, v_2, v_3, \dots, v_N$ of all the N individual molecules. The equation for mean speed is

$$\bar{v} = \left(\frac{8kT}{\pi m} \right)^{\frac{1}{2}} = \left(\frac{8RT}{\pi M} \right)^{\frac{1}{2}}$$

where k is Boltzmann's constant, R is the gas constant, T is the absolute temperature, m is the weight of 1 molecule, and M is the molecular weight.

The most probable speed is obtained from the plot of $f(v)$ versus v , where $f(v)$ is the fraction of molecules with speed v . The equation of this curve

$$f(v) = \left(\frac{m}{2\pi kT} \right)^{3/2} e^{-mv^2/2kT} 4\pi v^2$$

is differentiated with respect to v and set equal to zero

$\left(\frac{df(v)}{dv} = 0 \right)$. The shape of the curve of $f(v)$ versus v is indicated in the accompanying figure.

$$\frac{df(v)}{dv} = \left(\frac{m}{2\pi kT} \right)^{3/2} e^{-mv^2/2kT} \left(8\pi v + 4\pi v^2 \left(\frac{-mv}{kT} \right) \right)$$

This derivative vanishes at $v = v_p$, called the most probable speed, where

$$v_p = \left(\frac{2kT}{m} \right)^{\frac{1}{2}} = \left(\frac{2RT}{M} \right)^{\frac{1}{2}}$$

The root-mean-square speed, v_{rms} , is defined by

$$v_{rms} = \left(\frac{1}{N} \sum_{i=1}^N v_i^2 \right)^{\frac{1}{2}}$$

The symbol $\sum v_i^2$ indicates the sum of the squares of the velocities, $v_1, v_2, v_3, \dots, v_N$ of all the N individual molecules.

Since the velocity distribution is continuous, the root-mean-square velocity is obtained by multiplying each velocity squared by the probability of that velocity, integrating over all velocities, and taking the square root:

$$v_{\text{rms}} = \left(\int_0^{\infty} v^2 f(v) dv \right)^{\frac{1}{2}}$$

Substituting for $f(v)$ and simplifying one obtains

$$v_{\text{rms}} = \left(\frac{3kT}{m} \right)^{\frac{1}{2}} = \left(\frac{3RT}{M} \right)^{\frac{1}{2}}$$

At any given temperature, these velocities are inversely proportional to the square root of the molecular weight. Lighter molecules move more rapidly so that their average kinetic energies are exactly equal to those for the heavier molecules.

To solve this problem for hydrogen at 0°C one must 1) find the molecular weight of hydrogen in kg, 2) convert 0°C into the absolute temperature scale, 3) substitute these values into the equations for v_p , \bar{v} , and v_{rms} .

The molecular weight of hydrogen is 2.016 g/mole, which is 2.016×10^{-3} kg/mole. The temperature in degrees Kelvin is 273°K. Other pertinent information $R = 8.314$ J/K-mole, $\pi = 3.142$. Thus,

$$\begin{aligned} v_p &= \left(\frac{2RT}{M} \right)^{\frac{1}{2}} \\ &= [(2)(8.314 \text{ J/K-mol})(273^\circ\text{K}) / (2.016 \times 10^{-3} \text{ Kg/mol})]^{\frac{1}{2}} \\ &= 1.50 \times 10^3 \text{ m/s} \end{aligned}$$

$$\begin{aligned} \bar{v} &= \left(\frac{8RT}{\pi M} \right)^{\frac{1}{2}} \\ &= [(8)(8.314 \text{ J/K-mol})(273^\circ\text{K}) / (3.142)(2.016 \times 10^{-3} \text{ Kg/mol})]^{\frac{1}{2}} \\ &= 1.69 \times 10^3 \text{ m/s.} \end{aligned}$$

$$\begin{aligned} v_{\text{rms}} &= \left(\frac{3RT}{M} \right)^{\frac{1}{2}} \\ &= [(3)(8.314 \text{ J/K-mol})(273^\circ\text{K}) / (2.016 \times 10^{-3} \text{ Kg/mol})]^{\frac{1}{2}} \\ &= 1.84 \times 10^3 \text{ m/s.} \end{aligned}$$

The relative ratios of $v_p : \bar{v} : v_{\text{rms}}$ are 1 : 1.13 : 1.23.

• PROBLEM 101

Given that the average speed of an O_2 molecule is 1,700 km/h at $0^\circ C$, what would you expect the average speed of a CO_2 molecule to be at the same temperature?

Solution: The speed of a molecule is related to the mass and temperature by the equation

$$\frac{1}{2} m v^2 = \frac{3}{2} k T$$

where m is the mass, v is the speed, k is Boltzmann's constant, and T is the absolute temperature. Since k and T are constant $\frac{3}{2} k T$ is constant for this system. Thus

$$\frac{1}{2} m_{O_2} v_{O_2}^2 = \frac{1}{2} m_{CO_2} v_{CO_2}^2$$

where m_{O_2} is the mass of O_2 , v_{O_2} is the speed of O_2 , m_{CO_2} is the mass of CO_2 , and v_{CO_2} is the speed of CO_2 . The masses are equal to the molecular weights here. Solving for v_{CO_2} :

$$v_{CO_2} = \sqrt{\frac{\frac{1}{2} m_{O_2} v_{O_2}^2}{\frac{1}{2} m_{CO_2}}} \quad \begin{array}{l} m_{O_2} = 32 \\ m_{CO_2} = 44 \\ v_{O_2} = 1700 \text{ km/h.} \end{array}$$

$$v_{CO_2} = \sqrt{\frac{\frac{1}{2} (32) (1700)^2}{\frac{1}{2} (44)}}$$

$$= 1700 \sqrt{\frac{32}{44}} = 1700 \times (.85)$$

$$= 1450 \text{ km/h.}$$

• PROBLEM 102

A chemist possesses 1 cm^3 of O_2 gas and 1 cm^3 of N_2 gas both at Standard Temperature and Pressure (STP). Compare these gases with respect to (a) number of molecules, and (b) average speed of molecules.

Solution: This problem requires the use of the kinetic theory of gases and Avogadro's principle.

Avogadro proposed the principle that equal volumes of gases at the same temperature and pressure contain equal numbers of molecules. Both gases have equal volumes, 1 cm^3 , and are at the same temperature and pressure (STP). According to Avogadro's Principle, both have the same

number of molecules. The average speed of these molecules can be found from kinetic theory of gases. The rate of diffusion of a gas is directly proportional to the average speed of the molecules. According to Graham's law, there is an inverse proportionality between diffusion rate and the square root of the molecular weight (mass). In other words,

$$\frac{\text{rate of diffusion of } N_2}{\text{rate of diffusion of } O_2} = \frac{V_{N_2}}{V_{O_2}} = \sqrt{\frac{M_{O_2}}{M_{N_2}}},$$

where V = velocity and M = mass, $M_{N_2} = 28$, $M_{O_2} = 32$. In this case, mass = molecular weight. Thus,

$$\frac{V_{N_2}}{V_{O_2}} = \sqrt{\frac{32}{28}} = 1.07.$$

Thus, if the average speed of O_2 is one and the average speed of N_2 is 1.07.

• PROBLEM 103

Calculate (a) the number of collisions per second per molecule, (b) the number of collisions per cubic meter per second, and (c) the number of moles of collisions per liter per second of oxygen at a temperature of 25°C and pressure of 1 atm. The molecular diameter of oxygen is 3.61×10^{-10} m.




Figure A

Solution: Molecules of real gas attract one another at large distances and repel one another at very short distances. Assuming that molecules are rigid, noninteracting spheres with diameter, σ , one can derive an approximate equation for the number of collisions per second. It is also assumed that all the molecules travel with the same speed, the arithmetic mean velocity \bar{v} .

If two identical molecules just touch each other, then the distance separating their centers is the molecular diameter σ . Thus, a moving molecule collides with other molecules whose centers come within a distance of σ . (Hence, the effective collision radius is equal to σ , see figure a.) The quantity $\pi\sigma^2$, is called the collision cross section for the rigid spherical molecule. A molecule moving with a constant velocity of \bar{v} will sweep out $(\pi\sigma^2\bar{v})$ (meters)³/sec and strike $\pi\sigma^2\bar{v}n$ molecules per second, where n is the concentration of molecules. In these calculations, we have assumed several things some of which are not completely accurate. Actually the number of collisions per second is off by a factor of $\sqrt{2}$. Thus, the number of collisions per molecule per second is:

$$z = \sqrt{2} \pi\sigma^2\bar{v}n$$

Since there are n molecules per unit volume, there will be $\sqrt{2} \pi\sigma^2\bar{v}n^2$ collisions per unit time. The number of collisions per unit time per unit volume is given by

$$Z = \frac{\sqrt{2}}{2} \pi\sigma^2\bar{v}n^2$$

The factor of one-half comes from the fact that the number of collisions is one-half the number of total collisions. This is true because the number of total collisions is equal to the sum of all of the collisions of all of the particles, but in each collision two molecules are involved which means that in the total number each collision is really counted twice.

To solve the problem we must: 1) find the arithmetic mean speed, 2) find the concentration of molecules, n , using the ideal gas law, 3) substitute the values in the equations for $z = \sqrt{2} \pi\sigma^2\bar{v}n$ and $Z = \frac{\sqrt{2}}{2} \pi\sigma^2\bar{v}n^2$. The information known:

$$\sigma = 3.61 \times 10^{-10} \text{ m} \quad P = 1 \text{ atm}$$

$$T = 25^\circ\text{C} + 273^\circ = 298^\circ\text{K}; M = 32 \times 10^{-3} \text{ Kg/mole} = \text{molecular weight of oxygen}$$

$$R = 0.082 \frac{\text{liter} \cdot \text{atm}}{\text{K} \cdot \text{mole}} = 8.312 \text{ J/K} \cdot \text{mole}.$$

$$\begin{aligned} 1) \quad \bar{v} &= \left(\frac{8RT}{\pi M} \right)^{\frac{1}{2}} \\ &= [(8)(8.314 \text{ J/K} \cdot \text{mole})(298) / \pi(32 \times 10^{-3} \text{ Kg/mol})]^{\frac{1}{2}} \\ &= 444 \text{ m/s.} \end{aligned}$$

$$2) \quad n = \frac{N}{V} = \frac{\text{number of molecules}}{\text{m}^3} = \text{concentration}$$

According to the ideal gas law, $PV = \bar{N}RT$, where \bar{N} = moles or $\frac{\bar{N}}{V} = \frac{P}{RT}$. Avogadro's number, $N_A = 6.02 \times$

$10^{23} \frac{\text{molecules}}{\text{mole}}$, so that $\frac{N}{V} = \frac{PNa}{RT}$. But $N/V = n$ so that

$$\frac{N}{V} = n = \frac{PNa}{RT} = \frac{(1 \text{ atm}) \left(6.022 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \right) \left(10^{-3} \frac{\text{liter}}{\text{m}^3} \right)}{\left(0.082 \frac{\text{liter atm}}{\text{K-mole}} \right) (298 \text{ K})}$$

$$= 2.46 \times 10^{25} \frac{\text{molecules}}{\text{m}^3}$$

3) $z = \sqrt{2} \pi \sigma^2 \bar{v} n$

$$= (1.414) (3.14) (3.6 \times 10^{-10} \text{ m})^2 (444 \text{ m/s}) \left(2.46 \times 10^{25} \frac{\text{molecules}}{\text{m}^3} \right)$$

$$= 6.32 \times 10^9 \frac{\text{molecules}}{\text{s}}$$

This is the number of collisions per molecule per second.

$$Z = \frac{\sqrt{2}}{2} \pi \sigma^2 \bar{v} n^2$$

$$= \frac{z n}{2} = \frac{\left(6.32 \times 10^9 \frac{\text{molecules}}{\text{s}} \right) \left(2.46 \times 10^{25} \frac{\text{molecules}}{\text{m}^3} \right)}{2}$$

$$= 7.77 \times 10^{34} \frac{\text{collisions}}{\text{m}^3 \text{ s}}$$

To answer part c of the problem, we use the conversion factor $\frac{10^{-3} \text{ m}^3/\text{liter}}{6.02 \times 10^{23} \text{ molecules/mole}}$. Thus

$$Z = \frac{(7.77 \times 10^{34} \text{ collisions/m}^3\text{-s}) (10^{-3} \text{ m}^3/\text{l})}{(6.022 \times 10^{23} \text{ molecules/mole})}$$

$$= 1.29 \times 10^8 \text{ moles/liter-s,}$$

which is the number of moles of collisions per liter per second.

• PROBLEM 104

Calculate the mean free path for oxygen at 25°C at (a) 1 atm pressure and (b) 10^{-1} torr.

Solution: The mean free path is the average distance traversed by a molecule between collisions. In a second, a molecule will, on the average, traverse \bar{v} meters and collide Z times (where $Z = \sqrt{2} \pi \sigma^2 \bar{v} n$). Thus, the mean free path is

$$\lambda = \frac{v}{z} = \frac{1}{\sqrt{2} \pi \sigma^2 n}$$

Substituting $n = P/kT$ from the ideal gas law, we have

$$\lambda = \frac{kT}{\sqrt{2} \pi \sigma^2 P}$$

To solve this problem, we must 1) find n , the concentration, 2) substitute the known values into the equation for the mean free path

$$a) P = 1 \text{ atm} \quad \sigma = 3.61 \times 10^{-10} \text{ m}$$

$$T = 25^\circ\text{C} + 273 = 298^\circ\text{K} \quad R = 0.082 \text{ liter-atm/K-mole}$$

$$n = \frac{PN_A}{RT} = \frac{(1 \text{ atm}) \left[6.022 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \right] \left[10^3 \frac{\text{liter}}{\text{m}^3} \right]}{\left[0.082 \frac{\text{liter-atm}}{\text{K-mole}} \right] (298^\circ\text{K})}$$

$$n = 2.46 \times 10^{25} \frac{\text{molecules}}{\text{m}^3}$$

$$\text{Thus, } \lambda = \frac{1}{\sqrt{2} \pi \sigma^2 n}$$

$$= [(1.414) (3.14) (3.61 \times 10^{-10} \text{ m})^2 (2.46 \times 10^{25})]^{-1}$$

$$= 7.02 \times 10^{-8} \text{ m}$$

$$b) P = 10^{-3} \text{ torr} = \frac{10^{-3} \text{ torr}}{760 \frac{\text{torr}}{\text{atm}}} = 1.3157 \times 10^{-6} \text{ atm.}$$

$$T = 298^\circ\text{K} \quad \sigma = 3.61 \times 10^{-10} \text{ m}$$

Here, it is necessary to convert torr units to units of atmospheres.

$$n = \frac{(1.3157 \times 10^{-6} \text{ atm}) \left[6.022 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \right] \left[10^3 \frac{\text{liter}}{\text{m}^3} \right]}{\left[.082 \frac{\text{liter-atm}}{\text{K-mole}} \right] (298^\circ\text{K})}$$

$$= 3.24 \times 10^{15} \text{ molecules/m}^3.$$

$$\text{Thus, } \lambda = \frac{1}{\sqrt{2} \pi \sigma^2 n}$$

$$= [(1.414) (3.14) (3.61 \times 10^{-10} \text{ m})^2 (3.24 \times 10^{15} / \text{m}^3)]^{-1}$$

$$= 0.053 \text{ m} = 5.3 \text{ cm.}$$

• PROBLEM 105

In water vapor at 25°C, what is the average speed of a water molecule in meters/sec? Recall that 1 Joule = 1 Kg - m²/sec². The Boltzmann constant, $k = 1.3806 \times 10^{-23}$ J/deg.

Solution: Recall, that for all bodies in motion, kinetic energy (K.E.) = $\frac{1}{2} M \bar{V}^2$, where M = mass of body and \bar{V} = average velocity of the molecules. Thus, if K.E. and the mass are known, then \bar{V} can be solved. To find K.E., note that for gases average K.E. is proportional to temperature. This can be written as K.E. = $\frac{3}{2} kT$, where k is the Boltzmann constant 1.3806×10^{-23} J/deg and T = temperature in Kelvin (Celsius plus 273°). You are given the temperature of the system (25°C or 298°K). Thus, K.E. = $\frac{3}{2}(1.38 \times 10^{-23})(298) = 6.17 \times 10^{-21}$ J/molecule.

Since K.E. = $\frac{1}{2} M \bar{V}^2$, $6.17 \times 10^{-21} = \frac{1}{2} M \bar{V}^2$. If you knew the mass of one water molecule, you could substitute it into this expression and find V. The mass of the H₂O molecule can be found by using the molecular weight (M.W.) and Avogadro's number (6.02×10^{23} /mole). H₂O has an M.W. of 18 grams/mole. Thus, the mass of one molecule =

$$\frac{18 \text{ g/mole}}{6.02 \times 10^{23} \text{ molecule/mole}} = 2.99 \times 10^{-23} \text{ grams/molecule.}$$

Substituting and solving for V,

$$V = \sqrt{\frac{2(K.E.)}{M}} = \sqrt{\frac{2 \left[6.17 \times 10^{-21} \frac{\text{kg} \cdot \text{m}^2}{\text{sec}^2 \cdot \text{molecule}} \right]}{\left[2.99 \times 10^{-23} \frac{\text{g}}{\text{molecule}} \right] \left[.001 \frac{\text{kg}}{\text{g}} \right]}}$$

$$V = 642 \text{ m/sec.}$$

(Note: .001 kg/g is a conversion factor.)

• PROBLEM 106

The root mean square (rms) speed of hydrogen (H₂) at a fixed temperature, T, is 1600 m/sec. What is the rms speed of oxygen (O₂) at the same temperature?

Solution: To solve this problem we make use of the following equation for the rms speed, v_{rms} ,

$$v_{\text{rms}} = \sqrt{\frac{3 RT}{M}},$$

where R is the gas constant, T the absolute temperature, and M the molecular weight. For oxygen and hydrogen we have

$$v_{\text{rms}(\text{O}_2)} = \sqrt{\frac{3 RT}{M_{\text{O}_2}}} \quad \text{and} \quad v_{\text{rms}(\text{H}_2)} = \sqrt{\frac{3 RT}{M_{\text{H}_2}}}$$

Dividing the first of these equations by the second gives

$$\frac{v_{\text{rms}}(\text{O}_2)}{v_{\text{rms}}(\text{H}_2)} = \frac{\sqrt{\frac{3 RT}{M_{\text{O}_2}}}}{\sqrt{\frac{3 RT}{M_{\text{H}_2}}}} = \sqrt{\frac{3 RT/M_{\text{O}_2}}{3 RT/M_{\text{H}_2}}}$$

or, since R is a constant and T is the same for both gases,

$$\frac{v_{\text{rms}}(\text{O}_2)}{v_{\text{rms}}(\text{H}_2)} = \sqrt{\frac{M_{\text{H}_2}}{M_{\text{O}_2}}}$$

Solving for $v_{\text{rms}}(\text{O}_2)$,

$$\begin{aligned} v_{\text{rms}}(\text{O}_2) &= \sqrt{\frac{M_{\text{H}_2}}{M_{\text{O}_2}}} \times v_{\text{rms}}(\text{H}_2) \\ &= \sqrt{\frac{2.0 \text{ g/mole}}{32.0 \text{ g/mole}}} \times 1600 \text{ m/sec} = 400 \text{ m/sec.} \end{aligned}$$

• PROBLEM 107

Graham's law states that the rate at which gas molecules escape through a small orifice (rate of effusion) is inversely proportional to the square root of the density of the gas. Derive Graham's law from the following assumptions: (a) temperature is directly proportional to the average kinetic energy of the molecules; (b) the rate of effusion is directly proportional to the root mean square speed of the molecules; (c) the density of a gas at constant temperature and pressure is directly proportional to the molecular mass.

Solution: Consider the assumptions individually. The first states that temperature, T, is proportional to the average kinetic energy $\frac{1}{2} m \bar{u}^2$, where m = molecular mass, u = speed, and the bar over u^2 indicates the average. Hence

$$T = k_1 \times \frac{1}{2} m \bar{u}^2$$

where k_1 is a constant.

The second assumption states that the rate of effusion is directly proportional to the root mean square speed of the molecules, $\sqrt{\bar{u}^2}$, or

$$\text{rate} = k_2 \sqrt{\bar{u}^2}$$

where k_2 is a constant.

The third assumption states that the density, p , is directly proportional to the molecular mass, or

$$p = k_3 m$$

where k_3 is a constant. Thus,

$$T = k_1 \times \frac{1}{2} m \bar{u}^2 = k_1 \times \frac{1}{2} \times \frac{p}{k_3} \times \bar{u}^2.$$

Taking the square root of both sides we obtain

$$T^{\frac{1}{2}} = \left(\frac{k_1}{2k_3} \right)^{\frac{1}{2}} p^{\frac{1}{2}} \sqrt{\bar{u}^2}. \text{ Multiplying by } \left(\frac{2k_3}{k_1} \right)^{\frac{1}{2}} \text{ this becomes}$$

$$\left(\frac{2k_3}{k_1} \right)^{\frac{1}{2}} T^{\frac{1}{2}} = p^{\frac{1}{2}} \sqrt{\bar{u}^2}. \text{ At constant temperature, the expression}$$

$$\left(\frac{2k_3}{k_1} \right)^{\frac{1}{2}} T^{\frac{1}{2}} \text{ is a constant, say } k_4. \text{ Then, } k_4 = p^{\frac{1}{2}} \sqrt{\bar{u}^2} = p^{\frac{1}{2}} \times \frac{\text{rate}}{k_2}. \text{ Multiplying by } \frac{k_2}{p^{\frac{1}{2}}} \text{ and letting } k_2 \times k_4 = k_5,$$

$$\text{a constant, we obtain: rate} = \frac{k_2 \times k_4}{p^{\frac{1}{2}}} = \frac{k_5}{p^{\frac{1}{2}}}, \text{ which states}$$

that the rate of effusion is inversely proportional to the square root of the density, which is Graham's law.

CHAPTER 4

AVOGADRO'S HYPOTHESIS; CHEMICAL COMPOUNDS AND FORMULAS

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 95 to 143 for step-by-step solutions to problems.

Avogadro's Law

Avogadro's hypothesis, first introduced by the Italian Renaissance physicist Amedeo Avogadro, states that for two gases at the same temperature and pressure, equal volumes will contain equal numbers of molecules. This number will be the same for all gases.

Avogadro's hypothesis has been so extensively verified that it is now widely known as Avogadro's Law. It follows, from the ideal gas law, that a mole of any gas will contain the same number of molecules. This number, 6.02×10^{23} , is called Avogadro's number and applies to all matter, not just to gases. One mole of any species will contain Avogadro's number of molecules. One mole of any species is the amount of that species whose mass in grams is numerically equal to the molecular or atomic weight.

In order to calculate the number of molecules in a given mass of material, it is necessary to first compute the number of moles. To calculate the number of moles of a substance, divide the mass in grams by the molecular weight as shown in Equation 4-1.

$$n \text{ (moles)} = m \text{ (grams)} / M \text{ (grams/mole)} \quad 4-1$$

The number of molecules is simply the number of moles times Avogadro's number (the molecules per mole for any substance).

$$\text{Number of molecules} = n \times N_A \qquad 4-2$$

Definitions of Atomic Weight, Molecular Weight, and Equivalent Weight

Calculations involving chemical reactions require the understanding of the basic ideas of atomic weight, molecular weight, equivalent weight, and moles. You cannot proceed until you master these concepts and they become so familiar that their use is automatic. The definitions of these terms follow.

Atomic Weight:	The atomic weight of an atom is the mass of the atom compared to Carbon-12 which is defined as 12 amu or 12 atomic mass units. Atomic weights are tabulated for all elements. Unless reference is made to a specific isotope, the tabulated values of atomic weights are average values for the mixture of isotopes that occur naturally.
Molecular Weight:	The molecular weight is the sum of the atomic weights of all the atoms which form the molecule.
Equivalent Weight:	The equivalent weight is the molecular weight divided by the valence—i.e., the number of electrons/molecule which enter the reaction in question. Equivalent weight depends on the reaction and a substance may have different equivalent weights for different reactions.
Mole:	The mass of a substance (in grams) numerically equal to the molecular weight. Sometimes lb moles, kg moles, etc., are defined as the mass (in lb, kg, etc., respectively) numerically equal to the molecular weight. These quantities are not moles, but can be used for convenience in calculations employing those units. Note, however, that Avogadro's number of molecules constitutes only a (gram) mole.

The key to determining a chemical formula from weight percent data of the elemental constituents is the reduction of the amount of each element to moles. Then, from the molar ratio, the empirical or simplest formula will usually become obvious. When amounts of gases are expressed as volumes, the number of moles can be determined from the ideal gas law as discussed in Chapters 2 and 3.

Step-by-Step Solutions to Problems in this Chapter, "Avogadro's Hypothesis; Chemical Compounds and Formulas"

• PROBLEM 108

How many mercury atoms would there be in a 100 g piece of swordfish said to contain 0.1 ppm (part per million by weight) of mercury?

Solution: One can find the number of atoms present by multiplying the number of moles of mercury by the number of atoms in 1 mole. The number of atoms in one mole is equal to Avogadro's Number or 6.02×10^{23} atoms/mole. To solve for the number of mercury atoms present find:

- (1) the amount of mercury present
- (2) the number of moles of mercury
- (3) the number of atoms of mercury.

Solving:

(1) Because there is 0.1 ppm of mercury in the swordfish, one multiplies 0.1×10^{-6} by the amount of fish present.

$$\text{amount of mercury} = 0.1 \times 10^{-6} \times 100 \text{ g} = 1.0 \times 10^{-5} \text{ g.}$$

(2) The number of moles is found by dividing the weight of the mercury present by the weight of one mole, the molecular weight (MW = 200.6)

$$\text{no. of moles} = \frac{1.0 \times 10^{-5} \text{ g}}{200.6 \text{ g/mole}} = 4.99 \times 10^{-8} \text{ moles.}$$

(3) The number of atoms is found by multiplying the number of moles by the number of atoms in one mole, 6.02×10^{23} atoms.

$$\begin{aligned}\text{no. of atoms} &= 4.99 \times 10^{-8} \text{ moles} \times 6.02 \times 10^{23} \text{ atoms/mole} \\ &= 3.00 \times 10^{16} \text{ atoms.}\end{aligned}$$

• PROBLEM 109

If the dot under a question mark has a mass of 1×10^{-6} g, and you assume it is carbon, how many atoms are required to make such a dot?

Solution: Two facts must be known to answer this question. You must determine the number of moles in the carbon dot. You must also remember the number of atoms in a mole of any substance, 6.02×10^{23} atoms/mole (Avogadro's number).

A mole is defined as the weight in grams of a substance divided by the atomic weight (or molecular weight).

The atomic weight of carbon is 12 g/mole. Therefore, in the dot you have

$\frac{1 \times 10^{-6} \text{ g}}{12 \text{ g/mole}}$ moles of carbon. Therefore, the number of atoms in such a dot is the number of moles \times Avogadro's number,

$$\left(\frac{1 \times 10^{-6} \text{ g}}{12 \text{ g/mole}} \right) 6.02 \times 10^{23} \text{ atoms/mole} = 5 \times 10^{16} \text{ atoms.}$$

• PROBLEM 110

What is the approximate number of molecules in a drop of water which weighs 0.09 g?

Solution: The number of molecules in a mole is defined to be 6.02×10^{23} molecules. Thus, to find the number of molecules in a drop of water, one must know the number of moles making up the drop. This is done by dividing the weight of the drop by the molecular weight of H_2O . (MW of H_2O = 18)

$$\text{number of moles} = \frac{0.09 \text{ g}}{18 \text{ g/mole}} = .005 \text{ moles.}$$

The number of molecules present is now found by multiplying the number of moles by Avogadro's number (6.02×10^{23}).

$$\begin{aligned}\text{no. of molecules} &= .005 \text{ moles} \times 6.02 \times 10^{23} \text{ molecules/mole} \\ &= 3.01 \times 10^{21} \text{ molecules.}\end{aligned}$$

• PROBLEM 111

What is the difference between the number of carbon atoms in 1.00 g of C-12 isotope (atomic mass = 12.000 g/mole) and 1.00 g of C-13 isotope (atomic mass = 13.003 g/mole)?

Solution: The difference in the number of carbon atoms in each sample is equal to the difference in the number of moles times Avogadro's number 6.02×10^{23} . Hence, we must begin by calculating the number of moles of C-12 and of C-13 in 1.00 g samples of each.

The number of moles is equal to the mass divided by the atomic weight. Therefore,

$$\text{moles C-12} = \frac{\text{mass C-12}}{\text{atomic mass C-12}} = \frac{1.00 \text{ g}}{12.000 \text{ g/mole}} = 0.083 \text{ mole}$$

$$\begin{aligned} \text{and moles C-13} &= \frac{\text{mass C-13}}{\text{atomic mass C-13}} \\ &= \frac{1.00 \text{ g}}{13.003 \text{ g/mole}} = 0.077 \text{ mole.} \end{aligned}$$

The difference in the number of moles between the two samples is (moles C-12) - (moles C-13) = 0.083 - 0.077 = 0.006 mole. Multiplying by Avogadro's number gives the difference in the number of carbon atoms in the two samples:

$$\begin{aligned} \text{number of carbon atoms} &= 0.006 \text{ mole} \times 6.02 \times 10^{23} \frac{\text{molecules}}{\text{mole}} \\ &= 3.61 \times 10^{21} \text{ molecules.} \end{aligned}$$

• PROBLEM 112

During a moon landing, one of the experiments performed was the measurement of the intensity of solar wind. As a collector, an aluminum strip of about 3000 cm² area was used. It was found that in 100 min, a mass of 3.0×10^{-10} g of H atoms was collected (by the sticking of H atoms to the strip). What was the intensity of the solar wind (in numbers of atoms per cm² per second)?

Solution: The intensity of the solar wind is the number of H atoms striking the unit area in unit time,

$$\text{intensity} = \frac{\text{number of atoms}}{\text{area} \times \text{time}} .$$

We must determine the number of atoms. This is accomplished by converting mass to moles and then multiplying by Avogadro's number.

The number of moles is given by

$$\begin{aligned}\text{moles} &= \frac{\text{mass of H atoms}}{\text{atomic weight of H}} = \frac{3.0 \times 10^{-10} \text{ g}}{1.0 \text{ g/mole}} \\ &= 3.0 \times 10^{-10} \text{ mole.}\end{aligned}$$

Since there are an Avogadro's number of atoms in one mole, the number of atoms in 3.0×10^{-10} mole is

$$\begin{aligned}\text{number of atoms} &= \text{moles} \times \text{Avogadro's number} \\ &= 3.0 \times 10^{-10} \text{ mole} \times 6 \times 10^{23} \text{ atoms/mole} \\ &= 18 \times 10^{13} \text{ atoms.}\end{aligned}$$

The intensity of the solar wind is then

$$\begin{aligned}\text{intensity} &= \frac{\text{number of atoms}}{\text{area} \times \text{time}} = \frac{18 \times 10^{13} \text{ atoms}}{3000 \text{ cm}^2 \times 100 \text{ min}} \\ &= \frac{18 \times 10^{13} \text{ atoms}}{3000 \text{ cm}^2 \times 100 \text{ min} \times 60 \text{ sec/min}} \\ &= 1 \times 10^7 \text{ atoms/cm}^2\text{-sec.}\end{aligned}$$

• PROBLEM 113

The most abundant element in sea water is chlorine (Cl , atomic weight = 35.5 g/mole), which is present as chloride ion, Cl^- , in a concentration of 19 g Cl^- per 1 kg of sea water. If the volume of the earth's oceans is 1.4×10^{21} liters, how many moles of Cl^- are present in the oceans? Assume that the density of sea water is 1.0 g/cm^3 .

Solution: The total number of moles of Cl^- is equal to the total mass of Cl^- divided by its atomic weight. Hence we must find the total mass of Cl^- . This will be done by determining the total mass of the oceans and using this value in conjunction with the concentration of Cl^- to find the total mass of Cl^- .

The volume of the oceans is 1.4×10^{21} liters = 1.4×10^{21} liters $\times 1000 \text{ cm}^3/\text{liter} = 1.4 \times 10^{24} \text{ cm}^3$. Multiplying this volume by the density of sea water gives the total mass of the oceans, or $1.4 \times 10^{24} \text{ cm}^3 \times 1.0 \text{ g/cm}^3 = 1.4 \times 10^{24} \text{ g}$. Expressing this total mass as kilograms, we obtain

$$1.4 \times 10^{24} \text{ g} = 1.4 \times 10^{24} \text{ g} \times 10^{-3} \text{ kg/g} = 1.4 \times 10^{21} \text{ kg.}$$

For every kg of sea water, there are 19 g of Cl^- . Hence, for $1.4 \times 10^{21} \text{ kg}$ there are $1.4 \times 10^{21} \text{ kg} \times 19 \text{ g Cl}^-/\text{kg of sea water} = 2.7 \times 10^{22} \text{ g Cl}^-$.

Therefore, the number of moles of Cl^- is

$$\begin{aligned}\text{moles Cl}^- &= \frac{\text{mass Cl}^-}{\text{atomic weight Cl}^-} \\ &= \frac{2.7 \times 10^{22} \text{ g}}{35.5 \text{ g/mole}} = 7.6 \times 10^{20} \text{ moles.}\end{aligned}$$

• PROBLEM 114

An experiment to measure Avogadro's number involves the collection of a beam of alpha particles to which electrons are added to give neutral helium. Suppose that for a particular run it takes a number of electrons equivalent to 6.40×10^{-5} coulomb/sec to neutralize the beam, and that after 24 hours of collection 1.14×10^{-4} g of neutral helium are collected. From this data, determine Avogadro's number. The charge on one electron is 1.602×10^{-19} coulombs.

Solution: Remembering that it takes two electrons to neutralize one alpha particle, i.e., one positively charged He atom, we will calculate the total charge required during the 24 hour run and, from this, the number of electrons. The number of alpha particles neutralized, or the number of helium atoms produced, is then equal to one-half the number of electrons. Using the mass of helium collected and the atomic mass of helium (4.0026 g/mole), we will calculate the corresponding number of moles. Dividing the number of atoms by the number of moles will give the number of atoms per mole, or Avogadro's number.

24 hours is equivalent to $24 \text{ hours} \times 60 \text{ min/hr} \times 60 \text{ sec/min} = 86,400 \text{ sec}$. Thus, in 24 hours it took

$$6.40 \times 10^{-5} \text{ coulomb/sec} \times 86,400 \text{ sec} = 5.530 \text{ coulomb}$$

to neutralize the beam. Since the charge on one electron is 1.602×10^{-19} coulomb, dividing this number into the total number of coulombs gives the equivalent number of electrons, or

$$\begin{aligned}\text{number of electrons} &= \frac{\text{coulombs}}{\text{coulombs per electron}} \\ &= \frac{5.530 \text{ coulomb}}{1.602 \times 10^{-19} \text{ coulomb/electron}} \\ &= 3.452 \times 10^{19} \text{ electrons.}\end{aligned}$$

It takes two electrons to neutralize one alpha particle, hence the number of alpha particles neutralized by this number of electrons is

$$\begin{aligned}\text{number of alpha particles} &= \frac{\text{number of electrons}}{2} \\ &= \frac{3.452 \times 10^{19}}{2} = 1.726 \times 10^{19} \text{ alpha particles.}\end{aligned}$$

But one alpha particle produces one helium atom, so that the beam produced 1.726×10^{19} helium atoms.

The mass of helium corresponding to this number of helium atoms produced by the beam is given as 1.14×10^{-4} g. Dividing by the atomic mass of helium, we convert this mass to moles:

$$\begin{aligned}\text{moles helium} &= \frac{\text{mass helium}}{\text{atomic mass helium}} = \frac{1.14 \times 10^{-4} \text{ g}}{4.0026 \text{ g/mole}} \\ &= 2.84 \times 10^{-5} \text{ mole.}\end{aligned}$$

Hence, 1.726×10^{19} helium atoms correspond to 2.84×10^{-5} mole of helium. Dividing these two numbers gives Avogadro's number:

$$\begin{aligned}\text{Avogadro's number} &= \frac{\text{number of atoms}}{\text{moles}} \\ &= \frac{1.726 \times 10^{19}}{2.84 \times 10^{-5} \text{ moles}} \\ &= 6.08 \times 10^{23} \text{ atoms/mole.}\end{aligned}$$

Comparing this with the currently accepted value of 6.02×10^{23} atoms/mole, we see that the estimate calculated is in error by

$$\begin{aligned}&\frac{6.08 \times 10^{23} - 6.02 \times 10^{23}}{6.02 \times 10^{23}} \times 100 \% \\ &= \frac{6.08 - 6.02}{6.02} \times 100 \% = \frac{0.06}{6.02} \times 100 \% = 1 \%\end{aligned}$$

• PROBLEM 115

A chemist wants to calculate Avogadro's number by the inspection of a solid cube of AgCl. The density of the cube is 5.56 g/cm^3 . The spacing between the Ag^+ and Cl^- ions in the cube is $2.773 \times 10^{-8} \text{ cm}$ from their centers. From these data, perform this calculation.

Solution: To solve this problem, the following quantities must be determined: the volume of one mole of AgCl, cubic edge length of one mole, the number of ions on an edge, and the total number of ions. The total number of ions

divided by 2 will yield Avogadro's number. From the atomic weights of Ag and Cl (107.868 amu and 35.453 amu) one mole of AgCl weighs 143.321 grams. Because density = mass/volume, the volume of one mole =

$$= \frac{143.321 \text{ g/mole}}{5.56 \text{ g/cm}^3} = 25.78 \text{ cm}^3/\text{mole}.$$

The volume of a cube = $(e)^3$, where "e" is the length of the cube edge. Therefore,

$$e = \sqrt[3]{25.78 \text{ cm}^3/\text{moles}} = 2.954 \text{ cm}/\sqrt[3]{\text{mole}}.$$

Because the spacing between ions is 2.77×10^{-8} cm, the number of ions along the edge is

$$\frac{2.954 \text{ cm}/\sqrt[3]{\text{mole}}}{2.773 \times 10^{-8} \text{ cm}} = 1.065 \times 10^8 \text{ ions}/\sqrt[3]{\text{mole}}$$

The total number of ions in 1 mole of AgCl must be

$$(1.065 \times 10^8 / \sqrt[3]{\text{mole}})^3 = 1.209 \times 10^{24} / \text{mole}.$$

This number is not Avogadro's number. Because you have two ions per formula unit, there is one Avogadro's number of Ag^+ and one Avogadro's number of Cl^- . It follows, therefore, that you must divide by two. As such,

$$\frac{1.209 \times 10^{24} / \text{mole}}{2} = 6.04 \times 10^{23} / \text{mole} = \text{Avogadro's number}.$$

ATOMIC AND MOLECULAR WEIGHTS

• PROBLEM 116

1.3625 g of an unknown metal X reacts with oxygen to form 1.4158 g of the oxide X_2O . What is the atomic mass of X?

Solution: This problem will be solved by calculating the mass and, from this, the number of moles of oxygen in the compound. The corresponding number of moles of metal is twice this amount. The molecular weight is the number of moles of metal divided by the weight of metal that was used.

The mass of oxygen in the compound is

$$\text{mass O} = \text{mass oxide} - \text{mass X}$$

$$= 1.4158 \text{ g} - 1.3625 \text{ g} = 0.0533 \text{ g}.$$

The corresponding number of moles is

$$\text{moles O} = \text{mass O} \div \text{atomic mass O}$$

$$= 0.0533 \text{ g} \div 15.9994 \text{ g/mole} = 0.0033 \text{ mole.}$$

Since, in the oxide, there are two moles of X per mole of O, the number of moles of X present is $2 \times 0.0033 \text{ mole} = 0.0066 \text{ mole X}$. Hence, 1.3625 g of X corresponds to 0.0066 mole X. The atomic mass of X is then:

$$\text{atomic mass X} = \frac{\text{mass X}}{\text{moles X}} = \frac{1.3625 \text{ g}}{0.0066 \text{ mole}} = 206.4394 \text{ g/mole.}$$

• PROBLEM 117

How many moles are contained in 196 g of H_2SO_4 ?

Solution: To calculate the number of moles of a given compound from a certain number of grams of that compound one should use the following relation:

$$\text{number of moles} = \frac{\text{number of grams}}{\text{molecular weight}}$$

When determining the molecular weight of a compound one must first know the atomic weights of the elements within the compound. One then determines the molecular weight by adding together the weights of the individual elements making up the compound. The molecular weights of the elements contained in H_2SO_4 are MW of H = 1, MW of S = 32, and MW of O = 16. If there is more than one atom of a particular element, the molecular weight of the element must be multiplied by the number of atoms present in the compound.

Calculating the molecular weight of H_2SO_4 :

H	2	1.0	= 2
S	1	32	= 32
O	4	16	= 64
Total			98

The molecular weight of H_2SO_4 is thus 98 g. The number of moles of H_2SO_4 in 196 g can now be calculated.

$$\text{number of moles} = \frac{196 \text{ g}}{98 \text{ g/mole}} = 2.0 \text{ moles.}$$

• PROBLEM 118

How many moles are present in 100 g quantities of each of the following? (a) CaCO_3 , (b) H_2O , (c) HCl , (d) $\text{Al}_2(\text{SO}_4)_3$?

Solution: One can find the number of moles of a specific compound in a certain number of grams of that compound by dividing the number of grams present by the molecular weight. The molecular weight is defined as the weight of one mole or Avogadro's Number (6.02×10^{23}) of particles.

$$\text{number of moles} = \frac{\text{weight in grams of sample}}{\text{molecular weight}}$$

One calculates the molecular weight of a compound by adding together the molecular weights of the elements present. When calculating the molecular weight, one must take into account the number of atoms of each element in the compound. This is done by multiplying the molecular weight of the element by the number of atoms present of the particular element. This method will be used in the following examples. Once the molecular weight is determined, the number of moles present in 100 g of the compound can be found by using the equation

$$\text{number of moles in 100 g} = \frac{100 \text{ g}}{\text{molecular weight}}$$

(a) CaCO_3

There is one atom of Ca present in CaCO_3 . Thus, the molecular weight of Ca, 40, is multiplied by one. There is one atom of C. Thus the molecular weight, 12, of C is multiplied by one. Because there are 3 atoms of O present, the molecular weight of O, 16, is multiplied by 3.

1 atom of Ca	$1 \times 40 =$	40
1 atom of C	$1 \times 12 =$	12
3 atoms of O	$3 \times 16 =$	<u>48</u>

$$\text{molecular weight of } \text{CaCO}_3 = 100$$

The number of moles of CaCO_3 in 100 g may now be found.

$$\text{number of moles in 100 g} = \frac{100 \text{ g}}{100 \text{ g/mole}} = 1 \text{ mole.}$$

(b) H_2O

(1) Calculation of molecular weight. The MW of H is 1 and the MW of O is 16.

2 atoms of H	$2 \times 1 =$	2
1 atom of O	$1 \times 16 =$	<u>16</u>
molecular weight of H_2O		= 18

(2) Calculation of number of moles in 100 g of H_2O .

$$\text{number of moles in 100 g} = \frac{100 \text{ g}}{18 \text{ g/mole}} = 5.55 \text{ moles.}$$

(c) HCl

$$\text{MW of H} = 1, \text{ MW of Cl} = 35.5.$$

$$\begin{array}{rcl}
 1 \text{ atom of H} & 1 \times 1 & = 1 \\
 1 \text{ atom of Cl} & 1 \times 35.5 & = 35.5 \\
 \text{molecular weight of HCl} & & = 36.5
 \end{array}$$

$$\text{number of moles of HCl in 100 g} = \frac{100 \text{ g}}{36.5 \text{ g/mole}} = 2.74 \text{ moles}$$

(d) $\text{Al}_2(\text{SO}_4)_3$

$$\text{MW of Al} = 27 \quad \text{MW of S} = 32 \quad \text{MW of O} = 16$$

$$\begin{array}{rcl}
 2 \text{ atoms of Al} & 2 \times 27 & = 54 \\
 3 \text{ atoms of S} & 3 \times 32 & = 96 \\
 12 \text{ atoms of O} & 12 \times 16 & = 192 \\
 \text{molecular weight of } \text{Al}_2(\text{SO}_4)_3 & & = 342
 \end{array}$$

$$\begin{aligned}
 \text{number of moles of } \text{Al}_2(\text{SO}_4)_3 \text{ in 100 g} &= \frac{100 \text{ g}}{342 \text{ g/mole}} \\
 &= .292 \text{ moles.}
 \end{aligned}$$

• PROBLEM 119

What is the molecular weight of a substance, each molecule of which contains 9 carbon atoms and 13 hydrogen atoms and 2.33×10^{-23} g of other components?

Solution: The molecular weight of a compound is the sum of the weights of the components of the compound. It is the weight of one mole of the substance, thus this compound weighs the sum of the weight of 9 moles of C, 13 moles H and $(6.02 \times 10^{23}) \times (2.33 \times 10^{-23} \text{ g})$. Because each molecule of the third substance weighs $2.33 \times 10^{-23} \text{ g}$, one mole of it weighs $(6.02 \times 10^{23}) \times (2.33 \times 10^{-23} \text{ g})$. There are 6.02×10^{23} molecules of this other substance in one mole of the compound (Avogadro's number).

$$\begin{aligned}
 \text{molecular weight} &= (9 \times \text{MW of C}) + (13 \times \text{MW of H}) + \\
 &\quad (2.33 \times 10^{-23} \text{ g}) (6.02 \times 10^{23} / \text{mole}) \\
 &= (9 \times 12.01 \text{ g/mole}) + (13 \times 1.00 \text{ g/mole}) + \\
 &\quad (14.03 \text{ g/mole}) \\
 &= 108.09 \text{ g/mole} + 13.0 \text{ g/mole} + \\
 &\quad 14.03 \text{ g/mole} \\
 &= 135.12 \text{ g/mole.}
 \end{aligned}$$

• PROBLEM 120

Determine the relative abundance of each isotope in naturally occurring gallium from the following data:
At. wt. Ga = 69.72. Masses of isotopes ^{69}Ga = 68.926, ^{71}Ga = 70.925.

Solution: The relative abundance of the various isotopes of gallium can be found by using the following equation:

$$\begin{aligned} \text{atomic weight of Ga} &= \% \text{ of } ^{69}\text{Ga} \times \text{at. wt. of } ^{69}\text{Ga} + \\ &\quad \% \text{ of } ^{71}\text{Ga} \times \text{at. wt. of } ^{71}\text{Ga} \end{aligned}$$

It is given that Ga consists only of the two isotopes ^{69}Ga and ^{71}Ga . Thus, if one lets x = fraction of ^{69}Ga , then $1 - x$ = fraction of ^{71}Ga . Using the above equation one can solve for x .

$$69.72 = (x \times 68.926) + (1 - x) \times 70.925$$

$$69.72 = 68.926x + 70.925 - 70.925x$$

$$- 1.205 = - 1.999x$$

$$x = 1.2/2.0 = 0.60.$$

This means that the ^{69}Ga makes up 60 % of Ga.
 ^{71}Ga makes up $1 - .60$ or 40% of Ga.

• PROBLEM 121

One method for determining the molecular weight of large, biologically important molecules, is by measuring the density by standard procedures and determining the average volume occupied by a single molecule by X-ray crystallographic analysis. If a biochemist measures the density of a sample of deoxyribonucleic acid (DNA) as 1.1 g/cm^3 and X-ray analysis of the same sample estimates the volume of a single DNA molecule as $0.91 \times 10^{-15} \text{ cm}^3$, what is the molecular weight of this type of DNA?

Solution: The number density (molecules/ cm^3) is the reciprocal of the volume of a single molecule and is related to the density (g/cm^3) by the following formula:

$$\begin{aligned} \text{number density} &= \frac{1}{\text{volume per molecule}} \\ &= \frac{\text{density}}{\text{molecular weight}} \times \text{Avogadro's number} \end{aligned}$$

The validity of this formula is readily seen by considering the dimensions of the quantities involved.

Dividing the density (g/cm^3) by the molecular weight (g/mole) gives a quantity with units of

$$\frac{\text{g/cm}^3}{\text{g/mole}} = \text{mole/cm}^3 \text{ (molar density).}$$

Multiplying this number by Avogadro's number (molecules/mole) gives a quantity with dimensions of

$$\frac{\text{mole}}{\text{cm}^3} \times \frac{\text{molecules}}{\text{mole}} = \text{molecules/cm}^3 \text{ (number density).}$$

Solving for the molecular weight in the above expression we obtain

$$\begin{aligned} \text{molecular weight} &= \text{density} \times \text{Avogadro's number} \times \text{volume} \\ &\quad \text{per molecule} \\ &= 1.1 \text{ g/cm}^3 \times 6 \times 10^{23} \text{ molecules/mole} \times \\ &\quad \times 0.91 \times 10^{-15} \text{ cm}^3/\text{molecule} \\ &= 6 \times 10^8 \text{ g/mole.} \end{aligned}$$

• PROBLEM 122

When metal M is heated in halogen X_2 , a compound MX_n is formed. In a given experiment, 1.00 g of titanium reacts with chlorine to give 3.22 g of compound. What is the corresponding value of n ?

Solution: In the compound MX_n , there is one mole of M and n moles of X. In the compound here, one is given that 1.00 g of it is Ti (titanium) and that the compound weighs 3.22 g. This means that 3.22 g - 1.00 g or 2.22 g of the compound is Cl. To find the simplest formula for the compound, one must first determine the number of moles of each component present. This is done by dividing the weight present by the molecular weight

$$\text{no. of moles} = \frac{\text{no. of grams}}{\text{MW}}$$

$$\text{For Ti: (MW} = 47.9)$$

$$\text{no. of moles} = \frac{1.00 \text{ g}}{47.9 \text{ g/mole}} = 2.09 \times 10^{-2} \text{ moles.}$$

$$\text{For Cl: (MW} = 35.5)$$

$$\text{no. of moles} = \frac{2.22 \text{ g}}{35.5 \text{ g/mole}} = 6.25 \times 10^{-2} \text{ moles.}$$

Since the simplest formula for the compound is TiCl_n , n can be found using the following ratio.

$$\frac{1 \text{ Ti}}{n} = \frac{2.09 \times 10^{-2} \text{ moles Ti}}{6.25 \times 10^{-2} \text{ moles Cl}}$$

$$n = \frac{1 \text{ Ti} \times 6.25 \times 10^{-2} \text{ Cl}}{2.09 \times 10^{-2} \text{ Ti}} = 2.99 \text{ Cl}$$

The formula for the compound is therefore TiCl_3 . 2.99 Cl is rounded off to the nearest whole number, 3.

• PROBLEM 123

If the density of ethylene is 1.25 g/liter at S.T.P. and the ratio of carbon to hydrogen atoms is 1 : 2, what is molecular weight and formula of ethylene?

Solution: This problem is solved once you know that at S.T.P. (Standard Temperature and Pressure, 0°C , and 1 atm) one mole of any gas occupies 22.4 liters. Assume that one mole of ethylene gas is present. Density = mass/volume. As such the mass of the gas = $(1.25 \text{ g/l})(22.4 \text{ l}) = 28.0 \text{ g}$. Therefore one mole of ethylene weighs 28 g.

From the ratio given in the question you know the molecular formula can be represented as $(\text{CH}_2)_x$. To obtain the actual molecular formula, look for a compound that has a molecular weight of 28 g yet maintains the carbon : hydrogen ratio of 1 : 2. By looking at the periodic table for atomic weights and through some arithmetic, you will find that the only formula that meets these requirements is $(\text{CH}_2)_2$ or C_2H_4 . This formula can also be found by dividing the weight of 1 CH_2 into 28 g. This solves for x in the expression $(\text{CH}_2)_x$. MW of $\text{CH}_2 = 14$.

$$\text{no. of CH}_2 = \frac{28 \text{ g}}{14 \text{ g/CH}_2} = 2 \text{ CH}_2 = \text{C}_2\text{H}_4.$$

• PROBLEM 124

Phosphorus (atomic weight = 30.97) combines with another element such that 1 g of phosphorus requires 0.7764 g of the other element. If the atomic ratio of phosphorus to the other element is 4 : 3, what is the atomic weight of the unknown element?

Solution: First determine how many moles of phosphorus reacted. From this, one can calculate the number of moles

of the unknown element that reacted using the atomic ratio. This ratio represents the relative mole amounts. If the number of moles of the unknown element is determined, the atomic weight can be calculated because the number of moles is equal to number of grams/atomic weight. (At. wt. of P = 30.97).

One can now determine the atomic weight of the unknown element. The number of moles of phosphorus that reacted

$$= \frac{1.0 \text{ g}}{30.97 \text{ g/mole}} = 0.03229 \text{ moles.}$$

Let x = the number of moles of the unknown element. The atomic ratio of phosphorus to the unknown element is 4 : 3. Thus,

$$\frac{4 \text{ atoms}}{3 \text{ atoms}} = \frac{.03229 \text{ moles}}{x} \quad \text{Solving,}$$

$$x = 0.02422 \text{ moles unknown element.}$$

Therefore, the atomic weight of the unknown element is equal to

$$\frac{0.7764 \text{ g}}{0.02422 \text{ moles}} = 32.06 \text{ g/mole.}$$

• PROBLEM 125

1.0g of scandium (Sc) combines with oxygen (O) to form 1.5338 g of oxide. Assuming the oxide contains two atoms of scandium for every three atoms of oxygen, calculate the atomic weight of Sc. Oxygen has an atomic weight of 15.9994 amu.

Solution: This problem can be solved once a proportion is set up.

The oxide weighs 1.5338 g and the Sc, 1.0 g. Therefore, the weight of the oxygen that reacted must be $1.5338 - 1 = .5338 \text{ g}$.

The problem stated that 3 oxygen atoms reacted for every 2 Sc atoms. Therefore, the total atomic weight of the oxygen in the compound is 3×15.9994 .

Let y be the atomic weight of Sc, then $2y$ is the total weight involved in the formation of the oxide. Therefore, there is .5338 g of oxygen whose atoms (3 of them) have an atomic weight of 3×15.9994 , reacting with 1 g of Sc whose atoms weigh $2y$. This can be represented as

$$\frac{.5338 \text{ g}}{3(15.9994)} = \frac{1.00 \text{ g}}{2y}$$

Solving for y, results in 44.96, which is the atomic weight of scandium.

• PROBLEM 126

A given sample of pure compound contains 9.81 g of zinc, 1.8×10^{23} atoms of chromium and 0.60 mole of oxygen atoms. What is the simplest formula?

Solution: The simplest formula for this compound is found from the simplest ratio of moles Zn : moles Cr : moles O. One is given the number of moles of O but must find the number of moles of Zn and Cr. One is given that there is 9.81 g of Zn. One can find the number of moles by dividing 9.81 g by the MW of Zn. (MW of Zn = 65.4).

$$\text{no. of moles} = \frac{9.81 \text{ g}}{65.4 \text{ g/mole}} = 0.15 \text{ moles}$$

One is given that there are 1.8×10^{23} atoms of Cr present. The number of moles can be found by dividing the number of atoms by the number of atoms in one mole, 6.02×10^{23} .

$$\text{no. of moles} = \frac{1.8 \times 10^{23} \text{ atoms}}{6.02 \times 10^{23} \text{ atoms/mole}} = 0.30 \text{ moles}$$

The ratio of Zn : Cr : O is .15 : .30 : .60 or 1 : 2 : 4. The simplest formula is ZnCr_2O_4 .

• PROBLEM 127

Two different compounds of elements A and B were found to have the following composition: first compound, 1.188 g of A combined with 0.711 g of B; second compound, 0.396 g of A combined with 0.474 g of B. (a) Show that these data are in accord with the law of multiple proportions. (b) If the formula for the first compound is AB_2 , what is the formula for the second?

Solution: (a) The law of multiple proportions can be stated: When two elements combine to form more than one compound, the different weights of one that combine with a fixed weight of the other are in the ratio of small whole numbers. This means that if one solves for the expected amount of B that is used in forming the second compound from the ratio of A : B in experiment one, the experimental amount should be a multiple of the calculated value. This is seen more clearly after looking at the data.

In experiment 1, A combines with B in a ratio of 1.188 g A : 0.711 g B or 1 : .598. In experiment 2, A

combined with B in a ratio of 0.396 A : 0.479 B or 1 : 1.20. The law of multiple proportions states that .598 should be a small multiple of 1.20. $1.20/.598 = 2$, thus the law is supported.

(b) If the formula for the first compound is AB_2 , one knows that the proportion of the number of moles of A to B is 1 : 2. Thus 1 unit volume of A weighs 1.188 g and 2 units of B weigh 0.711 g. Therefore, 1 unit of B weighs $0.711 \text{ g}/2 = 0.356 \text{ g}$. Using this data, one solves for the number of units of A in the second compound by dividing the weight of A in compound 2 by 1.188 g.

$$\text{no. of units of A in compound 2} = \frac{.396 \text{ g}}{1.188 \text{ g/unit}} = .331 \text{ units}$$

$$\text{no. of units of B in compound 2} = \frac{.474 \text{ g}}{.356 \text{ g/unit}} = 1.331 \text{ units}$$

The ratio of A to B in compound 2 can now be found.
Let x = no. of B atoms

$$\frac{A}{B} = \frac{.331}{1.331} = \frac{1}{x}$$

$$x = \frac{1.331}{.331} = 4.0$$

Therefore, the second compound is AB_4 . This result could also have been obtained by using the data from part (a). It was determined that twice as much B is present in the second compound. Thus, if the first compound is AB_2 , then the second compound must be AB_4 .

• PROBLEM 128

When 10.00 g of phosphorus was reacted with oxygen, it produced 17.77 g of a phosphorus oxide. This phosphorus oxide was found to have a molecular weight of approximately 220 in the vapor state. Determine its molecular formula.

Solution: The molecular formula of a substance indicates the relative number of atoms in a molecule of the substance. Therefore, to solve this problem, you must first calculate the ratios of the gram-atoms to each other, the empirical formula, and then extrapolate to the molecular formula via the molecular weight.

The number of gram-atoms of phosphorus (P) is

$$\frac{\text{wt. in grams of P}}{\text{atom weight}} = \frac{10}{30.97} = 0.323 \text{ gram-atoms P.}$$

$$\text{For oxygen we have } \frac{7.77 \text{ g}}{16.00} = .484 \text{ gram-atoms}$$

The weight in grams of oxygen is 7.77 because the final product weighs 17.77 g and the phosphorus weighs 10.00 g. Since, the only other element is oxygen, its weight must be the difference.

The ratio of the gram-atoms of P and O is respectively 1 : 1.5 or 2 : 3. Therefore, the empirical formula of the oxide is P_2O_3 .

To calculate the molecular formula, we must use the stated molecular weight of 220. We must look for a formula that totals to this molecular weight AND maintains the 2 : 3 ratio of P : O as expressed in the empirical formula. With some arithmetic, we find that the only formula that meets these two requirements is P_4O_6 . 4 : 6 is the same as 2 : 3. The atomic weight of P and O is respectively 30.97 and 16. We have four P atoms for a total of 123.88 and we have 6 O atoms for a total of 96. Now add: $123.88 + 96 = 219.88$, which is approximately 220.

Another method for determining the molecular formula is to divide the molecular weight of the molecule, 220, by the weight of 1 P_2O_3 , 110.

$$\text{no. of } P_2O_3 = \frac{220 \text{ g}}{110 \text{ g/mole of } P_2O_3} = 2 \text{ moles of } P_2O_3.$$

The formula is, therefore $2 \times P_2O_3$ or P_4O_6 .

• PROBLEM 129

A chemist reacts metal "B" with sulfur and obtains a compound of metal and sulfur. Assuming metal "B" weighed 2.435 g (MW = 121.75 amu), and the compound weighs 3.397 g, what is the simplest or empirical formula of the compound? The atomic weight of sulfur is 32.06 g/mole.

Solution: To calculate the simplest formula, one must know the mole ratio of the elements that reacted.

A mole is defined as the weight in grams of a substance divided by its atomic weight. The number of moles of "B"

$$= \frac{2.435 \text{ g}}{121.75 \text{ g/mole}} = .0200 \text{ moles of "B".}$$

The number of grams of sulfur that reacted must be $3.397 - 2.435 = .962 \text{ g}$.

The increase in weight can only be derived from the addition of sulfur.

The number of moles of sulfur is

$$\frac{0.962 \text{ g}}{32.06 \text{ g/mole}} = .0300 \text{ moles of sulfur.}$$

$$\text{Experiment 1} \quad \frac{6.08 \text{ g A}}{4.00 \text{ g B}} = 1.5$$

$$\text{Experiment 2} \quad \frac{18.24 \text{ g A}}{12.00 \text{ g B}} = 1.5$$

$$\text{Experiment 3} \quad \frac{3.04 \text{ g A}}{2.00 \text{ g B}} = 1.5$$

The Law of Definite Proportions is illustrated here.

(b) From the Law of Definite Proportions, one can find the number of grams of B that will combine with 71 g of C. After this weight is found, one can find the number of grams of A that will react with 71 g of C by finding the amount of A that reacts with that amount of B. It is assumed that the amount of A that reacts with 71 g of C is equal to the amount of A that will react with the amount of B that reacts with 71 g of C. The amount of A that will react with this amount of B can be found by remembering, from the previous section of this problem, that A reacts with B in a ratio of 1.5.

(1) Finding the amount B that would react with 71 g of C.

One is told that 80 g of B reacts with 355 g of C. By the Law of Definite Proportions, a ratio can be set up to calculate the number of grams of B that will react with 71 g of C.

Let x = the number of grams of B that will react with 71 g of C.

$$\frac{80 \text{ g B}}{355 \text{ g C}} = \frac{x \text{ g B}}{71 \text{ g C}} \quad \text{of C.}$$

$$x = \frac{71 \times 80}{355} = 16.$$

16 grams of B will react with 71 g of C.

(2) It is assumed that the same amount of A that will react with 16 g of B will react with 71 g of C. Therefore, using the fact that the ratio of the amount of A that reacts to the amount of B is equal to 1.5 (this fact was obtained in part (1)), one can calculate the amount of A that will react with 71 g of C.

Let x = the number of grams of A that will react with 16 g of B.

$$\frac{x \text{ g A}}{16 \text{ g B}} = 1.5 \quad \text{of B.}$$

$$x = 16 \times 1.5 = 24 \text{ g.}$$

24 g of A will react with 16 g of B or 71 g of C.

(c) In finding the equivalent weight of C when B is taken to be oxygen, the Law of Definite Proportions is used again. The equivalent weight of oxygen is 8. Knowing that 16 g of B react with 71 g of C, one can set up the following ratio

x = weight of C if the weight of B is taken to be 8.

$$\frac{71 \text{ g C}}{16 \text{ g B}} = \frac{x \text{ g C}}{8 \text{ g B}}$$

$$x = \frac{8 \times 71}{16} = 35.5 \text{ g.}$$

The equivalent weight of C when B is taken to be oxygen is 35.5 g.

EQUIVALENT WEIGHTS

• PROBLEM 131

If the atomic weight of oxygen was 50, what would its equivalent weight be?

Solution: The relationship of the atomic weight and the equivalent weight is

$$\frac{\text{Atomic weight}}{\text{Equivalent weight}} = \text{valence number}$$

The valence number is a measure of the number of atoms of hydrogen that will combine with one atom of the element. Two hydrogen atoms combine with one oxygen. The valence number of oxygen is therefore 2. One can now solve for the equivalent weight of oxygen when the atomic weight is taken as 50.

$$\frac{50}{\text{equivalent weight}} = 2$$

$$\text{equivalent weight} = \frac{50}{2} = 25.$$

• PROBLEM 132

A compound of vanadium and oxygen is analyzed and found to contain 56.0 % vanadium. What is the equivalent weight of vanadium in this compound?

Solution: To find the equivalent weight of vanadium one can use the Law of Definite Proportions. This law states that when elements combine to form a given compound, they do so in a fixed and invariable ratio by weight. This means that the ratio of the weight of vanadium to the weight

of the oxygen that reacts is equal to the ratio of the equivalent weight of the vanadium to the equivalent weight of the oxygen.

$$\frac{\text{weight of V}}{\text{weight of O}} = \frac{\text{equivalent weight of V}}{\text{equivalent weight of O}}$$

Solving for the equivalent weight of vanadium in this compound, one assumes he has 100 g of the compound for calculations. Because 56 % of this compound is vanadium, it means that 56 g of it is vanadium. This indicates that $(100 \% - 56 \%) = 44 \%$ of the compound is oxygen and that in 100 g of the compound there are 44 g of oxygen.

Here one will assume that the equivalent weight of oxygen is its atomic number, 8. Solving for the equivalent weight of vanadium:

$$\frac{56 \text{ g}}{44 \text{ g}} = \frac{\text{equivalent weight of V}}{8}$$

$$\text{equivalent weight of V} = \frac{8 \times 56 \text{ g}}{44 \text{ g}} = 10.2$$

The equivalent weight of vanadium in this compound is 10.2.

• PROBLEM 133

2.0 g of molybdenum (Mo) combines with oxygen to form 3.0 g of a molybdenum oxide. Calculate the equivalent weight of Mo in this compound.

Solution: The equivalent weight of a compound can be determined by use of the Law of Definite Proportions. This law states that when elements combine to form a given compound, they do so in a fixed and invariable ratio by weight. This means that the following ratio is maintained:

$$\frac{\text{weight of Mo in reaction}}{\text{weight of O in reaction}} = \frac{\text{equivalent weight of Mo}}{\text{equivalent weight of O}}$$

The equivalent weight of oxygen is equal to 8. The weight of oxygen in the reaction can be determined by using the Law of Conservation of Mass. This law states that there is no detectable gain or loss of mass in a chemical change. The problem states that 2.0 g of Mo is added to an unknown amount of O to form 3.0 g of a molybdenum oxide. Using the Law of Conservation of Mass the following relation is true:

$$\text{weight of compound} = \text{weight of O} + \text{weight of Mo}$$

$$3.0 \text{ g} = \text{weight of O} + 2.0 \text{ g}$$

$$1.0 \text{ g} = \text{weight of O}$$

One can now use the Law of Definite Proportions to solve for the equivalent weight of Mo.

$$\frac{\text{weight of Mo}}{\text{weight of O}} = \frac{\text{equivalent weight of Mo}}{\text{equivalent weight of O}}$$

$$\frac{2.0 \text{ g}}{1.0 \text{ g}} = \frac{\text{equivalent weight of Mo}}{8}$$

$$\text{equivalent weight of Mo} = 8 \times \frac{2.0 \text{ g}}{1.0 \text{ g}} = 16.$$

• PROBLEM 134

It was found that a magnesium oxide contained .833 g of oxygen and 1.266 g of magnesium. Calculate the gram-equivalent weight of magnesium.

Solution: You begin this problem by establishing what is meant by the term gram-equivalent weight.

Gram-equivalent weight may be defined as the number of grams of an element that will involve a gain or loss of N electrons, i.e., the Avogadro's number (6.02×10^{23}) of electrons, when the element enters into chemical combination with another element. In this problem, you know that oxygen is present. Oxygen has an oxidation state of -2 . In a reaction, one mole of oxygen will gain two moles of electrons. The molecular weight of oxygen is 16 g/mole. Its equivalent weight becomes $16/2 = 8.00$ g, since 2 moles of electrons will be gained and, by definition, equivalent weight is the amount of a substance that will gain or lose one mole of electrons.

The gram-equivalent weight of magnesium is that amount of the element that combines with 8.00 g of oxygen. You are told that 1.266 g Mg combines with .833 g O_2 . If you let x = grams of Mg that will combine with 8.00 of oxygen, you can set up the following proportion:

$$\frac{1.266 \text{ g Mg}}{.833 \text{ g } O_2} = \frac{x \text{ g Mg}}{8.00 \text{ g } O_2}$$

Solving for x , you obtain $x = 12.16$ g of Mg, which is its gram-equivalent weight.

• PROBLEM 135

A chemist forms magnesium oxide by burning magnesium in oxygen. The oxide obtained weighed 1.2096 grams. It was formed from .7296 g of magnesium. Determine the mass equivalent of magnesium in this reaction.

Solution: An equivalent is defined as that mass of oxidizing or reducing agent that picks up or releases the Avogadro number of electrons in a particular reaction. One equivalent of any reducing agent reacts with one equivalent of any oxidizing agent. In this problem, the key is to determine the number of equivalents of oxygen involved. Once this is known, you also know the number of equivalents of magnesium.

Since the oxide weighed 1.2096 g and the magnesium weighed .7296 g, the mass of the combined oxygen must be $1.2096 - .7296 = .4800$ g. Before the oxygen reacted, its oxidation state was zero. After the reaction, however, it was - 2. As such, each oxygen atom gained 2 electrons. Therefore, the Avogadro number of electrons will be taken up by one half of a mole of O atoms.

It follows, therefore, that there are 8.000 g per equivalent for oxygen, since 1 mole of oxygen atoms weighs 16 grams. It was found, however, that there were .4800 g of oxygen. As such

$$\frac{.4800 \text{ g}}{8.00 \text{ g/equiv}} = .0600 \text{ equiv of oxygen.}$$


This means that magnesium also has .06 equiv. 0.7296 g of Mg participated in the reaction. Therefore, the grams per equivalent of Mg =

$$\frac{.7296 \text{ g Mg}}{.060 \text{ equiv}} = 12.16 \text{ g/equiv.}$$

• PROBLEM 136


For the oxidation of VO by Fe_2O_3 to form V_2O_5 and FeO , what is the weight of one equivalent of VO and of Fe_2O_3 ?

Solution: The equation for this reaction is



Here V is a reducing agent and Fe is an oxidizing agent. One equivalent of an oxidizing agent is defined as the mass of the substance that picks up the Avogadro number of electrons. One equivalent of a reducing agent is defined as that mass of the substance that releases the Avogadro number of electrons. The oxidation state of O is always - 2, thus the oxidation state of V in VO is + 2.

The oxidation state of V in V_2O_5 is + 5 because 5 O contribute $(5 \times - 2)$ or - 10, thus 2V must be + 10 and V is + 5. The half-reaction for V is



This means that there are 3 equivalents per mole of

VO. One finds the weight of one equivalent by dividing the molecular weight by 3. (MW of VO = 66.94).


$$\text{weight of 1 equiv of V} = \frac{66.94 \text{ g/mole}}{3 \text{ equiv/mole}} = 22.31 \text{ g}$$

One uses a similar method for Fe. The oxidation state of O in Fe_2O_3 is (3×-2) or -6 , this means the 2Fe must be $+6$ and Fe must be $+6/2$ or $+3$. The oxidation state of O in FeO is -2 , thus the oxidation state of Fe is $+2$.

The half-reaction for the Fe is then



In Fe_2O_3 , there are 2 moles of Fe, therefore the half-reaction becomes



There are thus 2 equiv per mole of Fe_2O_3 . The weight of one equivalent is equal to the weight of one mole Fe_2O_3 divided by 2. (MW of Fe_2O_3 = 159.70)

$$\text{wt of 1 equiv} = \frac{159.7 \text{ g/mole}}{2 \text{ equiv/mole}} = 79.85 \text{ g/equiv.}$$

• PROBLEM 137

In acting as a reducing agent a piece of metal M, weighing 16.00 g, gives up 2.25×10^{23} electrons. What is the weight of one equivalent of the metal?

Solution: One equivalent of a reducing agent is defined as that mass of the substance that releases the Avogadro number of electrons. Avogadro's number is 6.02×10^{23} , thus one can find the number of equivalents in 16.00 g of the metal by dividing 2.25×10^{23} by 6.02×10^{23} .

$$\text{no. of equiv} = \frac{2.25 \times 10^{23} \text{ electrons}}{6.02 \times 10^{23} \text{ electrons/equiv}} = .374 \text{ equiv.}$$

Thus, .374 equiv weigh 16.00 g, one can find the weight of one equivalent by dividing 16.0 g by 0.374 equiv.

$$\text{weight of 1 equiv} = \frac{16.0 \text{ g}}{0.374 \text{ equiv}} = 42.78 \text{ g/equiv.}$$

CHEMICAL COMPOSITION-WEIGHT AND VOLUME PERCENT

• PROBLEM 138

What is the simplest formula of a compound that is composed of 72.4 % iron and 27.6 % oxygen by weight?

Solution: For purposes of calculation, let us assume that there is 100 g of this compound present. This means that there are 72.4 g of Fe and 27.6 g O. The simplest formula for this compound is Fe_nO_m , where n is the number of moles of Fe present and m is the number of moles of O. One finds the number of moles by dividing the number of grams by the molecular weight.

$$\text{number of moles} = \frac{\text{number of grams}}{\text{MW}}$$

For Fe: n = number of moles present. $\text{MW} = 55.8$.

$$n = \frac{72.4 \text{ g}}{55.8 \text{ g/mole}} = 1.30 \text{ moles}$$

For O: m = number of moles present. $\text{MW} = 16.0$.

$$m = \frac{27.6 \text{ g}}{16.0 \text{ g/mole}} = 1.73 \text{ moles.}$$

One solves for the simplest formula by finding the ratio of Fe : O.

$$\frac{\text{Fe}}{\text{O}} = \frac{1.30}{1.73} = .75 = \frac{3}{4}$$

Therefore, $n = 3$ and $m = 4$.

The simplest formula is Fe_3O_4 .

• PROBLEM 139

An unknown compound consists of 82.98 % potassium and 17.02 % oxygen. What is the empirical formula of the compound?

Solution: The empirical formula of any compound is the

ratio of the atoms that make up the compound by weight. It is the simplest formula of a material that can be derived solely from its components. Therefore, we must determine the ratio of gram-atoms of potassium (K) to the number of gram-atoms of oxygen (O).

The number of gram-atoms of a substance equals the weight of the substance in grams divided by the weight per gram-atom of the substance. In other words,

$$\text{number of gram-atoms} = \frac{\text{weight in grams}}{\text{weight per gram-atom}}$$

In this problem, we are given the percentages of the elements that make up the compound. These percentages are, in reality, the weight in grams, since in the definition of weight, we imply percentage. The weight per gram atom is the atomic weight of the element which can be found in the periodic table of elements.

Therefore, the number of gram-atoms for potassium is

$$\frac{82.98 \text{ g (wt of K)}}{39.10 \text{ (atomic weight)}} = 2.120 \text{ moles.}$$

For oxygen, the number of gram-atoms is

$$\frac{17.02 \text{ g (wt of O)}}{16.00 \text{ (atomic wt)}} = 1.062 \text{ moles.}$$

Recall, the empirical formula is the ratio of the elements by weight. Consequently, the ratio of potassium to oxygen is 2 : 1, since the gram - atom ratios are respectively 2.120 : 1.062. Therefore, the empirical formula is K_2O .

• PROBLEM 140

A certain hydrate analyzes as follows: 29.7 % copper, 15.0 % sulfur, 2.8 % hydrogen, and 52.5 % oxygen. Determine the empirical formula of this hydrate from these percentages.

Solution: A hydrate is the chemical combination of water with another compound. For example, copper sulfate combines with water to form the hydrate of the composition $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$



One can find the empirical formula for a compound, when given the weight percents of the various elements making up the compound. This is done by finding the number of moles of each element in 100 g of the compound. The weight of each element is equal to the percent weight.

The number of moles is equal to the weight divided by the molecular weight of the element. In 100 g of this compound, there are 29.7 g Cu, 15.0 g S, 2.8 g H, and 52.5 g O. One can now determine the number of moles of each element present.

$$\text{number of moles} = \frac{\text{number of grams}}{\text{MW}}$$

MW of Cu = 63.5, MW of S = 32, MW of H = 1, MW of O = 16.

$$\text{number of moles of Cu} = \frac{29.7 \text{ g}}{63.5 \text{ g/mole}} = 0.47 \text{ moles}$$

$$\text{number of moles of S} = \frac{15.0 \text{ g}}{32.0 \text{ g/mole}} = 0.47 \text{ moles}$$

$$\text{number of moles of H} = \frac{2.8 \text{ g}}{1 \text{ g/mole}} = 2.8 \text{ moles}$$

$$\text{number of moles of O} = \frac{52.5 \text{ g}}{16 \text{ g/mole}} = 3.28 \text{ moles.}$$

To determine the empirical formula for this hydrate one must look at the ratio of Cu : S : H : O.

The ratio of the number of moles of these elements is

Cu : S : H : O

.47 : .47 : 2.8 : 3.28

To find the empirical formula, these numbers should be made into integers. This is done by making the lowest number equal to 1 and solving for the other three.

$$\frac{.47}{1} = \frac{.47}{x} \quad x = \frac{.47}{.47} = 1$$

$$\frac{.47}{1} = \frac{2.8}{x} \quad x = \frac{2.8}{.47} = 6$$

$$\frac{.47}{1} = \frac{3.28}{x} \quad x = \frac{3.28}{.47} = 7$$

The ratio now becomes

Cu : S : H : O

1 : 1 : 6 : 7.

The empirical formula for the compound is CuSH_6O_7 or $\text{CuSO}_4 \cdot 3\text{H}_2\text{O}$.

• PROBLEM 141

A chemist finds that an unknown compound contains

50.05 % S and 49.95 % O by weight. Calculate its simplest formula.

Solution: To calculate a compound's simplest formula, you need the relative number of moles of atoms in the compound. The percentages by weight of the elements allow for this calculation.

If you had 100 g of the unknown compound, it would consist of 50.05 g of S and 49.95 g of O. A mole is defined as weight in grams/atomic weight.

Therefore, the number of moles of sulfur

$$(S) = \frac{50.05 \text{ g}}{32.06 \text{ g/mole}} = 1.561 \text{ moles.}$$

The number of moles of oxygen =

$$\frac{49.95 \text{ g}}{15.999 \text{ g/mole}} = 3.122 \text{ moles.}$$

You see that the ratio of S to O is

$$\frac{1.561}{3.122} = \frac{1}{2} ; 1 : 2.$$

Therefore, the simplest formula can be expressed as SO_2 .

• PROBLEM 142

A sample of the poisonous compound nicotine extracted from cigarette smoke was found to contain 74.0 % by weight of carbon (C, atomic weight = 12.0 g/mole), 8.65 % by weight of hydrogen (H, atomic weight = 1.01 g/mole), and 17.3 % by weight of nitrogen (N, atomic weight = 14.0 g/mole). What is the empirical formula of nicotine?

Solution: The empirical formula states the relative ratio of the atoms of the various elements in any substance; the actual number of atoms (molecular formula) is not implied. Problems of this sort are solved by assuming a sample of some convenient mass and then calculating the number of moles of each atom in this sample. Once the number of moles has been calculated for each component, these numbers are divided by the greatest common factor in order to obtain the proportions in which the various components appear in the molecule. There is no error in assuming a sample of definite mass, since the size of a sample of molecules does not affect the composition of the molecules.

Assume a sample weighing 100 g. Then the masses of C, H, and N in this sample are:

$$\text{mass C} = 74.0 \% \times 100 \text{ g} = 74.0 \text{ g}$$

$$\text{mass H} = 8.65 \% \times 100 \text{ g} = 8.65 \text{ g}$$

$$\text{mass N} = 17.3 \% \times 100 \text{ g} = 17.3 \text{ g}$$

We convert the mass of each element to the corresponding number of moles by dividing by the atomic mass of that element. Thus we obtain the following number of moles of each element:

$$\text{moles C} = 74.0 \text{ g} / 12.0 \text{ g/mole} = 6.2 \text{ moles}$$

$$\text{moles H} = 8.65 \text{ g} / 1.01 \text{ g/mole} = 8.6 \text{ moles}$$

$$\text{moles N} = 17.3 \text{ g} / 14.0 \text{ g/mole} = 1.2 \text{ moles}$$

The greatest common factor is 1.2 moles. Dividing the number of moles of each element by 1.2 moles, we obtain

$$\frac{\text{moles C}}{1.2 \text{ moles}} = \frac{6.2 \text{ moles}}{1.2 \text{ moles}} \approx 5.$$

$$\frac{\text{moles H}}{1.2 \text{ moles}} = \frac{8.6 \text{ moles}}{1.2 \text{ moles}} \approx 7.$$

$$\frac{\text{moles N}}{1.2 \text{ moles}} = \frac{1.2 \text{ moles}}{1.2 \text{ moles}} = 1.$$

Hence, C, H, and N appear in the proportions 5, 7, and 1, respectively. The empirical formula of nicotine is thus $\text{C}_5\text{H}_7\text{N}$.

• PROBLEM 143

A compound subjected to analysis was found to have the following composition by weight: 69.96 % carbon (atomic weight = 12.0 g/mole), 7.83 % hydrogen (atomic weight = 1.01 g/mole), and 22.21 % oxygen (atomic weight = 16.0 g/mole). If the molecular weight of this compound is 360 g/mole, what is its molecular formula?

Solution: Molecular formula may be defined as the formula stating the actual number of each type atom in a particular compound. Problems of this sort are solved by assuming a sample of some convenient mass and then calculating the number of moles of each atom in this sample. Once the number of moles has been calculated for each component, these numbers are divided by the greatest common factor in order to obtain the proportions in which the various components appear in the molecule (empirical formula). From this, we check to see if the elements, in those proportions, give the actual molecular weight. If they do, we have the actual molecular formula. If they do not, we multiply by a factor until their weights do give the molecular weight. There is no error in assuming a sample of definite mass, since the size of a sample

of molecules does not affect the composition of the molecules.

Assume a sample weighing 100 g. Then the masses of C, H, and O in the sample are:

$$\text{mass C} = 69.96 \% \times 100 \text{ g} = 69.96 \text{ g}$$

$$\text{mass H} = 7.83 \% \times 100 \text{ g} = 7.83 \text{ g}$$

$$\text{mass O} = 22.21 \% \times 100 \text{ g} = 22.21 \text{ g}$$

We convert the mass of each element to the corresponding number of moles by dividing by the atomic weight of that element. Thus, we obtain the following number of moles of each element:

$$\text{moles C} = 69.96 \text{ g} / 12.0 \text{ g/mole} = 5.83 \text{ moles}$$

$$\text{moles H} = 7.83 \text{ g} / 1.01 \text{ g/mole} = 7.75 \text{ moles}$$

$$\text{moles O} = 22.21 \text{ g} / 16.0 \text{ g/mole} = 1.39 \text{ moles}$$

The greatest common factor is 1.39 moles. Dividing the number of moles of each element by 1.39 moles, we obtain

$$\frac{\text{moles C}}{1.39 \text{ moles}} = \frac{5.83 \text{ moles}}{1.39 \text{ moles}} \approx 4.2$$

$$\frac{\text{moles H}}{1.39 \text{ moles}} = \frac{7.75 \text{ moles}}{1.39 \text{ moles}} \approx 5.6$$

$$\frac{\text{moles O}}{1.39 \text{ moles}} = \frac{1.39 \text{ moles}}{1.39 \text{ moles}} = 1.$$

Multiplying these numbers by 5 in order to get whole numbers we obtain:

$$\text{Proportion of C atoms} = 4.2 \times 5 = 21$$

$$\text{Proportion of H atoms} = 5.6 \times 5 = 28$$

$$\text{Proportion of O atoms} = 1 \times 5 = 5.$$

Thus, the molecular formula of this compound is $\text{C}_{21}\text{H}_{28}\text{O}_5$. As a check, we determine the molecular weight of this compound to be

$$\begin{aligned} \text{molecular weight} &= (21 \times \text{atomic weight of C}) + (28 \times \\ &\quad \text{atomic weight of H}) + (5 \times \text{atomic} \\ &\quad \text{weight of O}) \\ &= (21 \times 12.0 \text{ g/mole}) + (28 \times 1.01 \text{ g/mole}) \\ &\quad + (5 \times 16.0 \text{ g/mole}) \end{aligned}$$

$$\approx 360 \text{ g/mole,}$$

which is the experimentally determined molecular weight.

• PROBLEM 144

The most common constituent of gasoline is iso-octane. It is a hydrocarbon, composed by weight of 84.12 % carbon, and 15.88 % hydrogen. Given that it contains 5.27×10^{21} molecules per gram, what is its molecular formula?

Solution: The molecular formula for iso-octane is $C_n H_m$, where n is the number of moles of C and m is the number of moles of H present in 1 mole of iso-octane. One can find n and m from the molecular weight of the compound. It is given that there are 5.27×10^{21} molecules/g present in one gram of the compound. One knows that there are 6.02×10^{23} molecules per mole. Thus,

$$MW = \frac{6.02 \times 10^{23} \text{ molecules/mole}}{5.27 \times 10^{21} \text{ molecules/g}} = 114.23 \text{ g/mole}$$

The C in this compound weighs 84.12 % of 114.23 g and the H weighs 15.88 % of 114.23 g.

$$\text{weight of C} = .8412 \times 114.23 \text{ g/mole} = 96.09 \text{ g/mole}$$

$$\text{weight of H} = .1588 \times 114.23 \text{ g/mole} = 18.14 \text{ g/mole.}$$

The number of moles of carbon present, n , is equal to 96.09 g divided by 12.01 g/mole, its atomic weight.

$$n = \frac{96.09 \text{ g}}{12.01 \text{ g/mole}} = 8.00 \text{ moles}$$

The number of moles of H, m , can be calculated in a similar manner.

$$m = \frac{18.14 \text{ g}}{1 \text{ g/mole}} = 18.14 \text{ mole,}$$

where 1 g/mole is the atomic weight of hydrogen.

The formula is $C_8 H_{18}$, 18.14 is rounded off to the nearest whole number.

• PROBLEM 145

Calculate the percentage composition of aluminum sulfate, $Al_2(SO_4)_3$.

Solution: Because the formula of a compound is constant, the percentage composition of each element present can be calculated by using the parts by weight of each element in one molecular weight of the compound. The molecular weight of the compound can be calculated by adding together the weights of the various elements contained in the compound.

There are 2 atoms of aluminum present, and the molecular weight of aluminum is 27, so the contribution of the aluminum to the compound's total molecular weight is found by multiplying 27 by 2.

$$2 \text{ atoms of aluminum weigh } 2 \times 27 = 54$$

The weight contributions of the other elements present can be calculated in the same way.

The molecular weight of sulfur is 32, and there are 3 sulfur atoms present in the compound.

$$3 \text{ atoms of sulfur weigh } 3 \times 32 = 96$$

The molecular weight of oxygen is 16, and there are 12 oxygen atoms present in the compound.

$$12 \text{ atoms of oxygen weigh } 12 \times 16 = 192$$

The weights of the three elements contained in the compound are added together to find the total molecular weight of the compound.

2 atoms of aluminum	54
3 atoms of sulfur	96
12 atoms of oxygen	<u>192</u>
molecular weight of $\text{Al}_2(\text{SO}_4)_3$	342

The percentage composition of each element present can now be calculated by considering what fraction by weight each element is of the total compound.

The weight of two atoms of aluminum is 54 so the fraction by weight of aluminum in the compound can be found by dividing 54 by 342, the molecular weight of the compound. The percentage is then found by multiplying this fraction by 100.

percentage composition of aluminum

$$= \frac{\text{weight of aluminum in compound}}{\text{molecular weight compound}} \times 100$$

$$\text{percentage composition of aluminum} = \frac{54}{342} \times 100 = 15.8 \%$$

The same method can be applied to the sulfur and oxygen.

$$\text{percentage composition of sulfur} = \frac{96}{342} \times 100 = 28.1\%$$

$$\text{percentage composition of oxygen} = \frac{192}{342} \times 100 = 56.1\%$$

If the percent compositions of all the elements in a compound are added together they will equal 100.0.

percent composition of aluminum	15.8
percent composition of sulfur	28.1
percent composition of oxygen	56.1
	<hr/>
	100.0

• PROBLEM 146

Two thirds of the atoms in a molecule of water (H_2O) are hydrogen. What percentage of the weight of a water molecule is the weight of the two hydrogen atoms? The atomic weight of hydrogen is 1.008 g/mole and of oxygen is 16.00 g/mole.

Solution: The most direct way to solve this composition problem is to consider the total weight of one molecule of water:

$$\begin{aligned}\text{mass}_{\text{H}_2\text{O}} &= 2 \text{ mass}_{\text{H}} + \text{mass}_{\text{O}} \\ &= 2(1.008 \text{ g/mole}) + 16.00 \text{ g/mole} \\ &= 18.016 \text{ g/mole}.\end{aligned}$$

The mass of two hydrogen atoms is $2 \times 1.008 = 2.016$ g/mole. Hence, the percentage mass of hydrogen in water is:

$$\begin{aligned}\% \text{ mass} &= \frac{2 \text{ mass}_{\text{H}}}{\text{mass}_{\text{H}_2\text{O}}} \times 100\% = \frac{2.016 \text{ g/mole}}{18.016 \text{ g/mole}} \times 100\% \\ &= 11.19\end{aligned}$$

• PROBLEM 147

Calculate the percent composition by weight of ether ($(\text{C}_2\text{H}_5)_2\text{O}$).

Solution: To understand this problem, one must first realize that the subscript 2 outside the brackets means that all elements inside the brackets are multiplied by 2. Carrying this procedure out, one obtains the formula

$C_4H_{10}O$ for ether.

Percent composition is found by dividing the individual weights of the components by the molecular weight. Therefore, first find molecular weight by using the table below.

Element	Number of Atoms	Atomic Weight	Total Atomic Weight
C	4	12	48
H	10	1	10
O	1	16	16

$$M.W. = 48 + 10 + 16 = 74 \text{ g/mole.}$$

Percent composition is then calculated from the formula

$$\% \text{ composition} = \frac{(\text{total atomic weight})}{M.W.} \times 100$$

$$\% C = \frac{48}{74} \times 100 = 64.86 \%$$

$$\% H = \frac{10}{74} \times 100 = 13.51 \%$$

$$\% O = \frac{16}{74} \times 100 = 21.63 \%$$

To double check, make sure that the total percent equals 100.

• PROBLEM 148

Calculate the percent weight of each element in magnesium chloride, $MgCl_2$.

Solution: As can be seen from the formula, for each molecule of Mg there are 2 molecules of Cl that must combine to form the compound.

The way to solve a percent composition problem is first to find the total molecular weight. Then divide the weight of each component by the total molecular weight to find the percentages.

First set up a table as shown below.

Element	Number of Atoms or Moles	Atomic Weight	Total Atomic Weight
Mg	1	24.3	24.3
Cl	2	35.45	70.9

$$\text{M.W. of MgCl}_2 = 24.3 + 70.9 = 95.2$$

To find percent composition use the formula

$$\% \text{ composition} = \frac{(\text{total atomic weight})}{\text{total molecular weight}} \times 100$$

$$\% \text{ Mg} = \frac{24.3}{95.2} \times 100 = 25.5 \%$$

$$\% \text{ Cl} = \frac{70.9}{95.2} \times 100 = 74.5 \%$$

To double check that the answer is correct, see if the sum of the percentages is 100 %.

• PROBLEM 149

Using the Periodic Table of Elements, find the following for sodium dihydrogen phosphate, NaH_2PO_4 : (a) formula weight, (b) percent composition of oxygen, (c) weight in grams of 2.7 moles, and (d) percentage composition of oxygen in 2.7 moles.

Solution: This problem encompasses work in chemical stoichiometry. With this in mind, you proceed as follows:

(a) The formula weight = molecular weight, which is the sum of atomic masses of all the atoms in the substance. $\text{Na} = 22.98$, $\text{H} = 1.008$, $\text{O} = 15.9999$, and $\text{P} = 30.97$. Thus, formula weight = $\text{Na} + 2\text{H} + \text{P} + 4\text{O} = 22.98 + 2(1.0080) + 30.97 + 4(15.9999) = 119.9$.

(b) Percentage composition of oxygen is

$$\frac{\text{total weight of oxygen in compound}}{\text{total weight of compound}}$$

$$= \frac{4 \text{ oxygen atoms} \times 15.9994 \text{ mass/oxygen atom}}{119.9}$$

$$= .5334 \text{ g of oxygen or } 53.34 \% \text{ by weight in } \text{NaH}_2\text{PO}_4.$$

(c) Mole = $\frac{\text{mass of substance}}{\text{molecular weight}}$. You are given that there are 2.7 moles and you calculated the molecular weight. Thus, the weight in grams of 2.7 moles = $(119.9 \text{ g/mole})(2.7 \text{ moles}) = 324 \text{ g of } \text{NaH}_2\text{PO}_4$.

(d) Following a procedure similar to the one in part (b),

$$\text{percent composition} = \frac{\text{no. of moles of O} \times \text{MW of O}}{\text{weight of comp/no. of moles}} \times 100$$

$$= \frac{4 \text{ moles} \times 15.9 \text{ g/mole}}{324 \text{ g}/2.7 \text{ moles}} \times 100$$

$$= 53.34 \%$$

The percent composition of any element in any compound does not change when the amount of the compound present is changed.

• PROBLEM 150

Calculate the weight of iron in 350 pounds (158.9 kgs) of Fe_2O_3 . First, calculate the percent of iron in the compound.

Solution: The percent composition of iron in the compound Fe_2O_3 can be calculated by first finding the molecular weight of the compound. After the molecular weight of the compound is found, the percent of this weight that is made up by the iron is found by dividing the weight of the iron by the molecular weight of the compound and then multiplying this fraction by 100.

The molecular weight of iron is 55.8, so that two atoms of iron weigh $2 \times 55.8 = 111.6$.

The molecular weight of oxygen is 16.0, so that 3 atoms of oxygen weigh $3 \times 16.0 = 48.0$

The molecular weight of Fe_2O_3 is found by adding the weights of the iron and oxygen together.

weight of iron	111.6
weight of oxygen	48.0
molecular weight of Fe_2O_3	159.6

The percent composition of iron =

$$\frac{\text{weight of iron in compound}}{\text{molecular weight of compound}} \times 100$$

$$\text{percent composition of iron} = \frac{111.6}{159.6} \times 100 = 69.9$$

Since the percent composition by weight of iron in the compound is known, the weight of iron in 350 pounds of this compound can now be found. Because the percent composition of iron in this compound is 69.9, the percent of iron in 350 pounds of Fe_2O_3 is also 69.9.

$$\text{Weight of iron in 350 lb of } \text{Fe}_2\text{O}_3 = 350 \times 0.699 = 245 \text{ lb.}$$

$$(111.07 \text{ kg})$$

What is the elemental percent composition (by weight) of a mixture that contains 20.0 g of $\text{KAl}(\text{SO}_4)_2$ and 60.0 g of K_2SO_4 ?

Solution: The elemental percent composition by weight is equal to the weight of the element present divided by the total weight of the mixture multiplied by 100.

$$\text{percent composition} = \frac{\text{wt of element}}{\text{total wt}} \times 100$$

The weight of the total mixture is equal to the sum of the weights of the $\text{KAl}(\text{SO}_4)_2$ and the K_2SO_4 .

$$\text{total weight} = 20.0 \text{ g} + 60.0 \text{ g} = 80.0 \text{ g}$$

The weights of the various elements present is found by dividing the molecular weight of the element multiplied by the number of atoms of that particular atom present in the molecule by the molecular weight of the compound, and then multiplying the quotient by the number of grams of the compound present.

Solving for the weights of the elements in $\text{KAl}(\text{SO}_4)_2$:
 MW = 258.12.

$$\text{For K: MW} = 39.1$$

$$\text{weight of K} = \frac{39.1 \text{ g/mole} \times 1}{258.12 \text{ g/mole}} \times 20.0 \text{ g} = 3.03 \text{ g.}$$

$$\text{For Al: MW} = 26.98$$

$$\text{weight of Al} = \frac{26.98 \text{ g/mole} \times 1}{258.12 \text{ g/mole}} \times 20.0 \text{ g} = 2.09 \text{ g.}$$

$$\text{For S: MW} = 32.06$$

$$\text{weight of S} = \frac{32.06 \text{ g/mole} \times 2 \text{ moles}}{258.12 \text{ g/mole}} \times 20.0 \text{ g} = 4.97 \text{ g.}$$

$$\text{For O: MW} = 16$$

$$\text{weight of O} = \frac{16 \text{ g/mole} \times 8 \text{ moles}}{258.12 \text{ g/mole}} \times 20.0 \text{ g} = 9.92 \text{ g.}$$

Solving for the weight of the elements in K_2SO_4 :
 MW = 174.26

$$\text{For K: MW} = 39.1$$

$$\text{weight of K} = \frac{39.1 \text{ g/mole} \times 2 \text{ moles}}{174.26 \text{ g/mole}} \times 60.0 \text{ g} = 26.93 \text{ g}$$

$$\text{For S: MW} = 32.06$$

$$\text{weight of S} = \frac{32.06 \text{ g/mole} \times 1 \text{ mole}}{174.26 \text{ g/mole}} \times 60.0 \text{ g} = 11.04 \text{ g}$$

For O: MW = 16

$$\text{weight of O} = \frac{16 \text{ g/mole} \times 4 \text{ moles}}{174.26 \text{ g/mole}} \times 60.0 \text{ g} = 22.04 \text{ g}.$$

One can find the total weights of the various elements by taking the sum of their weights from the two compounds.

$$\text{Total weight of K} = 3.03 \text{ g} + 26.93 \text{ g} = 29.96 \text{ g}$$

$$\text{Total weight of Al} = 2.09 \text{ g}$$

$$\text{Total weight of S} = 4.97 \text{ g} + 11.04 \text{ g} = 16.01 \text{ g}$$

$$\text{Total weight of O} = 9.92 \text{ g} + 22.04 \text{ g} = 31.96 \text{ g}.$$

One can now determine the elemental percent composition of the mixture.

$$\% = \frac{\text{weight of element}}{\text{total weight of mixture}} \times 100$$

$$\% \text{ of K} = \frac{29.96 \text{ g}}{80.0 \text{ g}} \times 100 = 37.45 \%$$

$$\% \text{ of Al} = \frac{2.09 \text{ g}}{80.0 \text{ g}} \times 100 = 2.61 \%$$

$$\% \text{ of S} = \frac{16.01 \text{ g}}{80.0 \text{ g}} \times 100 = 20.01 \%$$

$$\% \text{ of O} = \frac{31.96 \text{ g}}{80.0 \text{ g}} \times 100 = 39.95 \%$$

• PROBLEM 152

When a piece of magnesium ribbon weighing 0.32 g is burned in oxygen, the resultant oxide weighs 0.53 g. What is the percentage composition of the oxide?

Solution: To find the percent composition of magnesium and oxygen in the oxide, one must first know the weight of the magnesium and of the oxygen making up the compound. The percent composition of magnesium is equal to the weight of the magnesium divided by the weight of the compound multiplied by 100. The percent composition of the oxygen is found in the same manner by substituting the weight of oxygen for the weight of the magnesium. One is told (in this problem) that 0.32 g of magnesium is used, and that the weight of the oxide formed is 0.53 g. The weight of oxygen in the oxide is the difference in the weight of the oxide and the magnesium.

weight of oxygen = 0.53 g - 0.32 g = 0.21 g.

The oxygen in the oxide weighs 0.21 g. The percent composition of the oxide can now be determined.

$$\text{percent Mg in oxide} = \frac{0.32 \text{ g}}{0.53 \text{ g}} \times 100 = 60 \%$$

$$\text{percent O in oxide} = \frac{0.21}{0.53} \times 100 = 40. \%$$

The oxide is therefore 60 % Mg and 40 % O.

• PROBLEM 153

A 0.240 g sample of a compound of oxygen and element X was found by analysis to contain 0.096 g of X and 0.144 g of oxygen. (a) Calculate the percentage composition by weight. (b) Calculate from the above data, three possible atomic weights for X relative to oxygen (at. wt. = 16). (c) What additional information is needed to calculate the true atomic weight of X?

Solution: (a) The percent composition by weight of each element in the compound is found by dividing the weight of that element by the weight of the compound and then multiplying the quotient by 100. For O, therefore, you obtain the following percentage:

$$\frac{.144 \text{ g}}{0.240 \text{ g}} \times 100 = 60 \% \text{ O}$$

For X:

$$\frac{.096 \text{ g}}{0.240 \text{ g}} \times 100 = 40 \% \text{ X}$$

(b) If you assume that the compound is XO , you can solve for the molecular weight of X by using the following ratio. Let x = MW of X.

$$\frac{.144 \text{ g O}}{16.0 \text{ g/mole O}} = \frac{.096 \text{ g}}{\text{X g/mole } x}$$

$$\text{MW X} = 10.7$$

If the compound is taken to be X_2O , there are twice as many X atoms present as O atoms. In solving for the molecular weight of X in this case, use the following ratio

$$\frac{.144 \text{ g}}{\text{weight of 1 mole of O}} = \frac{.096}{\text{weight of 2 moles of X}}$$

Let x = MW of X

$$\frac{.144 \text{ g}}{16.0} = \frac{.096 \text{ g}}{x}$$

$$x = \frac{16.0 \times .096 \text{ g}}{.144 \text{ g}}$$

$$x = 5.35.$$

Assume also that the formula for the compound might be XO_2 . Then, the following ratio should be used: Let x = MW of X.

$$\frac{.144 \text{ g}}{\text{weight of 2 moles of O}} = \frac{.096}{\text{weight of 1 mole of X}}$$

$$\frac{.144 \text{ g}}{32.0} = \frac{.096 \text{ g}}{x}$$

$$x = \frac{.096 \text{ g} \times 32.0}{.144 \text{ g}} = 21.4.$$

(c) To solve for the true atomic weight of X, you must know the actual number of X atoms present per each O atom present.

• PROBLEM 154

A certain solution contains 5 % FeSO_4 . How many pounds of Fe could be obtained from 1 ton of this solution?

Solution: If a solution contains 5 % FeSO_4 , this means that 5 % of the total weight of the solution is FeSO_4 . In this problem, 5 % of a ton is FeSO_4 . To determine the weight of Fe in this amount of FeSO_4 , one must calculate the weight percent of Fe in FeSO_4 . This is done by dividing the molecular weight of Fe by the molecular weight of FeSO_4 and multiplying the quotient by 100.

To solve this problem:

- (1) Determine the percent weight of FeSO_4 .
- (2) Determine the weight of 5 % of a ton
- (3) Determine how much of this weight is Fe.

(1) The molecular weight of a compound is determined by adding together the weight contributed by the elements of which it is composed. MW of Fe = 55.8, MW of S = 32, MW of O = 16.

Thus, for FeSO_4 ,

$$\begin{array}{rcl} 1 \text{ atom of Fe} & 1 \times 55.8 & = 55.8 \\ 1 \text{ atom of S} & 1 \times 32 & = 32 \\ 4 \text{ atoms of O} & 4 \times 16 & = 64 \\ \hline \end{array}$$

$$\text{molecular weight of FeSO}_4 = 151.8$$

$$\begin{aligned} \text{percent weight of Fe in FeSO}_4 &= \frac{\text{weight of Fe}}{\text{weight of FeSO}_4} \times 100 \\ &= \frac{55.8}{151.8} \times 100 = 37 \%. \end{aligned}$$

37 % of FeSO_4 by weight is Fe.

(2) Determining 5 % of a ton. 1 ton = 2000 lbs.

$$5 \% \text{ of a ton} = .05 \times 2000 \text{ lbs} = 100 \text{ lbs.}$$

Thus, 5 % of a ton is 100 lbs.

(3) Determining the weight of Fe in 100 lbs. 37 % of this is Fe.

$$\text{weight of Fe} = .37 \times 100 \text{ lbs} = 37 \text{ lbs} (16.798 \text{ kgs})$$

• PROBLEM 155

Of the total number of atoms in the universe, approximately 93 % are hydrogen (H, atomic weight = 1.0 g/mole) and 7 % are helium (He, atomic weight = 4.0 g/mole). What percentage of the universe by weight is hydrogen?

Solution: The percentage by weight is equal to the total weight of H atoms divided by the total weight of the universe. Suppose that there are N atoms in the universe. Then the number of hydrogen atoms is $93 \% \times N = 0.93 N$ and the number of helium atoms is $7 \% \times N = 0.07 N$. The mass of hydrogen atoms is

$$\text{mass H} = \text{number H} \times \text{mass H} = 0.93 N \times 1.0 \text{ g/mole}$$

and the mass of helium atoms is

$$\text{mass He} = \text{number He} \times \text{mass He} = 0.07 N \times 4.0 \text{ g/mole.}$$

Then, on a per mole basis,

$$\begin{aligned} \% \text{ g/mole} &= \frac{\text{g/mole of H}}{\text{total g/mole}} \times 100 \% \\ &= \frac{0.93 N \times 1.0 \text{ g/mole}}{0.93 N \times 1.0 \text{ g/mole} + 0.07 N \times 4.0 \text{ g/mole}} \times 100 \% \end{aligned}$$

On a weight basis,

$$\begin{aligned}
 \% \text{ H by weight} &= \frac{0.93 \text{ N} \times 1.0 \text{ g/mole}}{0.93 \text{ N} \times 1.0 \text{ g/mole} + 0.07 \text{ N} \times 4.0 \text{ g/mole}} \\
 &\times \frac{1 \text{ mole}}{1 \text{ mole}} \times 100 \% \\
 &= \frac{0.93 \text{ N g}}{0.93 \text{ N g} + 0.28 \text{ N g}} \times 100 \% \\
 &= \frac{0.93 \text{ N g}}{1.21 \text{ N g}} \times 100 \% = \frac{0.93 \text{ g}}{1.21 \text{ g}} \times 100 \% = 77 \%.
 \end{aligned}$$

• PROBLEM 156

The density of a 25.0 % sugar solution is 1.208 g/ml. What weight of sugar would be contained in 1.00 liter of this solution?

Solution: When a solution is said to be 25.0 % sugar, it means that 25.0 % of the weight of the solution is made up by the sugar. Thus one can determine the weight of the sugar in this solution by multiplying the total weight of the solution by .25. Here one is not given the weight of the solution but the density and the volume. The density is the weight of 1 ml of the solution, thus the weight of 1.0 liter or 1000 ml is equal to the volume 1000 ml times the density (1.208 g/ml).

$$\text{weight of solution} = 1000 \text{ ml} \times 1.208 \text{ g/ml} = 1208 \text{ g}.$$

The weight of the sugar in the solution can now be found.

$$\begin{aligned}
 \text{weight of sugar} &= .25 \times \text{weight of solution} \\
 &= .25 \times 1208 \text{ g} = 302 \text{ g}.
 \end{aligned}$$

• PROBLEM 157

A lunar surface probe analyzed a sample of soil and found that 58 % of the atoms it contained were oxygen (O, atomic weight = 16 g/mole), 18 % were silicon (Si, atomic weight = 28 g/mole), 9 % were aluminum (Al, atomic weight = 27 g/mole), and 15 % consisted of other elements with an average atomic weight of 30 g/mole. Determine the percent oxygen by weight in this sample.

Solution: The percent oxygen by weight is equal to the mass of oxygen divided by the total mass, multiplied by 100 %. Hence, in order to determine the mass of O and the total mass, we must determine the mass of each element.

For each element, this is given by

$$\text{mass of element} = \% \text{ of atoms of element} \times \text{atomic weight} \times \text{number of moles in sample.}$$

Consider a sample containing one mole of atoms. We are justified in choosing a sample of definite size since no matter what the size of the sample, the percent composition is the same. Applying the above expression for the mass of each element, we obtain the following:

$$\text{mass O} = 58 \% \times 16 \text{ g/mole} \times 1 \text{ mole} = 9.28 \text{ g}$$

$$\text{mass Si} = 18 \% \times 28 \text{ g/mole} \times 1 \text{ mole} = 5.04 \text{ g}$$

$$\text{mass Al} = 9 \% \times 27 \text{ g/mole} \times 1 \text{ mole} = 2.43 \text{ g}$$

$$\text{mass others} = 15 \% \times 30 \text{ g/mole} \times 1 \text{ mole} = 4.5 \text{ g.}$$

The mass of O in our 1 mole sample is thus 9.28 g and the total mass of our sample is 9.28 g + 5.04 g + 2.43 g + 4.5 g = 21.25 g. Hence, the percent by weight of oxygen in the sample is

$$\frac{9.28 \text{ g}}{21.25 \text{ g}} \times 100 \% \approx 44. \%$$

• PROBLEM 158

What is the weight of 1.0 liter of carbon monoxide (CO) at STP?

Solution: At STP (Standard Temperature and Pressure, 0°C and 760 torr), a mole of any gas has a volume of 22.4 liters. This means that the gram molecular weight of a gas is contained in 22.4 liters. In this problem, one is looking for the weight of 1 liter of CO. The molecular weight of CO is 28 g. Because there are 28 g of CO in 22.4 liters of gas, 28 g must be divided by 22.4 to find the weight of one liter.

$$\text{weight of one liter} = \frac{\text{gram molecular weight}}{22.4}$$

$$\text{weight of 1 liter of CO} = \frac{28 \text{ g}}{22.4} = 1.25 \text{ g.}$$

• PROBLEM 159

Mammalian hemoglobin contains about 0.33 % iron (Fe, atomic weight = 56 g/mole) by weight. If the molecular

weight of hemoglobin is 68,000 g/mole, how many iron atoms are there in each molecule of hemoglobin?

Solution: To solve this problem, we consider a one mole sample of hemoglobin. This assumption introduces no error, since the size of the hemoglobin sample does not affect the number of Fe atoms per molecule of hemoglobin. We then calculate the mass of Fe atoms in one mole of hemoglobin and divide by the atomic weight of Fe to obtain the number of moles of Fe per mole of hemoglobin.

Since the molecular weight of hemoglobin is 68,000 g/mole, the weight of one mole of hemoglobin is

$$68,000 \text{ g/mole} \times 1 \text{ mole} = 68,000 \text{ g.}$$

The weight of Fe in one mole of hemoglobin is then

$$\text{weight Fe} = \% \text{ Fe by weight} \times \text{weight of 1 mole of hemoglobin}$$

$$= 0.33 \% \times 68,000 \text{ g} = 224.4 \text{ g.}$$

The number of moles of Fe corresponding to this weight is found by dividing this weight of Fe by the atomic weight of Fe to obtain

$$\text{moles Fe} = \frac{\text{weight Fe}}{\text{atomic weight Fe}} = \frac{224.4 \text{ g}}{56 \text{ g/mole}} = 4 \text{ moles.}$$

Hence, there are 4 moles of Fe per mole of hemoglobin. To convert from number of moles to number of atoms or molecules, we multiply by Avogadro's number: hence, 4 moles of Fe correspond to $4 \text{ moles} \times 6 \times 10^{23} \text{ atoms/mole} = 24 \times 10^{23} \text{ atoms of Fe}$ and 1 mole of hemoglobin corresponds to $1 \text{ mole} \times 6 \times 10^{23} \text{ molecules/mole} = 6 \times 10^{23} \text{ molecules of hemoglobin}$. Thus, the ratio of number of atoms of Fe to molecules of hemoglobin is

$$\frac{24 \times 10^{23} \text{ atoms of Fe}}{6 \times 10^{23} \text{ molecules of hemoglobin}} = 4 \text{ atoms Fe/molecule hemoglobin}$$

There are 4 atoms of Fe in every molecule of hemoglobin.

• PROBLEM 160

The average bromine content of sea water is 0.0064%.
(a) How much sea water, in cubic feet, would be required to obtain one pound of bromine? (b) What volume of chlorine gas, measured at STP, would be required to liberate the bromine from one ton of salt water? One cubic foot of sea water weighs about 63 pounds (28.35 kg)

Solution: (a) The weight of bromine in one cubic foot of sea water is found by multiplying 63 pounds (the weight of 1 cu.ft. of sea water) by 0.000064. This is because bromine composes 0.0064 % of the weight of sea water.

$$\begin{aligned}\text{weight of Br per 1 cubic ft (liter)} &= .000064 \times 63 \text{ lb} \\ &\quad (1.01 \text{ kg/liter}) \\ &= 4.032 \times 10^{-3} \text{ lb} \\ &\quad (6.46 \times 10^{-5} \text{ kg})\end{aligned}$$

One can find the number of cubic feet of sea water necessary to extract 1 lb of Br by dividing 1 lb by the number of pounds of Br in one cubic foot.

$$\begin{aligned}\text{no. of cubic ft} &= \frac{1 \text{ lb}}{4.032 \times 10^{-3} \text{ lb/cubic ft}} \\ &= 248.02 \text{ cubic feet (7022.93 liters)}\end{aligned}$$

(b) Since chlorine is more active than bromine, the latter may be liberated from its salt by treatment with chlorine.



One mole of Cl_2 will liberate 2 moles of Br^- . Therefore, to find the volume of Cl_2 gas necessary to liberate the bromine in one ton of sea water, one must first calculate the amount of Br^- present in 1 ton of sea water. One is given that 1 cubic foot of sea water weighs 63 pounds. Thus the number of cubic feet of sea water in 1 ton can be found. (1 ton = 2000 lbs).

$$\begin{aligned}\text{no. of cubic feet} &= \frac{2000 \text{ lbs}}{63 \text{ lbs/cubic feet}} = 31.75 \text{ cu.ft.} \\ &\quad (899 \text{ liters})\end{aligned}$$

In the previous section one found that each cubic foot contains 4.032×10^{-3} lb of Br^- , thus one finds the amount of Br^- in 31.75 cu. ft. by multiplying the number of cubic feet by the weight of one cubic foot.

$$\begin{aligned}\text{weight of Br}^- \text{ in 1 ton sea water} &= \\ &= 31.75 \text{ cu.ft.} \times 4.032 \times 10^{-3} \text{ lb/cu.ft.} \\ &= .128 \text{ lb (0.058 kg)}\end{aligned}$$

One finds the number of moles present by dividing .128 lb by the molecular weight in pounds. (MW of Br^- = 80 gr/mole.)

There are 454 g in 1 lb, therefore grams are converted to pounds by multiplying the number of grams by 1 lb/454 g.

$$\text{MW of Br}^- \text{ in lb} = 80 \text{ g/mole} \times \frac{1 \text{ lb}}{454 \text{ g}} = .1762 \text{ lb/mole}$$

One can now find the number of moles of Br^- present in one ton.

$$\text{no. of moles} = \frac{\text{weight of Br}^-}{\text{MW in lbs}}$$

$$\text{no. of moles} = \frac{.128 \text{ lbs}}{.1762 \text{ lbs/mole}} = .73 \text{ moles.}$$

From the equation one knows that $\frac{1}{2}$ as many moles of Cl_2 are needed as Br^- . Therefore, the amount of Cl_2 used is equal to $\frac{1}{2}$ of .73 moles or .365 moles. The volume of one mole of gas at STP (Standard Temperature and Pressure, 0°C and 1 atm) is defined to be 22.4 liters. Therefore, one can find the volume of Cl_2 gas required for the reaction by multiplying the number of moles of gas present by 22.4 liters.

$$\text{volume of Cl}_2 = .365 \times 22.4 \text{ liters} = 8.18 \text{ liters.}$$

• PROBLEM 161

What mass of calcium (Ca, atomic mass = 40.08 g/mole) must be combined with 1.00 g of phosphorus (P, atomic mass = 30.97 g/mole) to form the compound Ca_3P_2 ?

Solution: From the formula of the compound we are trying to form, Ca_3P_2 , we see that the ratio of moles of calcium to moles of phosphorus must be $3/2$. By calculating the number of moles of P in 1.00 g, we can determine the required number of moles of Ca and then convert this to a mass.

The number of moles of P in 1.00 g is

$$\text{moles P} = \frac{\text{mass P}}{\text{atomic mass P}} = \frac{1.00 \text{ g}}{30.97 \text{ g/mole}} = 0.0322 \text{ mole.}$$

Then, using the ratio $\frac{\text{moles Ca}}{\text{moles P}} = \frac{3}{2}$ or moles Ca = $\frac{3}{2}$ moles P, the number of moles of Ca required to combine with 0.0322 mole P in a $3/2$ ratio is

$$\text{moles Ca} = \frac{3}{2} \text{ moles P} = \frac{3}{2} \times 0.0322 \text{ mole} = 0.0483 \text{ mole.}$$

To convert this to a mass we multiply by the atomic mass, obtaining

$$\begin{aligned} \text{mass Ca} &= \text{moles Ca} \times \text{atomic mass Ca} \\ &= 0.0483 \text{ mole} \times 40.08 \text{ g/mole} = 1.94 \text{ g Ca.} \end{aligned}$$

• PROBLEM 162

Aluminum and oxygen react to form Al_2O_3 . This oxide has a

density = 3.97 g/ml and by chemical analysis is 47.1 weight-percent oxygen. The atomic mass of oxygen is 15.9999, what is the atomic mass of aluminum?

Solution: You can answer this question by setting up a proportion between the relative number of atoms in the oxide and the percentages by weight of the atoms in the compound. From Al_2O_3 , you see there must be 2 atoms of Al for every 3 atoms of O. You are told that the atomic mass of oxygen is 15.9999. There exist only 2 elements in Al_2O_3 . Thus, the weight-percent of the aluminum is $100 - 47.1 = 52.9$. You have,

$$\frac{2 \text{ Al}}{3 \text{ O}} = \frac{\text{weight-percent Al}}{\text{weight-percent O}} = \frac{2 \text{ Al}}{3(15.9999)} = \frac{52.9}{47.1}$$

Solving for Al, which is the atomic mass, you obtain

$$\text{Al} = \frac{3}{2} (15.9999) \left(\frac{52.9}{47.1} \right) = 26.9 \text{ g/mole.}$$

• PROBLEM 163

On being heated in air, a mixture of FeO and Fe_3O_4 picks up oxygen so as to be converted completely to Fe_2O_3 . If the observed weight gain is 5.00 percent of the initial weight, what must have been the composition of the initial mixture?

Solution: One should first determine the weight percent increase when the mixture is 100 % FeO or 100 % Fe_3O_4 . 5.0 % increase will be some mixture in between. 5 % will be equal to the sum of the products of the percent weight gained by each compound and the fractions of the mixture that each compound contributes.


The reaction of FeO and O to form Fe_2O_3 is



The percent weight increase for the reaction is found by dividing the weight of Fe_2O_3 by the weight of 2FeO . Weight of 2FeO is 144, weight of Fe_2O_3 is 160.

$$\frac{\text{weight of } \text{Fe}_2\text{O}_3}{\text{weight of } 2\text{FeO}} = \frac{160}{144} = 1.1111$$

The percent weight increase is equal to $(1.1111 - 1.0) \times 100 = 11.11$ %. This means that for each Fe_2O_3 formed by FeO there is an 11.11 % weight increase. One can solve for the weight increase for Fe_3O_4 by a similar method. The reaction is



The weight of $2\text{Fe}_3\text{O}_4$ is 464, the weight of $3\text{Fe}_2\text{O}_3$ is 480.

$$\frac{\text{weight of } 3\text{Fe}_2\text{O}_3}{\text{weight of } 2\text{Fe}_3\text{O}_4} = \frac{480}{464} = 1.0345$$

percent weight increase $= (1.0345 - 1.0) \times 100 = 3.45\%$.
Thus when Fe_2O_3 is formed from Fe_3O_4 , there is a 3.45 % weight increase. The final mixture must have a 5 % weight increase. Let x = fraction of mixture that is FeO .

$$\begin{aligned} \text{Mixture weight gain} &= (\text{fraction FeO}) \times (\text{wt gain percent} \\ &\quad \text{for FeO}) + (\text{fraction Fe}_3\text{O}_4) \times \\ &\quad (\text{wt. gain percent for Fe}_3\text{O}_4) \end{aligned}$$

$$5 = (x)(11.11) + (1 - x)(3.45)$$

$$5 = 11.11x + 3.45 - 3.45x$$

$$1.55 = 7.66x$$

$$\frac{1.55}{7.66} = x$$

$$.2024 = x$$

$$.7976 = 1 - x$$

The initial mixture is, therefore, 20.24 % FeO and 79.76 % Fe_3O_4 .

• PROBLEM 164

Ethanol ($\text{C}_2\text{H}_5\text{OH}$, molecular weight = 46 g/mole) unlike most ingested substances, is absorbed directly by the stomach lining. If 44 g of pure ethanol (4 oz of whiskey or 5.5 oz of a martini) is consumed, the resulting concentration of ethanol in the blood is 0.080 g ethanol/100 ml blood. What percent of the ingested ethanol is in the blood? Assume that the total blood volume of an adult is 7.0 liters.

Solution: We must first calculate the total mass of alcohol in the blood and then determine to what percent of the ingested mass of alcohol this corresponds.

The total mass of ethanol in the blood is equal to the product of the concentration of ethanol and the total blood volume, or 0.080 g ethanol/100 ml blood \times 7.0 liters blood = 0.080 g ethanol/100 ml blood \times 7000 ml blood = 5.6 g ethanol. Note: 7 liters = 7000 ml, since 1 l = 1000 ml. This corresponds to

$$\frac{5.6 \text{ g}}{44 \text{ g}} \times 100 \% = 13 \% \text{ of the ingested ethanol.}$$

It is known that, when exposed to air, beryllium does not corrode but barium does. One explanation is that beryllium (Be) forms a tightly protective oxide coat whereas barium (Ba) does not. The density of BeO = 3.01 g/cc and BaO = 5.72 g/cc, find what happens to the volume per atom when the metals become oxides. The density of Be = 1.86 g/cc and of Ba = 3.598 g/cc.

Solution: To solve this problem, first calculate the volume per atom of Be and Ba, and then compare it with the volume per atom of their oxides. From this comparison, expansion or shrinkage can be determined. The volume per atom of any element can be found by knowing the atomic weight, the number of atoms per mole and the density of the element.

Thus, there are Avogadro's number or 6.02×10^{23} atoms per mole. You are given the densities of the substances involved in this problem. Thus, by substitution you find volume of Be

$$= \frac{MW}{(6.02 \times 10^{23}) (\text{density})} = \frac{9.01 \text{ g/mole}}{6.02 \times 10^{23} \text{ a/mole} \times 1.86 \text{ g/cc}} \\ = 8.05 \times 10^{-24} \text{ cc/atom}$$

From similar computations, you find that

$$\text{BeO} = 1.38 \times 10^{-23} \text{ cc/atom}, \text{ Ba} = 6.34 \times 10^{-23} \text{ cc/atom}, \\ \text{and BaO} = 4.45 \times 10^{-23}.$$

One can find the comparative size of BeO to Be by subtracting the volume per atom of Be from that of BeO. This difference is then divided by the volume of Be and multiplied by 100. This product gives the percentage increase in size of the initial and final atoms when Be is oxidized. A similar process is used for Ba.

For Be:

$$\frac{1.38 \times 10^{-23} \text{ cc} - 8.05 \times 10^{-24} \text{ cc}}{8.05 \times 10^{-24}} \times 100 = 71 \%$$

For Ba:

$$\frac{4.45 \times 10^{-23} \text{ cc} - 6.34 \times 10^{-23}}{6.34 \times 10^{-23}} \times 100 = -30 \%$$

From a comparison of the atomic volumes of Be and BeO, you see that Be expands by 71 % when it forms the oxide. From a comparison of Ba and BaO, you see that Ba shrinks by 30 %, when it forms the oxide.

CHAPTER 5

STOICHIOMETRY/WEIGHT AND VOLUME CALCULATIONS

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 145 to 189 for step-by-step solutions to problems.

This chapter addresses problems of balancing chemical equations and calculating amounts, such as mass, volume, moles, etc., of species which enter chemical reactions. These kinds of calculations are called stoichiometric calculations. The key idea, as in Chapter 4, is to understand that chemical reactions are always written representing the number of molecules (or atoms for the elements) or moles of species which undergo reaction. Atomic or molecular ratios equal mole ratios. Conversion from moles to mass or, for gases, from moles to volume (see Chapters 3 and 4) must be understood to solve these problems.

Chemical reactions are represented by equations showing reactants and products for a particular reaction. The key to balancing chemical reaction equations lies in understanding that elements are conserved in chemical reactions (the exception is nuclear reactions or reactions involving the atomic nucleus. These very special kinds of reactions are discussed in Chapter 19). The same number of moles (or atoms) of each element must appear in both reactants and products. For simple reactions, which include all those in this text, the molecular ratios of reactants are small integers. These molecular ratios are the coefficients in chemical equations and are called the stoichiometric coefficients; they are usually small integers and are usually reduced to the smallest possible integers by dividing each coefficient by common factors.


For example, ammonia (NH_3) can be combined with oxygen (O_2) to form nitric oxide (NO) and water (H_2O). Write the balanced equation for this reaction (see Problem 173).



Conserving elements leads to the following requirements.

$$\begin{array}{ll} \text{For N} & a = c \\ \text{For H} & 3a = 2d \\ \text{For O} & 2b = c + d \end{array}$$

One additional relationship is needed to solve these equations for the four stoichiometric coefficients, a , b , c , and d . This relationship is arbitrary and is usually selected so that a , b , c , and d will be small integers. Simply pick a value of one coefficient — for example let $a = 1$. It follows that $c = a = 1$, $d = 3/2a = 3/2$, and $b = (c + d)/2 = 5/4$. If a , b , c , and d are multiplied by 4 (this is equivalent to initially selecting $a = 4$), the coefficients become



This is the same equation that appears in Problem 173.

In order to solve wt./wt. or vol./vol. or vol./wt. problems, it is critical to first change weights and/or volumes to moles.

The other key solution to problems in this chapter is the method of limiting reactants. Simply stated, when one of the reactants is used up, the reaction stops. Again, examples best illustrate the technique. Suppose a limited amount of ammonia (for example, 100 g) is burned in the atmosphere (21% O_2) which is, compared to 100 g, unlimited in supply. When 100 g of NH_3 is consumed, the reaction stops even though more oxygen is available.

Compute the weight of NO and H_2O produced when 100 g of ammonia is burned in the atmosphere. Note that the molecular weight of ammonia is 17 ($14 + 3 \times 1$). Therefore,

$$100/17 = 5.88 \text{ moles of } \text{NH}_3 \text{ are consumed.}$$

From Equation 5-2 on the previous page, $4/4(5.88) = 5.88$ moles of NO are produced and $6/4(5.88) = 8.82$ moles of H_2O are produced.

5.88 moles NO = 176.4 g (the molecular weight of NO = 30)

8.82 moles H_2O = 158.8 g (the molecular weight of H_2O = 18)

Now compute the volume (at standard conditions) of NH_3 , air, NO, and H_2O in this reaction. From Chapter 3, one mole of any gas at standard temperature and pressure occupies 22.4 liters.

For NH_3 and NO: 5.88 moles (22.4 liters/mole) = 131.7 liters

For air (21% O_2): $5/4(5.88)/.21$ moles (22.4 moles/liter) = 784 liters

For H_2O : 8.82 moles (22.4 liters/mole) = 197.6 liters

Step-by-Step Solutions to Problems in this Chapter, "Stoichiometry/Weight and Volume Calculations"

BALANCING EQUATIONS

• PROBLEM 166


Balance the equations: (a) $\text{Ag}_2\text{O} \rightarrow \text{Ag} + \text{O}_2$,
(b) $\text{Zn} + \text{HCl} \rightarrow \text{ZnCl}_2 + \text{H}_2$, (c) $\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O}$.

Solution: When balancing chemical equations, one must make sure that there are the same number of atoms of each element on both the left and right side of the arrow. For example, $\text{H}_2 + \text{O}_2 \rightarrow \text{H}_2\text{O}$ is not a balanced equation because there are 2 O's on the left side and only one on the right. $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$ is the balanced equation for water because there are the same number of H and O atoms on each side of the equation.


(a) $\text{Ag}_2\text{O} \rightarrow \text{Ag} + \text{O}_2$ is not a balanced equation because there are 2 Ag on the left and only one on the right, and because there is only one O on the left and two O on the right. To balance this equation one must first multiply the left side by 2 to have 2 O's on each side.



There are now 4 Ag on the left and only one on the right, thus the Ag, on the right must be multiplied by 4.



The equation is now balanced.




In this equation, there are 2 H and 2 Cl on the right and only one of each on the left, therefore, the equation can be balanced by multiplying the HCl on the left by 2.



Because there are the same number of Zn, Cl, and H on both sides of the equation, it is balanced.



Here, there are 1 Na, five O, 3 H and 1 S on the left and 2 Na, 1 S, five O, and 2 H on the right. To balance this equation, one can first adjust the Na by multiplying the NaOH by 2.



There are now 2 Na, six O, 4 H, and 1 S on the left and 2 Na, five O, 2 H, and 1 S on the right. Because there are two more H and one more O on the left than on the right, you can balance this equation by multiplying the H_2O by 2.



The equation is now balanced.

• PROBLEM 167


Balance the following by filling in missing species and proper coefficient: (a) $\text{NaOH} + \underline{\hspace{1cm}} + \text{NaHSO}_4 + \text{HOH}$,

(b) $\text{PCl}_3 + \underline{\hspace{1cm}} \text{HOH} + \underline{\hspace{1cm}} + 3\text{HCl}$, (c) $\text{CH}_4 + \underline{\hspace{1cm}} + \text{CCl}_4 + 4\text{HCl}$.

Solution: To balance chemical equations you must remember that ALL atoms (and charges) must be accounted for. The use of coefficients in front of compounds is a means to this end. Thus,



On the right side of the equation, you have 1 Na, 3 H's, 5 O's, and 1 S. This same number of elements must appear on the left side. However, on the left side, there exists only 1 Na, 1 O, and 1 H. You are missing 2 H's, 1 S, and 4 O's. The missing species is H_2SO_4 , sulfuric acid. You could have anticipated this since a strong base (NaOH) reacting with a strong acid yields a salt (NaHSO_4) and water. The point is, however, that H_2SO_4 balances the equation by supplying all the missing atoms.



Here, the left side has 1 P, 3 Cl's, 6 H's, and 3 O's. The right has 3 H's and 3 Cl's. You are missing 1 P, 3 O's and 3 hydrogens. Therefore, P(OH)_3 is formed.



Here, there are 1 C, 8 Cl's, and 4 H's on the right and 1 C and 4 H's on the left. The missing compound, therefore, contains 8 Cl's and thus it is 4Cl_2 . One knows that it is

4 Cl_2 rather than Cl_4 or 8Cl because elemental chlorine gas is a diatomic or 2 atom molecule.

CALCULATIONS USING CHEMICAL ARITHMETIC

• PROBLEM 168

Verify that the following data confirm the law of equivalent proportions: Nitrogen and oxygen react with hydrogen to form ammonia and water, respectively. 4.66 g of nitrogen is required for every gram of hydrogen in ammonia, and 8 g of oxygen for every gram of hydrogen in water. Nitrogen plus oxygen yields NO . Here, 14 g of nitrogen is required for every 16 g of oxygen.

Solution: To verify, you must show that when two elements (nitrogen and oxygen) combine with a third element (hydrogen), they will do so in a simple multiple of the proportions in which they combine with each other. Thus, the nitrogen to oxygen ratio in NO must be a ratio of small integers with the nitrogen and oxygen ratio in H_3N (ammonia) and H_2O (water).

$$\text{For ammonia and water, } \frac{\text{N}}{\text{O}} = \frac{4.66 \text{ g}}{8.00 \text{ g}} = .582$$

$$\text{For NO, } \frac{\text{N}}{\text{O}} = \frac{14 \text{ g}}{16 \text{ g}} = .875.$$

If .582 and .875 are a ratio of small integers to each other, you verify the law of equivalent proportions. Therefore,

$$\frac{.582}{.875} = .665 \approx \frac{2}{3},$$

concluding that they are a ratio.

• PROBLEM 169

In a chemical reaction requiring two atoms of phosphorus for five atoms of oxygen, how many grams of oxygen are required by 3.10 g of phosphorus?

Solution: Because the relationship between the phosphorus and oxygen is given in atoms, the relationship also holds for moles. There must be 2 moles of phosphorus for every 5 moles of oxygen. This is true because there is a set number of atoms in any one mole. Therefore, one must first find the number of moles of phosphorus present. From this, one can find the number of moles of oxygen present. From the number of moles of oxygen, one can find the weight by multiplying by the molecular weight.

The number of moles of phosphorus present is found by dividing the number of grams by the molecular weight (MW = 31).

$$\text{no. of moles} = \frac{3.10 \text{ g}}{31 \text{ g/mole}} = .10 \text{ moles}$$

Because there must be five moles of oxygen for every two moles of phosphorus, the following ratio can be used to determine the number of moles of oxygen. Let x = number of moles of O.

$$\frac{2 \text{ moles P}}{5 \text{ moles O}} = \frac{.10 \text{ moles P}}{x}$$

$$x = \frac{5 \text{ moles O} \times .10 \text{ moles P}}{2 \text{ moles P}} = .25 \text{ moles O}$$

The weight of the oxygen is then found by multiplying the number of moles by the molecular weight (MW = 16).

$$\text{no. of grams of O} = 16 \text{ g/mole} \times .25 \text{ mole} = 4.00 \text{ g O.}$$

• PROBLEM 170

Two atoms of scandium are to combine with three atoms of oxygen. If you start with 1 gram of scandium, how much oxygen is required? Scandium has an atomic weight of 44.96 g/mole. The at. wt. of oxygen is 15.999 g/mole.

Solution: The key to solving this problem is determining the number of moles that will be reacting. After this, one is able to calculate the mass by multiplying the atomic weight by the number of moles.

Two atoms of scandium react with 3 atoms of oxygen, which is equivalent to saying 2 moles of scandium react with 3 moles of oxygen. A mole is defined as weight in grams of the substance divided by the atomic weight or molecular weight.

If you start with 1 gram of scandium, then the number of moles of scandium is

$$\frac{1.00 \text{ g}}{44.96 \text{ g/mole}} = .0222 \text{ moles.}$$

Since the number of moles of oxygen must be in a 3 : 2 ratio to scandium, multiply $3/2$ by .0222 to determine the number of moles of oxygen that will react.

$.0222 \left(\frac{3}{2} \right) = .0333$ moles of oxygen atoms. Recalling the definition of a mole, the mass of the oxygen that reacts is

$$.0333 (15.999) = .533 \text{ g of oxygen.}$$

A metal has an atomic weight of 24. When it reacts with a non-metal of atomic weight 80, it does so in a ratio of 1 atom to 2 atoms, respectively. With this information, how many grams of non-metal will combine with 33.3 g of metal. If 1 g of metal is reacted with 5 g of non-metal, find the amount of product produced.

Solution: To answer this problem, write out the reaction between the metal and non-metal, so that the relative number of moles that react can be determined. You can calculate the number of grams of material that react or are produced. You are told that 1 atom of metal reacts with 2 atoms of non-metal. Let X = non-metal and M = metal. The compound is MX_2 . The reaction is $M + 2X \rightarrow MX_2$. Determine the number of moles that react. You have 33 g of M with an atomic weight of 24.

Therefore, the number of moles = $\frac{33 \text{ g}}{24 \text{ g/mole}} = 1.375$ moles.

The above reaction states that for every 1 mole of M , 2 moles of X must be present. This means, therefore, that $2 \times 1.375 = 2.75$ moles of X must be present. The non-metal has an atomic weight of 80. Thus, recalling the definition

a mole, $2.75 \text{ mole} = \frac{\text{grams}}{80 \text{ g/mole}}$. Solving for grams of X , you obtain $2.75 (80) = 220$ grams.

Let us consider the reaction with 1 g of M with 5 g of X to produce an unknown amount of MX_2 . The solution is similar to the other, except that here you consider the concept of a limiting reagent. The amount of MX_2 produced from a combination will depend on the substance that exists in the smallest quantity. Thus, to solve this problem you compute the number of moles of M and X present. The smaller number, (based on reaction equation) is the one you employ in calculating the number of moles of MX_2 that will be generated. You have, therefore,

$$M_{\text{moles}} = \frac{1}{24} = .04166 \text{ moles } M$$

$$X_{\text{moles}} = \frac{5}{80} = .0625 \text{ moles } X.$$

Using 0.0625 moles X , only .03125 moles of MX_2 will be produced, since the equation informs you that 1 mole of MX_2 is produced for every two moles of X . For M , it is a 1 : 1 ratio, so that .04166 moles of MX_2 would be generated. Therefore, X is the limiting reagent. The atomic weight of MX_2 is 184. Thus, the amount produced is

$$184 \text{ g/mole } (.03125 \text{ moles}) = 5.75 \text{ g.}$$


WEIGHT-WEIGHT PROBLEMS

• PROBLEM 172

Upon the addition of heat to an unknown amount of KClO_3 , .96 g of oxygen was liberated. How much KClO_3 was present?

Solution: The key to answering this question is to write a balanced equation that illustrates this chemical reaction. From this, you can employ the mole concept to determine the weights of the substances involved.

Given that oxygen is liberated and that oxygen gas exists as O_2 and not O , you can write the balanced equation



All atoms are accounted for. The coefficients indicate the relative number of moles that react. For example, every two moles of KClO_3 yield 3 moles of oxygen. A mole is defined as weight in grams of a substance divided by its atomic or molecular weight. Therefore, you have

$$\frac{.96 \text{ g}}{32 \text{ g/mole}} = .030 \text{ moles of oxygen. To calculate the}$$

weight of KClO_3 , you look at the balanced equation.

You find that the number of moles of KClO_3 is $\frac{2}{3}$ the number of moles of oxygen. As such, the number of

moles of $\text{KClO}_3 = .030 \left(\frac{2 \text{ moles } \text{KClO}_3}{3 \text{ moles } \text{O}_2} \right) = .020$ moles of KClO_3 . Recalling the definition of a mole, the number of grams of KClO_3 (MW of $\text{KClO}_3 = 122.55$) is

$$.020 \times (122.55 \text{ g/mole}) = 2.5 \text{ g.}$$

• PROBLEM 173

Given the balanced equation



how many grams of NH_3 will be required to react with 80 g of O_2 ?

Solution: In this problem, one is asked to find the number of grams of NH_3 required to react with 80 g of O_2 . From the equation, one can see that 4 moles of NH_3 react with 5 moles of O_2 . This means that one must first determine the number of moles of O_2 in 80 g. One can then use the following ratio to determine the number

of moles of NH_3 that will react with this number. This ratio holds because the ratio of NH_3 to O_2 is 4 : 5.

$$\frac{4}{5} = \frac{\text{number of moles of } \text{NH}_3}{\text{number of moles of } \text{O}_2}$$

After one knows the number of moles of NH_3 required, one can determine the weight of the NH_3 by multiplying the number of moles by the molecular weight.

To solve this problem:

- (1) calculate the number of O_2 in 80 g
- (2) calculate the number of moles of NH_3 that react with 80 g O_2
- (3) find the number of grams of NH_3 .

Solving:

(1) The number of moles of O_2 in 80 g is found by dividing 80 g by the molecular weight of O_2 . The molecular weight of O is 16, which means the molecular weight of O_2 is twice that amount, or 32.

$$\begin{aligned} \text{number of moles} &= \frac{\text{number of grams present}}{\text{molecular weight}} \\ \text{number of moles} &= \frac{80 \text{ g}}{32 \text{ g/mole}} = 2.5 \text{ moles} \end{aligned}$$

(2) The number of moles of NH_3 reacting with a certain number of moles of O_2 is in the ratio 4 : 5, as shown previously.

$$\begin{aligned} \frac{4}{5} &= \frac{\text{number of moles of } \text{NH}_3}{\text{number of moles of } \text{O}_2} \\ \frac{4}{5} &= \frac{\text{number of moles of } \text{NH}_3}{2.5 \text{ moles}} \end{aligned}$$

$$\text{number of moles of } \text{NH}_3 = \frac{4 \times 2.5 \text{ moles}}{5} = 2 \text{ moles}$$

2 moles of NH_3 are necessary to react with 2.5 moles or 80 g of O_2 .

(3) The number of grams of NH_3 in 2.0 moles is found by multiplying 2.0 moles by the molecular weight of NH_3 , MW of NH_3 = 17.

$$\text{number of grams} = \text{number of moles} \times \text{molecular weight}$$

$$\text{number of grams} = 2.0 \text{ moles} \times 17 \text{ g/mole} = 34 \text{ g.}$$

• PROBLEM 174

In the commercial preparation of hydrogen chloride gas, what weight of HCl in grams may be obtained by heating 234 g. of NaCl with excess H_2SO_4 ?

The balanced equation for the reaction is



Molecular weights: $\text{NaCl} = 58.5$, $\text{HCl} = 36.5$.

Solution: This problem can be solved by using either the mole method or the proportion method.

Mole method: According to the equation, 2 moles of NaCl produce 2 moles of HCl , thus 1 mole of HCl is obtained for every mole of NaCl used.

To use this method, one must first find the number of moles of NaCl reacted because the number of moles of NaCl reacted equals the number of moles of HCl formed.

The number of moles of NaCl reacted can be found by dividing the number of grams used by the molecular weight.

$$\text{number of moles} = \frac{\text{number of grams present}}{\text{molecular weight of NaCl}}$$


$$\text{number of moles} = \frac{234 \text{ g}}{58.5 \text{ g/mole}} = 4 \text{ moles NaCl}$$

Since 4 moles of NaCl are reacted, one knows that 4 moles of HCl will be formed. Thus, the weight of the HCl formed is equal to 4 times the molecular weight of HCl .

$$\text{weight of HCl formed} = \text{number of moles} \times \text{molecular weight}$$

$$\text{weight of HCl formed} = 4 \text{ moles} \times 36.5 \text{ g/mol} = 146 \text{ g HCl}$$

Proportion method: An alternate method of solution to this problem is also possible. In this method, the molecular weights (multiplied by the proper coefficients) are placed below the formula in the equation and the amounts of substances (given and unknown) are placed above.




X is the weight of HCl produced. Solving for X

$$\frac{234 \text{ g}}{117 \text{ gr}} = \frac{X}{73 \text{ g}}$$

$$X = \frac{(234 \text{ g}) \times (73 \text{ g})}{117 \text{ g}} = 146 \text{ g.}$$

A chemist decides to react 2 g of VO (vanadium oxide) with 5.75 g of Fe_2O_3 to produce V_2O_5 and FeO . How many grams of V_2O_5 can be obtained?

Solution: It is extremely important to write out a balanced equation that illustrates this chemical reaction. The balanced equation for this reaction is:



The coefficients tell us the relative number of moles of each reactant and the relative number of moles of each product. For example, 2 moles of VO will react for every three moles of Fe_2O_3 . To find the amount of V_2O_5 , therefore, you must first determine how many moles of V_2O_5 will be generated. To do this you must know the number of moles of the reactants.

$$\begin{aligned}\text{moles of VO} &= \frac{\text{weight in grams}}{\text{molecular weight}} = \frac{2.00 \text{ g}}{66.94 \text{ g/mole}} \\ &= .0299 \text{ moles}\end{aligned}$$

$$\text{moles of Fe}_2\text{O}_3 = \frac{5.75 \text{ g}}{159.69 \text{ g/mole}} = .0360 \text{ moles}$$

According to the equation, the number of moles of VO must be $2/3$ the number of moles of Fe_2O_3 . However, $2/3$ of $.0360 = .0240$, but you only have $.0299$ moles. This means that VO is in excess. In other words, for the amount of Fe_2O_3 present, there exists more VO than will react. As such, the products produced will depend on the number of moles of Fe_2O_3 , and not on VO. Fe_2O_3 is called the limiting reagent. In general, then, if you have two quantities that are known, and want to compute a third, you determine the limiting reagent. The unknown quantity will be dependent upon it only.

As stated earlier, the number of moles of Fe_2O_3 was $.0360$. According to the reaction, for every 3 moles of Fe_2O_3 , one mole of V_2O_5 is generated. Therefore, the number of moles of $\text{V}_2\text{O}_5 = 1/3 (.0360) = .0120$.

Recalling the definition of the mole, you have for V_2O_5 : $.0120 \text{ moles} = \text{no. of grams/molecular weight}$. The molecular weight of $\text{V}_2\text{O}_5 = 181.9$.

Therefore the number of grams of V_2O_5 obtained is
 $.0120 \text{ moles} (181.9 \text{ g/mole}) = 2.18 \text{ g}$.

Determine the weights of CO_2 and H_2O produced on burning 104 g. of C_2H_2 . Molecular weights are $\text{CO}_2 = 44$ and $\text{H}_2\text{O} = 18$. The equation for the reaction is



Solution: Two methods can be used to solve this problem. One is called the mole method and the other the proportion method.


Mole method: According to the equation, 2 moles of C_2H_2 react with 5 moles of O_2 to produce 4 moles of CO_2 and 2 moles of H_2O . Thus, in this problem, the first thing one has to determine is how many moles of C_2H_2 are contained in 104 g.

The molecular weight of C_2H_2 is 26, by dividing the amount of C_2H_2 present by the molecular weight of the compound, the number of moles present is found.

$$\text{number of moles of } \text{C}_2\text{H}_2 = \frac{\text{number of grams of } \text{C}_2\text{H}_2}{\text{molecular weight of } \text{C}_2\text{H}_2}$$

$$\text{number of moles of } \text{C}_2\text{H}_2 \text{ present} = \frac{104 \text{ g}}{26 \text{ g/mole}} = 4 \text{ moles.}$$

In the equation 2 moles of C_2H_2 are burned to form 4 moles of CO_2 and 2 moles of H_2O . Here there are 4 moles of C_2H_2 present, which is twice the amount in the empirical equation. Therefore, twice as much CO_2 and H_2O will be formed. 8 moles of CO_2 and 4 moles of H_2O will be formed.



To find the weight of 8 moles of CO_2 , the molecular weight of CO_2 is multiplied by 8.

$$\text{weight of } \text{CO}_2 = \text{number of moles present} \times \text{molecular weight of } \text{CO}_2.$$

$$\text{weight of } \text{CO}_2 = 8 \text{ moles} \times 44 \text{ g/mole} = 352 \text{ g.}$$

The same procedure can be followed for H_2O .


$$\text{weight of } \text{H}_2\text{O} = \text{number of moles present} \times \text{molecular weight of } \text{H}_2\text{O}$$

$$\text{weight of } \text{H}_2\text{O} = 4 \text{ moles} \times 18 \text{ g/mole} = 72 \text{ g.}$$

Proportion method:

An alternate method of solution to this problem is the proportion method in which molecular weights (multiplied by

the proper coefficients) are placed below the formula in the equation and the amounts of substances (given and unknown) are placed above. Here one has for CO_2 :




This becomes $\frac{104 \text{ g}}{52 \text{ g}} = \frac{X}{176 \text{ g}}$

Solving for X one has

$$X = \frac{(104 \text{ g})(176 \text{ g})}{(52 \text{ g})} = 352 \text{ gr CO}_2$$

A similar method can be applied to the H_2O



$$\frac{104 \text{ g}}{52 \text{ g}} = \frac{X}{36 \text{ g}}$$

$$X = \frac{(104 \text{ g})(36 \text{ g})}{(52 \text{ g})} = 72 \text{ g H}_2\text{O}.$$

• PROBLEM 177

Silver bromide, AgBr , used in photography, may be prepared from AgNO_3 and NaBr . Calculate the weight of each required for producing 93.3 lb of AgBr . (1 lb = 454 g.)

Solution: The reaction for the production of AgBr is written



This means that one mole of each AgNO_3 and NaBr are needed to form one mole of AgBr . In this problem, to determine the number of pounds of AgNO_3 and NaBr used to form 93.9 lbs of AgBr , one must first determine the number of moles of AgBr in 93.9 lbs. There will be one mole of each NaBr and AgNO_3 for each mole of AgBr formed. Once the number of required moles of NaBr and AgNO_3 are found, the weights of these compounds can be determined by multiplying the number of moles by the molecular weight. To solve this problem:

- (1) solve for the number of moles of AgBr in 93.9 lbs
- (2) determine the weights of NaBr and AgNO₃ used.

Solving:

(1) Molecular weights are given in grams, therefore, the grams should be converted to pounds for use in this problem. MW of AgBr = 188 g/mole.

There are 454 grams in one pound, thus grams can be converted to pounds by using the conversion factor 1 lb/454 grams.

$$\text{MW of AgBr in lbs} = 188 \times 1 \text{ lb}/454 \text{ g} = .41 \text{ lbs.}$$

The number of moles in 93.9 lbs can be found by dividing the 93.9 lbs by the molecular weight in pounds.

$$\text{no. of moles of AgBr} = \frac{93.9 \text{ lbs}}{.41 \text{ lbs/mole}} = 229 \text{ moles}$$

Therefore, 229 moles of each NaBr and AgNO₃ are needed to produce 93.9 lbs of AgBr.

(2) The weight of NaBr and AgNO₃ used is equal to the number of moles times the molecular weight. In this problem, one wishes to find the weight in pounds, not grams, thus the molecular weights must be converted to pounds before the conversion factor 1 lb/454 g can be used. MW of NaBr = 103, MW of AgNO₃ = 170.

$$\text{MW of NaBr in lbs} = 103 \text{ g/mole} \times 1 \text{ lb}/454 \text{ g} = .23 \text{ lbs/mole}$$

$$\begin{aligned} \text{MW of AgNO}_3 \text{ in lbs} &= 170 \text{ g/mole} \times 1 \text{ lb}/454 \text{ g} \\ &= .37 \text{ lbs/mole.} \end{aligned}$$

Since it has already been calculated that 229 moles of each of these compounds are needed, one can calculate the weight needed of each by multiplying the number of moles by the molecular weight.

$$\text{weight of NaBr} = .23 \text{ lbs/mole} \times 229 \text{ moles} = 52.7 \text{ lbs.}$$

$$\text{weight of AgNO}_3 = .37 \text{ lbs/mole} \times 229 \text{ moles} = 84.7 \text{ lbs.}$$

• PROBLEM 178


What weight of sulfur must combine with aluminum to form 600 lbs(272.4 kg) of aluminum sulfide?

Solution: In this problem one wants to find out how much reactant (sulfur) was needed to produce a given amount of product (aluminum sulfide). The balanced

equation for this reaction is:



The first method for solving this problem is the proportion method. This procedure involves the ratios of weights and molecular weights. Set up the balanced equation showing the weights and molecular weights.



M.W. of S is 32. M.W. of Al_2S_3 is calculated below.

Atom	Number of Atoms	Atomic Weight	Total Atomic Weight
Al	2	27	54
S	3	32	96

M.W. of Al_2S_3 = 150.

Use the proportion equation,

$$\frac{\text{weight}_{\text{reactant}}}{\text{moles} \times \text{M.W.}_{\text{reactant}}} = \frac{\text{weight}_{\text{product}}}{\text{moles} \times \text{M.W.}_{\text{product}}}$$

Solve for $\text{weight}_{\text{reactant}}$.

$$\begin{aligned} \text{weight}_{\text{reactant}} &= \frac{(3)(32)(600)}{150} \\ &= 384 \text{ lbs. (174.3 kg) sulfur} \end{aligned}$$

The second method for solving the problem is by the mole method. We see that 1 mole of Al_2S_3 requires 3 moles of S. Therefore, if one can calculate the actual number of moles present of 1 substance one can obtain the number of moles of the other substance by setting up a ratio. Then, knowing the number of moles, one can calculate the weight by using the equation $\text{moles} = \text{weight}/\text{M.W.}$

One is given that 600 lbs. of Al_2S_3 are present. Therefore, the number of moles is $600/150 = 4$ pound moles of Al_2S_3 . Setting up the ratio,

$$\frac{\text{S}}{\text{Al}_2\text{S}_3} = \frac{3}{1} = \frac{x}{4}$$


it is seen that $x = 12$ lb. moles of S.

Next, solve the mole equation for weight of S

$$\begin{aligned} \text{weight} &= \text{moles} \times \text{M.W.} = 12 \times 32 = 384 \text{ lbs.} \\ &= 384 \text{ lbs. (174.3 kg)} \end{aligned}$$

Heating of NaNO_3 decomposed it to NaNO_2 and O_2 . How much NaNO_3 would you have to decompose to produce 1.50 g of O_2 ?

Solution: The equation for this reaction is:



This means that for every mole of O_2 produced, 2 moles of NaNO_3 must be decomposed. One is given that 1.50 g of O_2 is formed, thus one should determine the number of moles. The number of moles of NaNO_3 needed will be twice this amount. One can solve for the weight by multiplying the number of moles by the molecular weight of the compound.

The number of moles of O_2 can be found by dividing 1.50 g by the molecular weight of O_2 (MW = 32.0).

$$\text{no. of moles} = \frac{1.50 \text{ g}}{32.0 \text{ g/mole}} = 4.69 \times 10^{-2} \text{ moles.}$$


The number of moles of NaNO_3 needed can be found by multiplying the number of moles of O_2 by 2.

$$\text{no. of moles} = 2 \times 4.69 \times 10^{-2} \text{ moles} = 9.38 \times 10^{-2} \text{ moles.}$$

One finds the weight of NaNO_3 needed by multiplying the number of moles by the molecular weight (MW = 85).

$$\text{weight of NaNO}_3 = 85 \text{ g/mole} \times 9.38 \times 10^{-2} \text{ moles} = 7.97 \text{ g.}$$

Baking powder consists of a mixture of cream of tartar (potassium hydrogen tartrate, $\text{KHC}_4\text{H}_4\text{O}_6$, molecular weight = 188 g/mole) and baking soda (sodium bicarbonate, NaHCO_3 , molecular weight = 84 g/mole). These two components react according to the equation



How much baking soda must be added to 8.0 g of cream of tartar for both materials to react completely?

Solution: From the equation, we know that one mole of NaHCO_3 reacts with one mole of $\text{KHC}_4\text{H}_4\text{O}_6$. Hence, if we convert 8.0 g of $\text{KHC}_4\text{H}_4\text{O}_6$ to moles, we know how many moles of NaHCO_3 must be added. Finally, all we need to do is to convert moles of NaHCO_3 to grams of NaHCO_3 .

In order to convert from grams to moles, we use the relationship $\text{moles} = \text{mass/molecular weight}$. The

number of moles of $\text{KHC}_4\text{H}_4\text{O}_6$ in 8.0 g is

$$\text{moles} = \frac{\text{mass}}{\text{molecular weight}} = \frac{8.0 \text{ g}}{188 \text{ g/mole}} = 4.3 \times 10^{-2} \text{ mole.}$$

Hence, we must add 4.3×10^{-2} mole of NaHCO_3 . Using the relationship $\text{mass} = \text{moles} \times \text{molecular weight}$, we find that 4.3×10^{-2} mole of NaHCO_3 corresponds to

$$4.3 \times 10^{-2} \text{ moles} \times 84 \text{ g/mole} = 3.6 \text{ g.}$$

Hence, 3.6 g of baking soda must be added.

• PROBLEM 181

Some solid CaO in a test tube picks up water vapor from the surroundings to change completely to $\text{Ca(OH)}_2(\text{s})$. An observed total initial weight ($\text{CaO} + \text{test tube}$) of 10.860 g goes eventually to 11.149 g. What is the weight of the test tube?

Solution: The equation for the reaction is



This means that one mole of H_2O reacts with one mole of CaO . The difference in the weights of the test tubes is the weight of the H_2O that the CaO absorbed.

$$\text{weight of H}_2\text{O} = 11.149 - 10.860 = .289 \text{ g.}$$

One should now solve for the number of moles of H_2O because the number of moles of water equals the number of moles of CaO present. From this one can find the weight of the CaO and the test tube. The number of moles equals the number of grams divided by the molecular weight (MW of $\text{H}_2\text{O} = 18.0$).

$$\text{number of moles of H}_2\text{O} = \frac{.289 \text{ g}}{18.0 \text{ g/mole}} = .0161 \text{ moles}$$

Therefore, .0161 moles of CaO were originally present in the test tube. One finds the number of grams by multiplying by the molecular weight of CaO (MW of $\text{CaO} = 56.08$).

$$\begin{aligned} \text{number of grams of CaO} &= .0161 \text{ moles} \times 56.08 \text{ g/mole} \\ &= .900 \text{ g.} \end{aligned}$$

The weight of the test tube is equal to .900 g subtracted from the original weight of the test tube and material.

$$\text{weight of test tube} = 10.860 - .900 = 9.960 \text{ g.}$$

"Hard" water contains small amounts of the salts calcium bicarbonate ($\text{Ca}(\text{HCO}_3)_2$) and calcium sulfate (CaSO_4 , molecular weight = 136 g/mole). These react with soap before it has a chance to lather, which is responsible for its cleansing ability. $\text{Ca}(\text{HCO}_3)_2$ is removed by boiling to form insoluble CaCO_3 . CaSO_4 is removed by reaction with washing soda (Na_2CO_3 , molecular weight = 106 g/mole) according to the following equation:



If the rivers surrounding New York City have a CaSO_4 concentration of 1.8×10^{-3} g/liter, how much Na_2CO_3 is required to "soften" (remove CaSO_4) the water consumed by the city in one day (about 6.8×10^9 liters)?

Solution: We must determine the amount of CaSO_4 present in 6.8×10^9 liters and, from this, the amount of Na_2CO_3 required to remove it.

The number of moles per liter, or molarity, of CaSO_4 corresponding to 1.8×10^{-3} g/liter is obtained by dividing this concentration by the molecular weight of CaSO_4 . Multiplying by 6.8×10^9 liters gives the number of moles of CaSO_4 that must be removed. Hence,

$$\begin{aligned} \text{moles CaSO}_4 &= \frac{\text{concentration (g/liter)}}{\text{molecular weight of CaSO}_4} \times 6.8 \times 10^9 \text{ liters} \\ &= \frac{1.8 \times 10^{-3} \text{ g/liter}}{136 \text{ g/mole}} \times 6.8 \times 10^9 \text{ liters} \\ &= 9.0 \times 10^4 \text{ moles.} \end{aligned}$$

From the equation for the reaction between CaSO_4 and Na_2CO_3 , we see that one mole of CaSO_4 reacts with one mole of Na_2CO_3 . Hence, 9.0×10^4 moles of Na_2CO_3 are required to remove all the CaSO_4 . To convert this to mass, we multiply by the molecular weight of Na_2CO_3 , and obtain

$$\begin{aligned} \text{mass Na}_2\text{CO}_3 &= \text{moles Na}_2\text{CO}_3 \times \text{molecular weight Na}_2\text{CO}_3 \\ &= 9.0 \times 10^4 \text{ moles} \times 106 \text{ g/mole} \\ &= 9.5 \times 10^6 \text{ g} = 9.5 \times 10^6 \text{ g} \times 1 \text{ kg}/1000 \text{ g} \\ &= 9.5 \times 10^3 \text{ kg,} \end{aligned}$$

which is about 10 tons.

How many pounds of air (which is 23.19% O_2 and 75.46% N_2 by weight) would be needed to burn a pound of gasoline by a reaction whereby C_8H_{18} reacts with O_2 to form CO_2 and H_2O ?

Solution: The equation for the reaction is



From the equation, one knows that 12.5 moles of O_2 are needed to burn 1 mole of gasoline. To solve for the number of pounds of air necessary to burn 1 pound of gasoline:

- (1) find the number of moles in 1 pound of C_8H_{18}
- (2) determine the number of moles of O_2 needed
- (3) solve for the number of moles of O_2 in 1 pound of the air
- (4) calculate the number of pounds of air needed.

Solving:

(1) The molecular weight of C_8H_{18} is 114.23 g/mole. There are 453.50 g in 1 lb, thus one can convert from grams to pounds by multiplying the number of grams by 1 lb/453.50 g.

$$\begin{aligned} \text{MW of } C_8H_{18} \text{ in lbs} &= 114.23 \text{ g/mole} \times \frac{1 \text{ lb}}{453.50 \text{ g}} \\ &= 2.52 \times 10^{-1} \text{ lbs/mole.} \end{aligned}$$

One can find the number of moles in 1 lb by dividing 1 lb by the molecular weight of C_8H_{18} in lbs.

$$\text{no. of moles} = \frac{1 \text{ lb}}{2.52 \times 10^{-1} \text{ lbs/mole}} = 3.97 \text{ moles}$$

- (2) One needs 12.5 times as much O_2 as C_8H_{18} .

$$\begin{aligned} \text{no. of moles of } O_2 \text{ needed} &= 12.5 \times \text{no. of moles } C_8H_{18} \\ &= 12.5 \times 3.97 \text{ moles} = 49.63 \text{ moles} \end{aligned}$$

(3) In 1 lb of the air, there is .2319 lb O_2 and .7546 lb N_2 . Thus, to find the number of moles of O_2 in one pound of the air, one must divide .2319 lb by the molecular weight of O_2 in pounds (MW of O_2 = 32 g/mole).

$$\text{MW in lbs of O}_2 = 32 \text{ g/mole} \times \frac{1 \text{ lb}}{453.5 \text{ g}}$$

$$= 7.06 \times 10^{-2} \text{ lb/mole}$$

$$\text{no. of moles} = \frac{.2319 \text{ lb}}{7.06 \times 10^{-2} \text{ lb/mole}} = 3.29 \text{ moles.}$$

(4) There are 3.29 moles of O₂ in one lb and 49.63 moles of O₂ are needed to burn 1 lb of gas.

$$\text{no. of lbs of air needed} = \frac{49.63 \text{ moles}}{3.29 \text{ moles/lb}} = 15.10 \text{ lbs.}$$

• PROBLEM 184

A lunar module used Aerozine 50 as fuel and nitrogen tetroxide (N₂O₄, molecular weight = 92.0 g/mole) as oxidizer. Aerozine 50 consists of 50 % by weight of hydrazine (N₂H₄, molecular weight = 32.0 g/mole) and 50 % by weight of unsymmetrical dimethylhydrazine ((CH₃)₂N₂H₂, molecular weight = 60.0 g/mole). The chief exhaust product was water (H₂O, molecular weight = 18.0 g/mole). Two of the reactions that led to the formation of water are the following:



If we assume that these reactions were the only ones in which water was formed, how much water was produced by the ascent of the lunar module if 2200 kg of Aerozine 50 were consumed in the process?

Solution: Aerozine 50 consists of N₂H₄ and (CH₃)₂N₂H₂. From the first reaction, we see that 2 moles of H₂O are produced per mole of N₂H₄ consumed, and, from the second reaction, we see that 4 moles of H₂O are produced per mole of (CH₃)₂N₂H₂ consumed. Thus, if we determine the number of moles of N₂H₄ and the number of moles of (CH₃)₂N₂H₂ in 2200 kg of Aerozine 50, we can calculate the number of moles of water, and, from this, the mass of water produced, since moles = grams (mass)/molecular weight.

Since N₂H₄ and (CH₃)₂N₂H₂ each form 50% of Aerozine 50 by weight, the mass of each component in 2200 kg of Aerozine 50 is

$$\text{mass N}_2\text{H}_4 = 50 \% \times 2200 \text{ kg} = 1100 \text{ kg} = 1.1 \times 10^6 \text{ g.}$$

$$\text{mass (CH}_3)_2\text{N}_2\text{H}_2 = 50 \% \times 2200 \text{ kg} = 1100 \text{ kg} = 1.1 \times 10^6 \text{ g.}$$

(there are 1000g per kg.)

To convert mass to moles, we divide by the molecular weight. Hence, 2200 kg of Aerozine 50 contains

$$\text{moles } \text{N}_2\text{H}_4 = 1.1 \times 10^6 \text{ g} / 32.0 \text{ g/mole} = 3.4 \times 10^4 \text{ moles } \text{N}_2\text{H}_4$$

$$\text{moles } (\text{CH}_3)_2\text{N}_2\text{H}_4 = 1.1 \times 10^6 \text{ g} / 60.0 \text{ g/mole}$$

$$= 1.8 \times 10^4 \text{ moles } (\text{CH}_3)_2\text{N}_2\text{H}_4.$$

3.4×10^4 moles of N_2H_4 produces $2 \times 3.4 \times 10^4 = 6.8 \times 10^4$ moles of H_2O and 1.8×10^4 moles of $(\text{CH}_3)_2\text{N}_2\text{H}_2$ produces $4 \times 1.8 \times 10^4 = 7.2 \times 10^4$ moles of H_2O .




The total number of moles of H_2O produced is $6.8 \times 10^4 + 7.2 \times 10^4 = 1.4 \times 10^5$ moles. To convert this to mass, we multiply by the molecular weight of water. Hence,

$$1.4 \times 10^5 \text{ moles} \times 18.0 \text{ g/mole} = 2.5 \times 10^6 \text{ g}$$

$$= 2.5 \times 10^3 \text{ kg of water were produced.}$$

• PROBLEM 185

It has been found that the following sequence can be used to prepare sodium sulfate, Na_2SO_4 :



If you performed this sequence of reactions, how many moles of Na_2SO_4 could possibly be produced if you start with 1 mole of sulfur? How many moles are possible with 4.5 g of water?

Solution: If you had the general equation,




a moles of A react with b moles of B to produce c moles of C and d moles of D. Thus, if you want to know how many grams of D will be produced, and you know how much A you have, calculate the number of moles of A present. From this, you can determine how many moles of D can be generated, since you know that a moles of A will produce d moles of D. With this in mind, you can proceed to answer these questions.

From the first equation, you see that if you start with 1 mole of sulfur (S), 1 mole of SO_2 can be generated (since all the coefficients are one, although 1 is not

written). Now that 1 mole of SO_2 is generated, you proceed to the second equation. It states that for every 2 moles of SO_2 , 2 moles of SO_3 are generated. Thus, in keeping with this relative ratio, 1 mole of SO_3 can be generated from one mole of SO_2 . In the third reaction you again have 1 mole of SO_3 yielding 1 mole of H_2SO_4 . At this point, you have 1 mole of H_2SO_4 (from 1 mole of starting sulfur). The last equation shows 1 mole of H_2SO_4 producing 1 mole of Na_2SO_4 . Therefore, if you were to start with 1 mole of S, you would obtain 1 mole of Na_2SO_4 .

Now let us consider H_2O . Water does not enter the sequence until the third equation. This means you start with



If you have 4.5 g of H_2O ($\text{MW} = 18 \text{ g/m}$), you possess $4.5 \text{ g} / 18 \text{ g/mole} = .25 \text{ moles H}_2\text{O}$. From the reaction, 1 mole of H_2O yields 1 mole of H_2SO_4 . Thus, .25 moles of H_2SO_4 are generated from .25 moles of water. You have the same 1 : 1 ratio.

The last equation is also a 1:1 ratio. Therefore, if you start with 4.5 grams of H_2O , .25 moles of Na_2SO_4 will be produced.

• PROBLEM 186

When 4.90 g of KClO_3 was heated, it showed a weight loss of 0.384 g. Find the percent of the original KClO_3 that had decomposed.

Solution: To solve this problem, you need to determine how many moles of KClO_3 decomposed. Once this is determined, calculate the initial number of moles of KClO_3 and divide to obtain a percentage. To do this, a balanced equation that illustrates this reaction must be written. Such an equation is



The weight loss of .384 g must be the amount of O_2 liberated. The number of moles of O_2 is this weight divided by its molecular weight, or

$$\frac{.384 \text{ g}}{32 \text{ g/mole}} = .0120 \text{ moles of } \text{O}_2.$$

Going back to the original equation, you find that the number of moles of KClO_3 is $2/3$ that of O_2 . Therefore, the number of moles of KClO_3 that reacted is

$$(.0120 \text{ mol of } \text{O}_2) \left(\frac{2 \text{ mol } \text{KClO}_3}{3 \text{ mol } \text{O}_2} \right) = .0080 \text{ moles of } \text{KClO}_3.$$

The number of moles of KClO_3 that you started with, however, is

$$\frac{4.90 \text{ g}}{122.6 \text{ g/mole}} = .0400 \text{ moles, where } 122.6 \text{ g/mole is}$$

the molecular weight of KClO_3 . Therefore, the percentage decomposition is


$$\frac{.0080}{.0400} \times 100 = 20\%.$$

• PROBLEM 187

An impure sample of aluminum sulfate, $\text{Al}_2(\text{SO}_4)_3$, is analyzed by forming a precipitate of insoluble barium sulfate, BaSO_4 , by reacting aluminum sulfate with an excess of BaCl_2 (to insure complete precipitation). After washing and drying, 2.000 g of BaSO_4 was obtained. If the original sample weighed 1.000 g, what was the per cent of aluminum sulfate in the sample?

Solution: The problem is solved by calculating the amount of BaSO_4 that would form if the sample were pure, and comparing this with the amount actually formed.

$\text{Al}_2(\text{SO}_4)_3$ reacts with BaCl_2 according to the equation



The number of moles of $\text{Al}_2(\text{SO}_4)_3$, assuming the sample is pure, is equal to the mass of the sample (1.000 g) divided by the molecular weight of $\text{Al}_2(\text{SO}_4)_3$, (342.14 g/mole),

$$\frac{1.000 \text{ g}}{342.14 \text{ g/mole}} = 0.00292 \text{ mole } \text{Al}_2(\text{SO}_4)_3.$$

Since 3 moles of BaSO_4 are formed for every mole of $\text{Al}_2(\text{SO}_4)_3$ reacted, as indicated by the coefficients in the reaction, 1.000 g of pure $\text{Al}_2(\text{SO}_4)_3$ would produce $3 \times 0.00292 \text{ mole} = 0.00876 \text{ mole } \text{BaSO}_4$.

The actual number of moles of BaSO_4 formed is equal to the mass of BaSO_4 obtained after washing and drying (2.000 g) divided by the molecular weight of BaSO_4 , (233.40 g/mole),

$$\frac{2.000 \text{ g}}{233.40 \text{ g/mole}} = 0.00857 \text{ mole } \text{BaSO}_4 \text{ formed.}$$

The purity of the sample is then given by the ratio of the number of moles BaSO_4 formed divided by the number of moles of BaSO_4 that would have formed from a pure sample, times 100 %:

$$\text{purity} = \frac{0.00857 \text{ mole}}{0.00876 \text{ mole}} \times 100 \% = 97.8 \%$$

The original sample, therefore, contained 97.8 % $\text{Al}_2(\text{SO}_4)_3$ and $100 \% - 97.8 \% = 2.2 \%$ impurities.

• PROBLEM 188

A silicious rock contains the mineral ZnS . To analyze for Zn, a sample of the rock is pulverized and treated with HCl to dissolve the ZnS (silicious matter is insoluble). Zinc is precipitated from solution by the addition of potassium ferrocyanide $\text{K}_4\text{Fe}(\text{CN})_6$. After filtering, the precipitate is dried and weighed. The reactions which occur are



If a 2 gram sample of rock yields 0.969 gram of $\text{Zn}_2\text{Fe}(\text{CN})_6$, what is the percentage of Zn in the sample? Atomic weight Zn = 65.4, molecular weight $\text{Zn}_2\text{Fe}(\text{CN})_6 = 342.6$.

Solution: In this problem, one wants to find the percent of Zn in the original rock sample. One is told that the rock contains some ZnS but not 100 %. One must find the weight of Zn in $\text{Zn}_2\text{Fe}(\text{CN})_6$ for which we are given the total amount present. Since no Zn is lost during the course of the reaction the final and initial weight of Zn is the same. Then, knowing the weight of Zn in the rock, we can calculate the % of Zn in the rock by dividing the Zn weight by the total weight of the rock.

The first step is to find the % of Zn in $\text{Zn}_2\text{Fe}(\text{CN})_6$. Use the formula:

$$\% \text{ Zn} = \frac{(\text{atomic weight Zn}) (\text{ number of Zn atoms})}{\text{M.W. of } \text{Zn}_2\text{Fe}(\text{CN})_6} \times 100$$

$$= \frac{65.4 (2)}{342.6} \times 100$$

$$= 38.2 \% \text{ or, as a fraction} = 0.382.$$


To find the weight of any part of a compound, multiply the decimal fraction by the total weight of com-

pound. In this case, weight of Zn = (0.382)(0.969 g) = 0.360 g. The weight of Zn is constant; thus % of Zn in rock is calculated by

$$100 \times \frac{\text{weight of Zn}}{\text{total weight of rock}} = \frac{.360}{2} \times 100 = 18 \% \text{ Zn.}$$

• PROBLEM 189

Clay contains 30 % Al_2O_3 , 55 % SiO_2 , and 15 % H_2O . What weight of limestone is required per ton of clay to carry out the following chemical change?



Solution: From the coefficients in the reaction equation, one sees that for every mole of Al_2O_3 and SiO_2 present, 6 moles of CaCO_3 (limestone) are required for the reaction to occur. As such, to determine how much limestone is necessary to react with 1 ton of clay, compute the number of moles of Al_2O_3 and SiO_2 present. From this figure, the number of moles of CaCO_3 can be found. Because a mole is defined as grams divided by molecular weight (MW), the weight (grams) can be found once the molecular weight is calculated.

It is given that 1 ton or 2000 lb of clay will be used. Because clay contains 30 % Al_2O_3 and 55 % SiO_2 , $.30 \times 2000 = 600$ lbs of Al_2O_3 and $.55 \times 2000 = 1100$ lbs of SiO_2 are present in one ton of clay. To calculate the mole amounts, one must convert lbs to grams. This can be done by using the conversion factor 454 g/lb. Thus, $600 \text{ lbs} \times 454 \text{ g/lb} = 2724 \times 10^2$ grams of Al_2O_3 and $1100 \times 454 \text{ g/lb} = 4994 \times 10^2$ grams of SiO_2 .

The MW of $\text{Al}_2\text{O}_3 = 102$ grams/mole. The MW of $\text{SiO}_2 = 60$ grams/mole. Therefore,

$$\text{moles of } \text{Al}_2\text{O}_3 = \frac{2724 \times 10^2 \text{ g}}{102 \text{ g/mole}} = 2.67 \times 10^3 \text{ moles}$$

$$\text{and moles of } \text{SiO}_2 = \frac{4994 \times 10^2 \text{ g}}{60 \text{ g/mole}} = 8.32 \times 10^3 \text{ moles}$$


Notice, in one ton of clay, the mole amounts of SiO_2 and Al_2O_3 are not equal. The mole amount of limestone required will be six times the mole amount of Al_2O_3 , 2.67×10^3 , and not SiO_2 . The reason stems from the fact that Al_2O_3 is the limiting reagent. The amount of any reagent required (or product produced) depends only on the limiting reagent. SiO_2 , with a mole of 8.3×10^3 , is in excess; the amount of limestone required will not depend on it.

Consequently, moles of limestone = $6(\text{moles of Al}_2\text{O}_3) = 6(2.67 \times 10^3) = 1.60 \times 10^4$ moles. The molecular weight of CaCO_3 (limestone) = 100 g/mole. Grams of CaCO_3 required per ton of clay = (moles of CaCO_3)(MW of CaCO_3)

$$= (1.60 \times 10^4)(100) = 1.60 \times 10^6.$$

• PROBLEM 190


A chemist has a mixture of KClO_3 , KHCO_3 , K_2CO_3 , and KCl . She heats 1,000 g of this mixture and notices that the following gases evolve: 18 g of water (H_2O), 132 g of CO_2 , and 40 g of O_2 according to the following reactions:



The KCl is inert under these conditions. Assuming complete decomposition, determine the composition of the original mixture.


Solution: The solution of this problem involves the use of the mole concept and the ability to employ it using chemical (balanced) equations. You need to determine the number of moles of the gases generated from the masses given and their molecular weights. A mole = mass in grams/molecular weight. Once this is known, you can calculate the number of moles of substances in the mixture that had to exist to produce the given quantities. You proceed as follows:

The only source of the 18 g of H_2O is from



The molecular weight of $\text{H}_2\text{O} = 18$. Thus, you have $18 \text{ g}/18 \text{ g/mole} = 1$ mole of H_2O . But the equation states that for every mole of H_2O , you originally had 2 moles of KHCO_3 . Therefore, the mixture must have had 2 moles of KHCO_3 . The molecular weight of KHCO_3 is 100.1 g. Therefore, the weight in grams of it was $2(100.1) = 200.21 \text{ g}$.

The O_2 is generated from only the reaction



You have 40 g of O_2 evolved. Molecular weight of $\text{O}_2 = 32$. Thus you have $40/32 = 1.25$ moles of O_2 . According to the equation, for every 3 moles of O_2 produced, there existed 2 moles of KClO_3 . Thus, you have $2/3(1.25)$ or .833 moles of KClO_3 . Molecular weight of $\text{KClO}_3 = 122.6$. Therefore, the number of grams = $(122.6)(.833) = 102.1 \text{ g}$.

The CO_2 gas has two sources:



From the water evolved, you already know that you have 200.21 g of KHCO_3 . Its molecular weight is 100.1g. Thus, the number of moles of it is, $200.2/100.1 = 2$. From the equation, however, you see that for every 2 moles of KHCO_3 , you obtain 2 moles of CO_2 . Maintaining this one to one ratio, 2 moles of CO_2 must be generated. The molecular weight of CO_2 is 44. Thus, from KHCO_3 , $2(44) = 88$ g of CO_2 produced. The total number of grams CO_2 evolved was given as 132 g. This means, therefore, that the other source of CO_2 , K_2CO_3 , must give off $132 - 88 = 44$ grams of CO_2 . You have



This equation shows a 1 : 1 mole ratio between K_2CO_3 and CO_2 . You have 44 grams of CO_2 released or 1 mole, since the molecular weight of CO_2 is 44. Therefore, you must have 1 mole of K_2CO_3 in the mixture. The molecular weight of K_2CO_3 is 138.21. Thus, the number of grams is (1)(138.21) for $\text{K}_2\text{CO}_3 = 138.21$ g.

In summary, you have 200.14 g of KHCO_3 , 102.12 g of KClO_3 , and 138.21 g of K_2CO_3 . The total mass of these substances = 440.47 g. The original mixture was 1000 g. Thus,

$$1000 - 440.47 = 559.53 \text{ g is the mass of the inert KCl.}$$

REACTIONS WITH LIMITING REAGENTS

• PROBLEM 191

Chromic oxide (Cr_2O_3) may be reduced with hydrogen according to the equation



- (a) What weight of hydrogen would be required to reduce 7.6 g of Cr_2O_3 ? (b) For each mole of metallic chromium prepared, how many moles of hydrogen will be required? (c) What weight of metallic chromium can be prepared from one ton of Cr_2O_3 ? 1 lb = 454 g.

Solution: (a) From the equation for the reaction, one knows that it takes three moles of H_2 to reduce one mole

of Cr_2O_3 . Thus, in solving this problem one should first determine the number of moles of Cr_2O_3 in 7.6 g, then using the ratio

$$\frac{3}{1} = \frac{\text{number of moles of H}_2}{\text{number of moles of Cr}_2\text{O}_3}$$

One can find the number of moles of H_2 necessary to reduce 7.6 g of Cr_2O_3 . After finding the number of moles of H_2 needed, one can obtain the weight by multiplying the number of moles by the molecular weight of H_2 .

(1) Solving for the number of moles of Cr_2O_3 in 7.6 g. This is done by dividing 7.6 g by the molecular weight of Cr_2O_3 (MW = 152).

$$\text{no. of moles} = \frac{7.6 \text{ g}}{152 \text{ g/mole}} = .05 \text{ moles}$$

(2) determining the number of moles of H_2 necessary. The ratio

$$\frac{3}{1} = \frac{\text{number of moles of H}_2}{\text{number of moles of Cr}_2\text{O}_3}$$

will be used. This ratio was made using the stoichiometric coefficients of the equation for the reaction

$$\frac{3}{1} = \frac{\text{number of moles of H}_2}{.05 \text{ moles}}$$

$$\text{no. of moles of H} = \frac{.05 \text{ moles} \times 3}{1} = .15 \text{ moles}$$

(3) solving for the weight of H_2 . (MW = 2.) The weight of a compound is found by multiplying the number of moles present by the molecular weight.

$$\text{weight of H}_2 = .15 \text{ moles} \times 2 \text{ g/mole} = .30 \text{ g.}$$

(b) From the equation for the reaction, it is seen that 3 moles of H_2 are needed to form 2 moles of Cr. This means that for every mole of Cr formed $3/2$ this amount of H_2 is needed. Thus, 1.5 mole of H_2 is necessary to form one mole of Cr.

(c) Using the equation for the reaction, one is told that for every mole of Cr_2O_3 , reduced 2 moles of Cr are formed. Thus, one must determine the number of moles in 1 ton of Cr_2O_3 ; there will be twice as many moles of Cr formed. After one knows the number of moles of Cr formed, one can determine its weight by multiplying the number of moles by the molecular weight of Cr.

(1) Determining the number of moles of Cr_2O_3 in 1 ton of the compound (1 ton = 2000 lbs). The number of moles is found by dividing the weight of the Cr_2O_3 present by its molecular weight. Because the molecular weight is given in grams per mole, it must be converted to pounds per mole

before using it to determine the number of moles present. There are 454 g in one pound, thus grams can be converted to pounds by multiplying the number of grams by the conversion factor 1 lb/454 g. (MW of Cr_2O_3 = 152.)

$$\begin{aligned}\text{MW of } \text{Cr}_2\text{O}_3 \text{ in lbs} &= 152 \text{ g/mole} \times 1 \text{ lb}/454 \text{ g} \\ &= .33 \text{ lbs/mole.}\end{aligned}$$

(2) determining the number of moles of Cr_2O_3 in 1 ton. The number of moles present can be found by dividing 1 ton (2000 lbs) by the molecular weight in pounds.

$$\text{number of moles} = \frac{2000 \text{ lbs}}{.33 \text{ lbs/mole}} = 6060 \text{ moles}$$

Thus, there are twice this many moles of Cr produced.

$$\text{number of moles of Cr} = 2 \times 6060 \text{ moles} = 12120 \text{ moles}$$

(3) finding the weight of Cr formed. One knows that 12,120 moles of Cr are produced. To find the weight of this quantity, the number of moles is multiplied by the molecular weight. To find the weight in pounds, one must first convert the molecular weight from grams to pounds. This is done by multiplying the molecular weight by the conversion factor 1 lb/454 g. This is used because there are 454 g in 1 lb (MW of Cr = 52).


$$\begin{aligned}\text{MW of Cr in pounds} &= 52 \text{ g/mole} \times 1 \text{ lb}/454 \text{ g} \\ &= 0.11 \text{ lb/mole.}\end{aligned}$$

The weight of the Cr formed is now found by multiplying this molecular weight by the number of moles present.

$$\text{weight} = 0.11 \text{ lb/mole} \times 12120 \text{ moles} = 1333 \text{ lbs.}$$

• PROBLEM 192

When 10.0 g of silicon dust, Si, is exploded with 100.0 g of oxygen, O_2 , forming silicon dioxide, SiO_2 , how many grams of O_2 remain uncombined? The reaction equation is



Solution: From the equation, it can be seen that Si and O_2 react in a 1 : 1 ratio. This means that 1 mole of Si will react with 1 mole of O_2 . To determine the amount of O_2 left unreacted after the reaction is performed, calculate the number of moles of O_2 and Si present and then subtract the number of moles of Si from the number of moles of O_2 . (MW of Si = 28, MW of O_2 = 32.)

$$\text{number of moles} = \frac{\text{number of grams}}{\text{MW}}$$

$$\text{number of moles of O}_2 = \frac{100.0 \text{ g}}{32 \text{ g/mole}} = 3.12 \text{ moles}$$

$$\text{number of moles of Si} = \frac{10.0 \text{ g}}{28 \text{ g/mole}} = 0.357 \text{ moles}$$


$$\text{number of moles of excess O}_2 = 3.12 - 0.357 = 2.763 \text{ moles}$$

$$\text{weight of excess O}_2 = \text{number of moles} \times \text{MW}$$

$$= 2.763 \text{ moles} \times 32 \text{ g/mole} = 88.5 \text{ g.}$$

• PROBLEM 193

How many moles of Al_2O_3 can be formed when a mixture of 0.36 moles of aluminum and 0.36 moles of oxygen is ignited? Which substance and how much of it is in excess of that required?



Solution: In this reaction 4 moles of Al and 3 moles of O_2 form 2 moles of Al_2O_3 . One is told that 0.36 moles of Al and of O_2 are available for the reaction. One can see from the equation that a greater number of moles of Al are needed for the reaction than O_2 . Thus, one should assume that all 0.36 moles of Al will be used, but not all 0.36 moles of the O_2 . Since 4 moles of Al are needed for every 3 moles of O_2 , the following ratio holds:

$$\frac{4}{3} = \frac{\text{number of moles of Al}}{\text{number of moles of O}_2}$$

If 0.36 moles of Al are used, one can solve for the number of moles of O_2 that are needed to react with them.

$$\frac{4}{3} = \frac{0.36 \text{ moles}}{\text{number of moles of O}_2}$$

$$\text{number of moles of O}_2 = \frac{3 \times 0.36}{4} = 0.27 \text{ moles}$$

Since only 0.27 moles of O_2 are needed, and there are 0.36 moles present, there is an excess of 0.09 moles of O_2 .

From the reaction, one knows that there are 2 moles of Al_2O_3 formed for every 4 moles of Al reacted. Therefore, a ratio can be set up to determine the number of moles of Al_2O_3 formed from 0.36 moles of Al.

$$\frac{4}{2} = \frac{\text{number of moles of Al}}{\text{number of moles of Al}_2\text{O}_3}$$

If 0.36 moles of Al are reacted, one can determine the number of moles of Al_2O_3 formed.

$$\frac{4}{2} = \frac{0.36 \text{ moles}}{\text{number of moles of Al}_2\text{O}_3}$$

$$\text{number of moles of Al}_2\text{O}_3 = \frac{0.36 \times 2}{4} = 0.18 \text{ moles}$$

Note: One does not determine the moles of Al_2O_3 from O_2 , since the latter is in excess.

• PROBLEM 194

Suppose the change $\text{HC}_2\text{O}_4^- + \text{Cl}_2 \rightarrow \text{CO}_3^{2-} + \text{Cl}^-$ is to be carried out in basic solution. Starting with 0.10 mole of OH^- , 0.10 mole of HC_2O_4^- , and 0.05 mole of Cl_2 , how many moles of Cl^- would be expected to be in the final solution?

Solution: The equation for this reaction is




From this equation one knows that 1 mole of HC_2O_4^- , 1 mole of Cl_2 , and 5 moles of OH^- are needed to form 2 moles of Cl^- .

One needs 5 times as much OH^- as HC_2O_4^- or Cl_2 . This means that if one wishes to react 0.10 moles of HC_2O_4^- or Cl_2 there must be 0.50 moles of OH^- present. There are 0.10 moles of OH^- present, thus one needs only .02 moles of each HC_2O_4^- and Cl_2 . Some of these reactants will remain unchanged. From the equation, one notes that there are 2 moles of Cl^- formed for each Cl_2 reacted. If .02 moles of Cl_2 are used then .04 moles of Cl^- will be formed. OH^- is called the limiting reagent.

• PROBLEM 195

What is the maximum weight of SO_3 that could be made from 25.0 g of SO_2 and 6.00 g of O_2 by the following reaction?



Solution: From the reaction, one knows that for every 2 moles of SO_3 formed, 2 moles of SO_2 and 1 mole of O_2 must react. Thus, to find the amount of SO_3 that can be formed, one must first know the number of moles of SO_2 and O_2 present. The number of moles is found by dividing

the number of grams present by the molecular weight.

$$\text{number of moles} = \frac{\text{number of grams}}{\text{MW}}$$

For O_2 : MW = 32

$$\text{no. of moles} = \frac{6.0 \text{ g}}{32.0 \text{ g/mole}} = 1.88 \times 10^{-1} \text{ moles}$$

For SO_2 : MW = 64.

$$\text{no. of moles} = \frac{25.0 \text{ g}}{64.0 \text{ g/mole}} = 3.91 \times 10^{-1} \text{ moles.}$$

Because 2 moles of SO_2 are needed to react with 1 mole of O_2 , 3.76×10^{-1} moles of SO_2 will react with 1.88×10^{-1} moles of O_2 . This means that $3.91 \times 10^{-1} - 3.76 \times 10^{-1}$ moles or $.15 \times 10^{-1}$ moles of SO_2 will remain unreacted. In this case, O_2 is called the limiting reagent because it determines the number of moles of SO_3 formed. There will be twice as many moles of SO_3 formed as there are O_2 reacting.


$$\begin{aligned} \text{no. of moles of } \text{SO}_3 \text{ formed} &= 2 \times 1.88 \times 10^{-1} \text{ moles} \\ &= 3.76 \times 10^{-1} \text{ moles.} \end{aligned}$$

The weight is found by multiplying the number of moles formed by the molecular weight (MW of SO_3 = 80).

$$\text{weight of } \text{SO}_3 = 3.76 \times 10^{-1} \text{ moles} \times 80 \text{ g/mole} = 30.1 \text{ g.}$$

• PROBLEM 196

A chemist reacts ferric sulfate with barium chloride and obtains barium sulfate and ferric chloride. He writes the following balanced equation to express this reaction:



(A) How much BaCl_2 should be used to react with 10 grams of $\text{Fe}_2(\text{SO}_4)_3$? (B) How much $\text{Fe}_2(\text{SO}_4)_3$ will be necessary to produce 100 g of BaSO_4 ? (C) From a mixture of 50 g of $\text{Fe}_2(\text{SO}_4)_3$ and 100 g of BaCl_2 , how much FeCl_3 can be produced?

Solution: To answer these questions, you must understand the mole concept, and how it is used to calculate the amount of material required in a chemical reaction.

A mole is defined as the number of grams of a substance divided by its molecular weight. In other words,

$$\text{a mole} = \frac{\text{amount in grams of substance}}{\text{molecular weight of substance}}$$

In the given equation:



the numbers before each compound are termed coefficients (the coefficient of $\text{Fe}_2(\text{SO}_4)_3$ is one, and by convention is not written). The equation is balanced. This means that all of the elements are equal in number on both sides of the equation. For example, we have a total of 12 oxygen atoms on the left and 12 on the right. Before doing any problem involving a chemical equation, we must always balance it.

All the atoms must be accounted for. The coefficients serve this purpose. After balancing, they also tell you the relative mole amounts that will react. In other words, in this reaction 1 mole of $\text{Fe}_2(\text{SO}_4)_3$ reacts with 3 moles of BaCl_2 . For the reaction to occur, we must have a mole ratio of 1 : 3 between $\text{Fe}_2(\text{SO}_4)_3$ and BaCl_2 . If such a condition exists, then the equation tells us that 3 moles of BaSO_4 will be produced per mole of $\text{Fe}_2(\text{SO}_4)_3$. In addition, 2 moles of FeCl_3 will also be produced. Therefore, the coefficients tell the relative number of moles of each substance that must be either present or produced.

With this information, we can now answer the questions.

(A) We have 10 grams of $\text{Fe}_2(\text{SO}_4)_3$, and want the number of grams of BaCl_2 that are required to react with it. The molecular weight of $\text{Fe}_2(\text{SO}_4)_3$ is 399.88 g/mole. This number is obtained by adding up all the atomic weights of the individual atoms in the formula. Recalling the definition of a mole, we have 10/399.88 moles of $\text{Fe}_2(\text{SO}_4)_3$.

The equation tells us that 3 moles of BaCl_2 react with one mole of $\text{Fe}_2(\text{SO}_4)_3$. Therefore, we must have 3 times the number of moles of $\text{Fe}_2(\text{SO}_4)_3$, or

$$3 \times \frac{10}{399.88} = \frac{30}{399.88} \quad \text{The number of moles of } \text{BaCl}_2$$

required is, therefore, $\frac{30}{399.88}$. The molecular weight of BaCl_2 is 208.24 g/mole. Recall, mole = number of grams/molecular weight. The number of grams of BaCl_2 required is, therefore,

$$\frac{30}{399.88} \text{ moles} \times 208.24 \text{ g/mole} = 15.62 \text{ grams}$$

(B) We want to produce 100 grams of BaSO_4 . The molecular weight of BaSO_4 is 233.40 g/mole. Therefore, we want to produce 100/233.40 moles of BaSO_4 . How much $\text{Fe}_2(\text{SO}_4)_3$ should be used? Again, we must go back to the equation and look at the coefficients to determine the mole requirements. We see that 3 moles of BaSO_4 are pro-

duced for every mole of $\text{Fe}_2(\text{SO}_4)_3$. Therefore, the required number of moles of $\text{Fe}_2(\text{SO}_4)_3$ is $1/3$ the number of moles of BaSO_4 or

$$\frac{1}{3} \times \frac{100}{233.40} = \frac{100}{700.20} \text{ . The molecular weight of}$$

$\text{Fe}_2(\text{SO}_4)_3$ is 399.88 g/mole. Consequently, the number of grams required of $\text{Fe}_2(\text{SO}_4)_3$ is

$$\frac{100}{700.20} \text{ mole} \times 399.88 \text{ g/mole} = 57.11 \text{ grams}$$

(C) In this question, we are working with a mixture of two substances to produce a third. The first thing is to compute the moles:

$$\text{Moles } \text{Fe}_2(\text{SO}_4)_3 = \frac{50.00 \text{ g}}{399.88 \text{ g/mole}} = .125 \text{ moles}$$

$$\text{Moles } \text{BaCl}_2 = \frac{100 \text{ g}}{208.24 \text{ g/mole}} = .48 \text{ moles}$$

Because we are dealing with two substances to yield one, we must also consider their mole ratios as well as with the product FeCl_3 . The equation calls for three moles of BaCl_2 to react with one mole of $\text{Fe}_2(\text{SO}_4)_3$. We have .125 moles of $\text{Fe}_2(\text{SO}_4)_3$. 3 times this amount is .375 moles. However, you have .48 moles of BaCl_2 . This means, therefore, that we have .48 - .375 or .105 moles of BaCl_2 that will not react. In other words, BaCl_2 is in excess. We always take the limiting reagent, which is $\text{Fe}_2(\text{SO}_4)_3$. The amount of FeCl_3 present will reflect the number of moles of $\text{Fe}_2(\text{SO}_4)_3$, and not BaCl_2 .

Because 2 moles of FeCl_3 will be produced for every mole of $\text{Fe}_2(\text{SO}_4)_3$, $.125 \times 2 = .25$ moles of FeCl_3 will be produced. The molecular weight of FeCl_3 is 162.20. Therefore, the maximum weight of FeCl_3 produced is $.25(162.20) = 40.56$ grams.

• PROBLEM 197

Through several successive reactions, a chemist uses carbon, CaO , HCl and H_2O to produce $\text{C}_6\text{H}_4\text{Cl}_2$. Assuming an efficiency of 65 %, how much $\text{C}_6\text{H}_4\text{Cl}_2$ can be produced from 500 grams of carbon? Assume that $1/3$ of the carbon is lost as 3 moles CO .

Solution: In solving this problem, you must account for all carbon atoms and employ the mole concept. You need not be concerned with the actual sequence of reactions nor the roles of CaO and H_2O .

Dichlorobenzene, $\text{C}_6\text{H}_4\text{Cl}_2$, consists of 6 carbon atoms.

You can determine that there were originally 9 moles of carbon present since 3 moles of CO are produced and the carbon present in the CO represents 1/3 of the original amount of carbon present.

A mole is defined as weight in grams/molecular weight. The molecular weight of carbon is 12. You started with 500 grams. Therefore, the number of moles of carbon is $500/12$. It is stated above, however, that for every 9 moles of C, 1 mole of $C_6H_4Cl_2$ was produced. Therefore, the number of moles of $C_6H_4Cl_2$ is 1/9 of the moles of carbon that you started with. Namely, the number of moles of $C_6H_4Cl_2$ is $1/9 \times 500/12$. The problem calls for an efficiency of 65 %. Therefore, we must multiply this number of moles by

$$\frac{65}{100} \quad \text{or,} \quad \frac{1}{9} \times \frac{500}{12} \times \frac{65}{100} .$$


The molecular weight of $C_6H_4Cl_2$ is 147. Recalling the definition of a mole, the weight of $C_6H_4Cl_2$ produced is

$$147 \times \frac{1}{9} \times \frac{500}{12} \times \frac{65}{100} = 442 \text{ grams.}$$

VOLUME-VOLUME PROBLEMS

• PROBLEM 198

Calculate the volume of O_2 necessary to burn 50 liters of CO completely. The balanced reaction is:



Also, calculate the volume of CO_2 formed.

Solution: This is a volume-volume problem and the technique shown applies only to gases. It is assumed that all the gases must be at the same temperature and pressure. No molecular weights or molecular volumes are needed. One solves this problem by setting up a ratio between the actual volumes present and the mole volumes of the reacting compounds as shown in the equation below.

$$\frac{\text{volume CO present}}{\text{mole volume CO}} = \frac{\text{volume } O_2 \text{ needed}}{\text{mole volume } O_2}$$

The mole ratio for this reaction is 2 : 1, therefore, the mole volume ratio is 2 : 1. Substitute, to obtain

$$\frac{50 \text{ liters}}{2 \text{ liters}} = \frac{\text{volume O}_2}{1 \text{ liter}}$$

$$\text{volume O}_2 = 25 \text{ liters.}$$

To find the volume of CO₂ produced, set up a similar proportion between CO and CO₂ or O₂ and CO₂. The mole ratio for CO and CO₂ is 1 : 1 and for O₂ and CO₂ is 1 : 2. Therefore use the equation

$$\frac{\text{volume reactant}}{\text{mole volume reactant}} = \frac{\text{volume CO}_2}{\text{mole volume CO}_2}$$

The values for CO as the reactant are:

$$\frac{50}{2} = \frac{\text{volume CO}_2}{2}$$

$$\text{volume CO}_2 = 50 \text{ liters.}$$


The values of O₂ as the reactant are:

$$\frac{25}{1} = \frac{\text{volume CO}_2}{2}$$

$$\text{volume CO}_2 = 50 \text{ liters.}$$

• PROBLEM 199

In the Ostwald process for the commercial preparation of nitric acid, ammonia gas is burned in oxygen in the presence of a Pt catalyst. The balanced equation is:



What volume of O₂ and what volume of NO is formed in the combustion of 500 liters of NH₃. All gases are under the same conditions of temperature and pressure.

Solution: In order to solve this volume-volume problem set up a ratio between the volumes involved. No molecular weights are needed in volume-volume problems. First, one wants to find out how much O₂ is required to react with 500 liters of NH₃. As can be seen from the stoichiometry of the reaction, the ratio of the mole volumes (i.e. how much NH₃ reacts with O₂) is 4 : 5. Using the equation

$$\frac{\text{volume NH}_3 \text{ present}}{\text{mole volume NH}_3} = \frac{\text{volume O}_2 \text{ required}}{\text{mole volume O}_2}$$

$$\frac{500 \text{ liters}}{4 \text{ liters}} = \frac{\text{volume O}_2}{5 \text{ liters}}$$

$$\text{volume O}_2 = 625 \text{ liters.}$$

To find the amount of NO produced, set up a similar equation. In this case, 4 mole volumes of NH_3 produce 4 mole volumes of NO as indicated by the stoichiometry of the reaction

$$\frac{\text{volume NH}_3 \text{ present}}{\text{mole volume NH}_3} = \frac{\text{volume NO produced}}{\text{mole volume NO}}$$

$$\frac{500 \text{ liters}}{4 \text{ liters}} = \frac{\text{volume NO produced}}{4 \text{ liters}}$$

$$\text{volume NO produced} = 500 \text{ liters.}$$

It was mentioned in the problem that a Pt catalyst was used. A catalyst is a substance which speeds up the rate at which a reaction occurs by lowering the amount of energy needed to start a reaction (i.e. lowering the activation energy). Catalysts have no effect on the final concentrations of the compounds involved in the reaction. Therefore, the presence of a catalyst does not change the values calculated in this question.

• PROBLEM 200

Calculate the volume of oxygen necessary to burn completely 100 cubic feet (1 cubic foot = 28.316 liters) of butane gas according to the equation



Solution: Since volumes are concerned, the procedure for solving this problem is to set up a ratio between the volumes present and the mole requirements. For this reaction, 2 moles of butane react with 13 moles of O_2 .

When two gases are under the same temperature and pressure conditions, a mole of either gas will occupy the same volume. Therefore, since 13/2 times as many moles of O_2 are required, the volume of O_2 must be 13/2 times that of methane.

Given the volume of methane is 1000 cu. ft., the volume of $\text{O}_2 = \frac{13}{2} (1000) = 6500 \text{ cubic feet.} (1.84 \times 10^5 \text{ liters})$

WEIGHT-VOLUME PROBLEMS

• PROBLEM 201

Glucose-1-phosphate, essential to the metabolism of carbohydrates in humans, has a molecular weight of

260 g/mole and a density of about 1.5 g/cm³. What is the volume occupied by one molecule of glucose-1-phosphate?

Solution: In general, volume = mass/density. Hence, in order to determine the volume, we must determine the mass of one molecule of glucose-1-phosphate.

One mole of glucose-1-phosphate weighs 260 g (this is the meaning of a molecular weight of 260 g/mole). Since there is an Avogadro's number of molecules in a mole (6.02×10^{23} molecules/mole), the mass of one molecule of glucose-1-phosphate is given by

$$\begin{aligned}\text{mass} &= \frac{\text{molecular weight}}{\text{Avogadro's number}} = \frac{260 \text{ g/mole}}{6 \times 10^{23} \text{ molecules/mole}} \\ &= 4.3 \times 10^{-22} \text{ g/molecule.}\end{aligned}$$

Hence, one molecule of glucose-1-phosphate weighs 4.3×10^{-22} g and has a volume of

$$\text{volume} = \frac{\text{mass}}{\text{density}} = \frac{4.3 \times 10^{-22} \text{ g}}{1.5 \text{ g/cm}^3} \approx 2.9 \times 10^{-22} \text{ cm}^3.$$

• PROBLEM 202

What is the mass of 1 liter of carbon monoxide (CO) at standard temperature and pressure (STP).

Solution: At STP, 1 mole of any gas occupies a volume of 22.4 l. The weight of any substance divided by its molecular weight (MW) is equal to the number of moles of that substance. Thus, the weight of 1 mole of any substance weighs its molecular weight (in grams).

The molecular weight of CO = 28 g/mole. Thus, 22.4 liters of CO at STP weigh 28 g. The mass of one liter of CO can then be found from the following proportion

$$\frac{28 \text{ g}}{22.4 \text{ liters}} = \frac{X \text{ grams of CO}}{1 \text{ liter}}$$

Solving for X, $X = 1.25 \text{ g}$.

• PROBLEM 203

What is the weight of 1,000 cubic feet of air at STP?

Solution: To solve this problem, one must first recognize that air is a mixture of 80% N₂ and 20% O₂. One should also know that 22.4 cubic feet of any gas at STP weighs its ounce molecular weight. For example, in this problem,

22.4 ft.³ of O₂ would weigh 32 oz.

22.4 ft.³ of N₂ would weigh 28 oz.

Using this information, if one knows what 22.4 ft.³ of air weighs, then one can determine how much 1000 ft.³ of air weighs.

Since air is a mixture, one can find its ounce molecular weight by multiplying the per cent composition of each gas times each gas' ounce molecular weight and adding these two values:

for N₂ : (0.8) (MW N₂) = (0.8) (28 oz) = 22.4 oz

for O₂ : (0.2) (MW O₂) = (0.2) (32 oz) = 6.4 oz.

The ounce molecular weight for air is 22.4 + 6.4 = 28.8 oz., and has a volume of 22.4 ft.³

If 22.4 ft.³ of air has a weight of 28.8 oz then, the weight of 1000 ft.³ can be found through the following proportion:

$$\frac{22.4 \text{ ft.}^3}{28.8 \text{ oz.}} = \frac{1000 \text{ ft.}^3}{\text{unknown weight}}$$

Solving for the unknown weight of air

$$\text{weight of air} = \frac{(1000 \text{ ft.}^3)(28.8 \text{ oz.})}{(22.4 \text{ ft.}^3)} = 1.29 \times 10^3 \text{ oz.}$$

• PROBLEM 204

Chlorine may be prepared by the action of KClO₃ on HCl, and the reaction may be represented by the equation:



Calculate the weight of KClO₃ which would be required to produce 1.0 liter of Cl₂ gas at STP. R = .082 liter-atm/mole-°K.

Solution: From the equation one can see that 1 mole of KClO₃ reacts to form 3 moles of Cl₂. If one can find the number of moles of KClO₃, which will react to form 1 liter of Cl₂, then one can find its weight by multiplying the number of moles by the molecular weight of KClO₃. STP

(Standard Temperature and Pressure) is defined as 0°C and 1 atm.

One can find the number of moles of Cl_2 in 1 liter by using the Ideal Gas Law

$$n = \frac{PV}{RT},$$

where P is the pressure, V is the volume, R is the gas constant (0.082 liter-atm/mole-°K), T is the absolute temperature and n is the number of moles. Here, T is given in °C; it can be converted to °K by adding 273 to it. You have, then,

$$T = 0 + 273 = 273^\circ\text{K}.$$

One can now solve for the number of moles of Cl_2 produced.

$$\begin{aligned} n &= \frac{PV}{RT} & P &= 1 \text{ atm} \\ & & V &= 1 \text{ liter} \\ & & R &= 0.082 \frac{\text{liter-atm}}{\text{mole-}^\circ\text{K}} \\ & & T &= 273 \end{aligned}$$

$$n = \frac{1 \text{ atm} \times 1 \text{ liter}}{0.082 \frac{\text{liter-atm}}{\text{mole-}^\circ\text{K}} \times 273^\circ\text{K}} = .045 \text{ moles.}$$

From the equation for the reaction, the following ratio is determined.

$$\frac{3}{1} = \frac{\text{number of moles of } \text{Cl}_2}{\text{number of moles of } \text{KClO}_3}$$

One knows that 1 liter contains .045 moles Cl_2 , and if you substitute this value into the above ratio, one can find the number of moles of KClO_3 reacted.

$$\frac{3}{1} = \frac{.045 \text{ moles}}{\text{number of moles of } \text{KClO}_3}$$

$$\text{number of moles of } \text{KClO}_3 = \frac{.045 \text{ moles} \times 1}{3} = .015 \text{ moles}$$

The weight of .015 moles of KClO_3 can be found by multiplying .015 moles by the molecular weight of KClO_3 . (MW of KClO_3 = 122.5.)

$$\text{weight of } \text{KClO}_3 = 122.5 \text{ g/mole} \times .015 \text{ moles} = 1.84 \text{ g.}$$

• PROBLEM 205

Lithium oxide (Li_2O , molecular weight = 30 g/mole) reacts with water (H_2O , molecular weight = 18 g/mole, density = 1.0 g/cm³) to produce lithium hydroxide (LiOH)

according to the following reaction:



What mass of Li_2O is required to completely react with 24 liters of H_2O ?

Solution: One mole of Li_2O reacts with exactly one mole of H_2O , as indicated by the coefficients in the reaction. Hence, if we determine the number of moles of H_2O in 24 liters, we then know the required number of moles of Li_2O , and, from this, can calculate the mass of Li_2O that we need.

The number of moles of water is $\text{moles} = \text{mass}/\text{molecular weight}$. But $\text{mass} = \text{density} \times \text{volume}$, so that the number of moles of water is

$$\begin{aligned}\text{moles} &= \frac{\text{mass}}{\text{molecular weight}} = \frac{\text{density} \times \text{volume}}{\text{molecular weight}} \\ &= \frac{1.0 \text{ g/cm}^3 \times 24 \text{ liters}}{18 \text{ g/mole}} = \frac{1.0 \text{ g/cm}^3 \times 24000 \text{ cm}^3}{18 \text{ g/mole}} \\ &= 1333.33 \text{ moles}\end{aligned}$$

Thus, we require 1333.33 moles of Li_2O to completely react with 24 liters of H_2O (= 1333.33 moles). Multiplying this by the molecular weight of Li_2O , we determine that

$$1333.33 \text{ moles} \times 30 \text{ g/mole} = 40,000 \text{ g} = 40 \text{ kg}$$

of Li_2O is needed.

• PROBLEM 206

What volume of hydrogen at STP is produced as sulfuric acid acts on 120 g. of metallic calcium. Equation for the reaction is



Solution: This problem may be solved by using either the mole method or the proportion method.

Mole method: In using the mole method, one looks at this equation and sees that for every mole of calcium acted on by H_2SO_4 , one mole of hydrogen is produced.

This means that to find how much hydrogen is produced, one must first find out how much calcium is present. There will be the same number of moles of hydrogen produced as there are calcium reacted. After one knows how many moles of hydrogen are produced, one can calculate the volume.


To calculate the number of moles of calcium present, one must divide the amount present by the molecular weight (molecular weight of Ca = 40).

$$\text{number of moles} = \frac{\text{number of grams present}}{\text{molecular weight}}$$

$$\text{number of moles of Ca} = \frac{120 \text{ g}}{40 \text{ g/mole}} = 3.00 \text{ moles}$$

Therefore, 3.00 moles of hydrogen gas are produced. At STP (Standard Temperature and Pressure), the volume of one mole of any gas occupies 22.4 liters. Thus, when 3 moles of gas are generated, as in this problem it occupies $3 \times 22.4 \text{ liters} = 67.2 \text{ liters}$.

Proportion method: In the proportion method, the molecular weights (multiplied by the proper coefficients) are placed below the formula in the equation and the amounts of substances (given and unknown) are placed above. In this case, because one is trying to find the volume of hydrogen and not its weight, the volume of one mole will be placed below the equation as shown.



$$\frac{120 \text{ g}}{40.0 \text{ g}} = \frac{X}{22.4 \text{ liters}}$$

X = unknown volume of H₂ produced.

Solving for X:

$$X = \frac{(22.4 \text{ liters})(120 \text{ g})}{40.0 \text{ g}} = 67.2 \text{ liters.}$$

• PROBLEM 207

The executioner in charge of the lethal gas chamber at the state penitentiary adds excess dilute H₂SO₄ to 196 g (about $\frac{1}{2}$ lb) of NaCN. What volume of HCN gas is formed at STP?

Solution: The equation for this reaction is



This means that for each NaCN reacted, one HCN is formed. After one knows the number of moles of HCN formed, one can find the volume that the gas occupies by multiplying the number of moles by 22.4 liters (volume of one mole of gas at STP). STP (Standard Temperature and Pressure) is de-

fined as 273°K and 1 atm.

The number of moles of NaCN reacted can be found by dividing 196 g by the molecular weight of NaCN. (MW of NaCN = 49.)

$$\text{number of moles} = \frac{196 \text{ g}}{49 \text{ g.mole}} = 4.0 \text{ moles}$$

From the stoichiometry of the reaction, we see there will be 4.0 moles of HCN formed.

The volume of the gas can now be found.

$$\text{volume} = 4.0 \text{ moles} \times 22.4 \text{ liters/mole} = 89.6 \text{ liters.}$$

• PROBLEM 208

A chemist decides to prepare some chlorine gas by the following reaction:



If he uses 100 g of MnO_2 , what is the maximum volume of chlorine gas that can be obtained at standard temperature and pressure (STP)?

Solution: The solution of this problem entails a combination of stoichiometry and gas law theory.

The first thing to determine is how many moles of chlorine will be produced. The volume can be determined using the fact that each mole, at STP, occupies 22.4 l.

Since the equation is balanced, you must look at the coefficients. The equation indicates that for every mole of MnO_2 , 1 mole of chlorine gas is produced. A mole is defined as being equal to weights in grams of material/molecular weight of material.

The molecular weight of MnO_2 is 86.9. It follows, then, that the number of moles of MnO_2 is 100/86.9. This will be equated to the number of moles of chlorine gas produced.

At STP one mole of gas occupies 22.4 liters. However, we do not have 1 mole, but 100/86.9 moles. To calculate the new volume, you multiply the STP volume by the number of moles of chlorine gas produced. In other words, the volume of the chlorine gas is

$$22.4 \times \frac{100}{86.9} = 25.8 \text{ liters.}$$

What volume of ammonia at STP can be obtained when steam is passed over 4000 g of calcium cyanamide? The balanced reaction is



(Molecular weight of $\text{CaCN}_2 = 80$, MW of $\text{NH}_3 = 17$.)

Solution: From the stoichiometry of the equation 1 mole of CaCN_2 produces 2 moles of NH_3 . Thus, if one calculates the number of moles of CaCN_2 , then one knows the number of moles of NH_3 . To calculate the number of moles of CaCN_2 divide its weight by its molecular weight, thus, the number of moles of CaCN_2 equals

$$\frac{4000 \text{ g CaCN}_2}{80 \text{ g/mole}} = 50 \text{ moles.}$$

Therefore, the number of moles of NH_3 produced equals 100 moles.

At STP, one mole of any gas occupies 22.4 liters. Hence, the total volume of NH_3 produced at STP is

$$V_{\text{NH}_3} = (22.4 \text{ liter/mole})(100 \text{ mole NH}_3) = 2240 \text{ liter.}$$

Iron (III) oxide is reacted with carbon monoxide in a blast furnace to produce iron. The balanced reaction is:



What volume of CO at STP is required to completely use up 31.94 kg of iron oxide? (MW of $\text{Fe}_2\text{O}_3 = 159.7$, MW of CO = 28.)

Solution: From the stoichiometry of the above reaction, one sees that 1 mole of Fe_2O_3 requires 3 moles of CO to react. Thus, three times as many moles of CO will react with a given number of moles of Fe_2O_3 . Once the number of moles of CO is known, one can use the fact that 1 mole of any gas at STP has a volume of 22.4 l to determine the volume of CO required.

The number of moles of Fe_2O_3 available for the reaction is the weight of Fe_2O_3 in grams divided by its molecular weight:

$$\text{moles Fe}_2\text{O}_3 = \frac{\text{weight Fe}_2\text{O}_3}{\text{MW Fe}_2\text{O}_3} = \frac{31940 \text{ g}}{159.7 \text{ g/mole}} = 200 \text{ moles.}$$

The number of moles of CO needed to completely react is $3(200 \text{ moles}) = 600 \text{ moles}$.

The volume at STP of 600 moles of CO is the number of moles of CO times the volume of 1 mole:

$$\begin{aligned}\text{volume CO at STP} &= (\text{moles CO}) (22.4 \text{ l/mole}) \\ &= (600 \text{ moles}) (22.4 \text{ l/mole}) = 13,440 \text{ l}.\end{aligned}$$

• PROBLEM 211

How many liters of phosphine (PH_3) gas at STP could be made from 30 g of calcium by use of the following sequence of reactions:



(Molecular weights: $\text{Ca} = 40$, $\text{PH}_3 = 34$.)

Solution: From the stoichiometry of these two equations 3 moles of Ca will yield 2 moles of PH_3 gas. Thus, if one knows the number of moles of Ca, then one can determine the number of moles of PH_3 produced. The number of moles of Ca given is its weight divided by its molecular weight. Therefore,

$$\text{number of moles Ca} = \frac{30 \text{ g Ca}}{40 \text{ g/mole}} = 0.75 \text{ moles}$$

From 0.75 moles of Ca, one produces $2/3$ as much or $2/3(0.75) = 0.50$ moles of PH_3 gas. At STP, one mole of any gas occupies 22.4 liters, hence, the total volume of PH_3 produced at STP

$$= (22.4 \text{ l/mole}) (0.50 \text{ moles } \text{PH}_3) = 11.2 \text{ l}.$$

• PROBLEM 212

Nitroglycerin ($\text{C}_3\text{H}_5(\text{NO}_3)_3$) explodes according to the following reaction:




producing only gaseous products. What is the total volume of gaseous products produced at standard temperature and pressure (STP) when 454 g of nitroglycerin explodes? The molecular weight of nitroglycerin is 227 g/mole.

Solution: This problem is an application of the ideal

gas equation, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = gas constant, and T = absolute temperature. Solving for V ,

$$V = \frac{nRT}{P}$$

STP is, by definition, 0°C ($= 273^\circ\text{K}$) and 1 atm pressure. Hence, $T = 273^\circ\text{K}$ and $P = 1$ atm. Also, $R = 0.082$ liter-atm/mole-deg. We must find the number of moles, n , of gaseous products. The number of moles of nitroglycerin we started with is equal to its mass divided by the molecular weight, or $454 \text{ g} / 227 \text{ g/mole} = 2$ moles of nitroglycerin. Dividing the equation for the reaction of nitroglycerin by 2 (so that $4\text{C}_3\text{H}_5(\text{NO}_3)_3$ becomes $2\text{C}_3\text{H}_5(\text{NO}_3)_3$), we obtain:



Thus, our 2 moles of nitroglycerin will produce a total of $6 + 3 + \frac{1}{2} + 5 = 14.5$ moles of gaseous products, so that $n = 14.5$ moles. Substituting the values of n , R , T , and P into the equation for V gives:

$$V = \frac{nRT}{P} = \frac{14.5 \text{ moles} \times 0.082 \text{ liter-atm/mole-deg} \times 273^\circ\text{K}}{1 \text{ atm}} \\ = 325 \text{ liters.}$$

• PROBLEM 213

A chemist performs the following reaction:



He collects the O_2 gas by water displacement at 20°C . He observes a pressure of 753 mm Hg. Assuming the pressure of water vapor is 17.5 mm Hg at 20°C and he started with 1.28 g of potassium chlorate (KClO_3), what volume of gas is produced? ($R = .0821 \text{ l-atm/mole } ^\circ\text{K}$.)

Solution: You are asked to find the volume of gas produced and can do so using the equation of state, $PV = nRT$, where P = pressure, V = volume, n = number of moles, R = universal gas constant and T = temperature in Kelvin (Celsius plus 273°).

In the problem you are told the pressure at which the gas is collected, which must be modified due to the presence of water vapor and the temperature. If you knew n , you could substitute for the values in the equation of state and solve for V .

Your procedure will be to find the number of moles of gas produced. This can be determined from the reaction e-

quation given and the use of stoichiometry. If you knew how many moles of KClO_3 you started with, you would know the number of moles of O_2 produced, since 3 moles of O_2 are generated for every 2 moles of KClO_3 . You can determine how many moles of KClO_3 you started with. You are told that the chemist has 1.28 g. Since the molecular weight of KClO_3 is 122.55 g, you have

$\frac{1.28}{122.55} = 1.045 \times 10^{-2}$ moles of KClO_3 . Therefore, from reaction there are $\frac{3}{2} (1.045 \times 10^{-2}) = 1.57 \times 10^{-2}$ moles of O_2 gas produced.

Now you go back to $PV = nRT$. Recall that the gas was collected by water displacement. Thus, it is saturated by water vapor. Hence, the pressure of O_2 is only 753 mm Hg - 17.5 mm Hg, or 735.5 mm where 17.5 mm Hg is the water vapor pressure. Since 1 atm = 760 mm,

$$(735.5 \text{ mm}) \left(\frac{1 \text{ atm}}{760 \text{ mm}} \right) = .968 \text{ atm.}$$

Now, substitute and solve for the volume V of O_2 produced. Rewriting and substituting,

$$V = \frac{nRT}{P} = \frac{(.0157)(.0821)(293^\circ\text{K})}{.968} = .390 \text{ liters or } 390 \text{ ml.}$$

CHAPTER 6

SOLIDS

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 191 to 211 for step-by-step solutions to problems.

Many solids form crystals or molecular structures in which the constituent atoms or molecules are in a repeating three-dimensional array. Metals, simple salts, and semiconducting solids like silicon are all crystalline. The most common non-crystalline or amorphous solid is glass. In this solid, the molecules are in a random orientation with respect to each other. Glasses do not have a sharp melting point, rather they have a melting range, thus leading some to classify glass as a supercooled or extremely viscous liquid.

A complete description of the possible crystalline structures of solids is beyond the scope of this text, but several common cubic crystalline structures, simple cubic, face centered cubic (fcc), and body centered cubic (bcc), are illustrated in the figures associated with Problems 218 and 219. A unit cell is the smallest repeating unit of the lattice.

A key solution in calculating the stoichiometric properties of unit cells is to ascertain how many cells share a particular atom or molecule. For cubic lattices, a corner species is shared by eight cells, an edge species by four cells, a face centered species by two cells, and a body centered species by only one cell.

In crystals, the atoms or molecules remain in a fixed orientation with respect to each other. The distances between atoms can be determined from x-ray diffraction patterns because the distances are comparable in magnitude to the wave lengths of x-rays. Hence x-ray diffraction patterns are determined by the regular spacings between the atoms or molecules.

The principles of stoichiometry discussed in Chapters 3 and 4 can be used, along with information on the size of unit cells, to calculate the densities, volumes, and many other properties of crystalline solids. For example, the theoretical density of a crystalline solid can be obtained by:

$$\rho = \frac{\text{mass of unit cell}}{\text{volume of unit cell}} \quad 5-3$$

This is the maximum value of a pure crystal; because of imperfections in the crystalline lattice, the measure density is usually slightly less than that calculated by this procedure.

Phase Diagrams

Phase diagrams are also discussed in this chapter. Phase diagrams are simply graphs, usually on pressure and temperature coordinates, showing the boundaries between phases. An example for a simple single-component phase diagram (e.g., water) is shown below and illustrates some very familiar as well as some lesser known properties of pure compounds.




Figure 1 — Phase diagram for single component system

The scales on the axes of the sample phase diagram shown above have deliberately been left off so that the diagram applies to any single-component

system which has only one solid phase*. If the diagram above were to apply for water, then the normal boiling point, T_b (the temperature where the liquid and gas phases are in equilibrium at 1 atm. total pressure), would be 100°C. The melting point, T_m , is 0°C, and at a pressure of one atmosphere, water is a liquid between 0°C and 100°C. At temperatures and pressures represented by points along the lines separating the phases, both phases can exist in equilibrium with each other.

The line BN represents the temperatures and pressures where the solid and gas phases are in equilibrium. Similarly, the line NA represents the points where the liquid and solid phases are in equilibrium; for water and many other species, it slants slightly to the left of vertical as shown in this figure illustrating the effect of pressure on lowering the melting point. Note that in a similar figure in Problem 231, the solid-liquid equilibrium boundary between the solid and liquid phases has a positive slope; either can occur. The line NC represents the points where the liquid and gas phases are in equilibrium and this line ends at point C , the critical point. Above the critical point (218 atm and 375°C for water), the liquid and gas phases are indistinguishable. In this part of the phase diagram the material is sometimes called a supercritical fluid since it is not quite correct to refer to it as either a liquid or gas. There is normally no corresponding end point at A where the solid and liquid phases become indistinguishable (except, perhaps, at extreme pressures).

Note that at point N (0.0098°C and 4.58 torr for water) the solid, liquid, and gas phases meet. At this point, called the triple point, all three phases can exist in equilibrium. The triple point is unique; that is, there is only one point where this can occur, and both the temperature and pressure must be exactly equal to those represented by the coordinates of the triple point. On the other hand, there are many values of temperature and pressure where both liquid and gas or liquid and solid or even gas and solid phases exist in equilibrium.

If the temperature is raised at constant pressure along the line rs , the material will change from a liquid to a gas. The same thing will happen if the pressure is lowered at constant temperature along the line mn . Similarly, the solid phase will sublime (change to a gas) if the appropriate state variable (temperature/pressure) changes along the lines pq or ut .


*Many single-component systems have more than one solid phase. These are typically different crystalline forms of the component. Sulfur, for example, has a monoclinic and orthorhombic solid crystalline phase. In a phase diagram for such a material, areas of the diagram exist which represent the temperatures and pressures where each of the solid phases is stable.

Step-by-Step Solutions to Problems in this Chapter, “Solids”

CRYSTAL STRUCTURE

• PROBLEM 214

Distinguish between crystalline and amorphous solid substances, using some specific examples. To what extent is the distinction useful?




Solution: Crystalline substances can be generally thought of as species composed of structural units with specific geometric patterns. The accompanying drawing of sodium chloride would be an example of such a pattern. The important point is that there exists a regularity in the arrangement of structural units. Structures with regularity generally show a sharp and characteristic melting point, which is the case with crystalline substances.

Amorphous substances, however, tend to be shapeless and without definite order. That is, you have a randomness. For example, glassy or glasslike materials such as Plexiglas and silicate glasses. In substances with a general lack of order, the melting points vary over a range or temperature interval. For amorphous substances, this is

exactly what you find. It would, however, be incorrect to state categorically that amorphous substances are without ANY order. For they do tend to have short range order even though they do contain long-range randomness.

• PROBLEM 215


Explain briefly the following terms; (a) isotropic and anisotropic solids, (b) general and directional properties of crystalline substances, (c) a plane, an axis, and a center of symmetry and (d) polymorphism and allotropism.



Plane of symmetry
(abcd)



Axis of symmetry
(ab)



Point of symmetry
(c)

Solution: These properties all pertain to the solid state.

(a) An isotropic solid is one in which there are no directional differences in structure. Isotropic solids usually do not form regular crystals, and examples of an isotropic amorphous solids are Plexiglass and glass. An anisotropic solid is one in which its properties depend, in general, upon the particular direction along which the measurement is made. Crystalline substances, for example, are anisotropic. Examples of anisotropic crystals are cubic and tetrahedral lattices. (b) By general properties of crystalline substances, you mean those such as density, specific heat, melting point and chemical composition. These are called "general" due to the fact that none of them involve any special direction. Directional properties refer to those such as refraction and absorption of light, cohesion, expansion and electrical and thermal conductivity, all of which are measured in a special direction. (c) Plane of symmetry is said to exist in a crystal if it can be divided by an imaginary plane passing through the center, such that the two equal parts formed are an exact mirror image of each other. An axis of symmetry is said to exist in a crystal if you can draw a line through the crystal's center and when you proceed to rotate the crystal about this line through 360° the same appearance of itself is produced more than once. A crystal will have a center of symmetry, if every crystal face has a twin equidistant on the opposite side of this center. The accompanying figures illustrate each of these. (d) Polymorphism refers to the

existence of substance having a definite composition yet occurring in more than one crystalline form. When the substance is an element rather a compound, this phenomenon is termed allotropy.

• PROBLEM 216

It is known that both metallic and ionic crystals are good conductors of electricity, yet many of their other properties differ significantly. Explain why.

Solution: The similarities and differences of the two will be seen after an investigation of their space lattices, i.e., those patterns of points describing the atomic or molecular arrangements of points in a unit of a crystal. For something to conduct electricity, it must have charges. In an ionic crystal, the lattice points are occupied by positive or negatively charged ions, each of which occupies a position exactly equivalent to every other species of like charge. The electric conductivity of ionic crystals is low but increases with Temp. because more electrons are excited into the conduction band of the crystal. Ionic crystals are held together by electrostatic forces between the ions.

In a metallic crystal, you have discrete atoms, not ions, at the lattice points. It seems that the species should not conduct electricity. It is important to realize, however, the valence or outer electrons of the metallic crystal are distributed over the crystal as a whole rather than being localized on each atom. The mobile electrons, therefore, which are able to move in an applied electric field, account for the electrical conductivity.

This mobility of electrons also explains why a metallic crystal is generally strong, lustrous and malleable. In the ionic crystal, you have strong electrostatic forces between the ions, which accounts for the high melting points. However, these crystals also tend to be hard and brittle because the crystal consists of parallel sheets of positively and negatively charged ions, lateral displacement may bring ions of like charge into the vicinity of each other, resulting in electrostatic repulsion of like charges. This, then, helps to explain why they possess a facile crystal fracture.

• PROBLEM 217

The properties of diamond and graphite differ vastly. Explain these differences on the basis of their fundamentally different space lattices.



**Diamond
Structure**



**Graphite
Structure**

Solution: The arrangement (location) of all atoms in a unit of a crystal is termed a space lattice, i.e., those patterns of atoms describing the atomic or molecular arrangements. This problem calls for a comparison of such lattices in a diamond and graphite. The accompanying figures show the space lattices in question.

The diamond lattice surrounds each carbon atom by 4 others in a tetrahedral configuration. There exist covalent bonds between the carbon atoms, which means they share electrons. These covalent forces are responsible for the hard nature and high melting point of the substance. On the other hand, graphite has covalent bonding in two-dimensional, planar, hexagonal rings. As the figure suggests, the sheets of these atoms are held together by weak forces of attraction. This means, that the layers can slide easily over each other. This, then, accounts for the lubricating properties and flakiness associated with this material.

LATTICE STRUCTURES AND UNIT CELLS

• PROBLEM 218

Excluding hexagonal unit cells, when counting the number of points inside a cell, a point on an edge is $\frac{1}{4}$ inside the cell, and a lattice point at a corner is $\frac{1}{8}$ inside the cell. Justify these fractions. Calculate, also, the net number of lattice points in the following unit cells: simple cubic, body-centered cubic, face-centered cubic, and tetragonal.


Solution: To solve this problem properly you need to know the definition of a unit cell and lattice point. You need, also, to know the actual structures of the unit cells given. You proceed as follows: A unit cell is that small fraction of a space lattice, which sets the pattern for the whole lattice. In other words, it is the smallest portion of the space lattice (which is just




simple cubic
(a)



body-centered cubic
(b)



face-centered cubic
(c)



(d)



tetragonal
(e)

the pattern of points which describes the arrangement of atoms or molecules in the crystal), which, when moved repeatedly a distance equal to its own dimensions along the various directions, generates the whole space lattice equivalent to the original lattice. The lattice is, of course, just points that denote atoms or molecules; thus, the term lattice point. The accompanying illustrations give you the structures in question. From this, you can obtain justification for the fractions and find the net number of lattice points. For example, take the simple cubic structure. In a space lattice, each atom is shared with three other unit structures as seen in figure d.

Note; This point is shared by 8 unit cells, if you place another 4 cubes on the side. Thus, each point is only $1/8$ in the cell. This same sort of procedure and reasoning can be used to justify all fractions in all unit cells given. Calculations now become easy. Simple cubic: Have 8 lattice points. Each is only $1/8$ inside cell. Thus, you have net of $(1/8)(8) = 1$.


Face-centered cubic: 8 lattice points at corners, each only $1/8$ inside cell. Have a point on each face. Each point is only $1/2$ inside cell. Have six points. Thus, net = $(8)(1/8) + (6)(1/2) = 4$.

Body-centered: Again have $(8)(1/8)$ from lattice points at corners. 1 point in center, which is entirely in cell. Thus net becomes $(8)(1/8) + (1)(1) = 2$.

Tetragonal: Only have a net of $(8)(1/8)$ from lattice points at corners.

• PROBLEM 219

Refer to the accompanying figure and determine the number of unit particles (atoms) in a unit cell of each of the three type lattices.



Unit cells in cubic lattices: (a) simple cubic lattice;
(b) body-centered lattice; (c) face-centered lattice.

Solution: The following rules may be used to determine the number of unit particles (atoms) associated with each type of cubic unit cell:

1) An atom at a corner contributes $1/8$ of its volume to each of eight adjacent cubes.

2) An atom on an edge of a cube contributes $1/4$ of its volume to each of 4 adjacent cubes.

3) An atom in the face of a cube contributes $1/2$ of its volume to each of 2 adjacent cubes.

4) An atom completely within a cube contributes all of its volume to the unit cell.

The number of unit particles in each of the three types of lattices can be found as follows:

Simple cubic: In the simple cubic lattice all eight units are of the same variety - they are all at corners, thus rule 1 is used.

$$\text{Total atoms in cell} = 8 \text{ corners} \times 1/8 \text{ atom/corner} = 1 \text{ atom}$$

Therefore the area of unit cube is equivalent to 1 atom.

Body-centered: In the body-centered lattice there are nine atoms contributing to the unit cube. There are 8 corners and one atom completely within the cube, thus, rules 1 and 4 are used.


$$\text{Total atoms in cell} = 8 \text{ corners} \times 1/8 \text{ atom/corner}$$

$$+ 1 \text{ atom in center} = 2 \text{ atoms.}$$

Face-centered: In the face-centered lattice there are 14 atoms contributing to the unit cube. There are 8 corners and 6 faces, thus, rules 1 and 3 will be used.

$$\begin{aligned} \text{Total atoms in a unit cell} &= 8 \text{ corners} \times 1/8 \text{ atom/corner} \\ &\quad + 6 \text{ faces} \times 1/2 \text{ atom/corner} \\ &= 4 \text{ atoms.} \end{aligned}$$

What fraction of the total space in a body-centered cubic unit cell is unoccupied? Assume that the central atom touches each of the eight corner atoms of the cube.



body-centered lattice

Figure 1





Figure 2



Solution: To determine the percent of the unit cube that is unoccupied by the atoms making up the lattice, one must subtract the volume taken up by the atoms from the volume of the cube. This volume is then divided by the volume of the cube and multiplied by 100 to find the percent of unoccupied space.

1) Determining the volume of the atoms: One must first determine the number of atoms contributing to the unit cube. In this lattice there are 8 corner atoms and 1 atom in the center. Corner atoms contribute $1/8$ of their volume to the cube. The atom in the center contributes its entire volume.

$$\begin{aligned} \text{no. of atoms in unit cube} &= (1/8 \text{ atom/corner} \times 8 \text{ corners}) \\ &+ 1 \text{ atom (in center)} = 2 \text{ atoms.} \end{aligned}$$

Thus, the volume in the unit cube taken up by the atoms is equal to the volume of two atoms. The radius of these atoms is taken to be 1. The volume of a sphere is $4/3 \pi r^3$.

$$\text{volume of 2 atoms} = 2 \times 4/3 \pi (1)^3 = 8.38$$

2) Volume of the cube: The corner atoms are assumed to be touching the central atom. The diagonal of the cube can be visualized as shown in Figure 2.

Because \overline{AB} is shown to be $4r$ or 4 the side of the cube \overline{AC} can be found using the Pythagorean Theorem. Once \overline{AC} is known the volume of the cube can be found. From geometry, it is known that

$$\overline{CB} = \overline{AC} \times \sqrt{2}$$

$$\overline{AB}^2 = \overline{AC}^2 + \overline{CB}^2$$

$$\overline{AB} = 4$$

$$4^2 = \overline{AC}^2 + 2(\overline{AC})^2$$

$$CB = \overline{AC} \times \sqrt{2}$$

$$4^2 = 3\overline{AC}^2$$

$$\frac{4}{\sqrt{3}} = \overline{AC}$$

The volume of the cube is equal to the length of the side cubed.

$$\text{volume of cube} = \left(\frac{4}{\sqrt{3}} \right)^3 = 12.32$$

3) The space in the cube is equal to the volume of the spheres subtracted from the volume of the cube.

$$\text{vol. of space} = \text{vol. of cube} - \text{vol. of spheres}$$

$$\text{vol. of space} = 12.32 - 8.38 = 3.94$$

$$\text{percentage of cube taken up by space} = \frac{3.94}{12.32} \times 100 = 32 \%$$

• PROBLEM 221

The spherical atomic nucleus of potassium has a radius of about 4×10^{-13} cm.; 6.02×10^{23} of these nuclei weigh about 39 g. Calculate the approximate density of the nucleus of potassium in tons per cubic centimeter. Volume of a sphere = $\frac{4}{3} \pi r^3$, where r is the radius.

Solution: The density is defined as the mass of a substance divided by its volume. To calculate the density one must know the weight and volume of the substance. Here one is given the weight in grams but the problem asks for the density in tons, thus one must convert grams to tons. The volume of the nuclei is not given, but can be found by using the radius of a nucleus, the formula for the volume of a sphere and the number of nuclei present.

1) Converting grams to tons

There are 454 g in one pound and 2000 lb in one ton, therefore, grams can be converted to tons by multiplying the number of grams by the following conversion factor $1 \text{ lb}/454 \text{ g} \times 1 \text{ ton}/2000 \text{ lb}$

$$\begin{aligned} \text{number of tons} &= 39 \text{ g} \times 1 \text{ lb}/454 \text{ g} \times 1 \text{ ton}/2000 \text{ lb} \\ &= 4.3 \times 10^{-5} \text{ tons.} \end{aligned}$$

2) Calculating volume of nuclei

The volume of a sphere is equal to $\frac{4}{3} \pi r^3$ where r is the radius of the sphere. One is given that the radius of a potassium nuclei is 4×10^{-13} cm.

$$\begin{aligned}\text{volume of 1 nuclei} &= \frac{4}{3} \pi (4 \times 10^{-13} \text{ cm})^3 \\ &= \frac{4}{3} (3.14) (64 \times 10^{-39} \text{ cm}^3) \\ &= 2.6 \times 10^{-37} \text{ cm}^3\end{aligned}$$

The total volume of the nuclei is equal to the number of nuclei times the volume of one nucleus

$$\text{total volume} = \text{no. of nuclei} \times \text{volume of one nucleus}$$

$$\begin{aligned}\text{total volume of potassium nuclei} &= 6.02 \times 10^{23} \times 2.6 \times 10^{-37} \text{ cm}^3 \\ &= 1.6 \times 10^{-13} \text{ cm}^3\end{aligned}$$

3) The weight in tons and the volume in cubic centimeters is now known. The density in tons per cm^3 can be calculated

$$\text{density} = \frac{\text{no. of tons (= mass)}}{\text{no. of cm}^3 \text{ (= volume)}}$$

$$\text{density} = \frac{4.3 \times 10^{-5} \text{ tons}}{1.6 \times 10^{-13} \text{ cm}^3} = 2.7 \times 10^8 \text{ tons/cm}^3.$$

• PROBLEM 222

The density of KF is 2.48 g/cm^3 . The solid is made up of a cubic array of alternate K^+ and F^- ions at a spacing of 2.665×10^{-8} cm. between centers. From these data, calculate the apparent value of the Avogadro number.

Solution: Avogadro's number is the number of particles in one mole of a substance. Here, one is given the dimensions of a cubic lattice made of $\frac{1}{2} \text{ K}^+$ ions and $\frac{1}{2} \text{ F}^-$ ions. Due to the fact that KF crystallizes like NaCl, it can be seen that there are eight ions or 4 formula units of KF in the cube. To calculate Avogadro's number from the data, one should find:

- (1) the volume of one cube
- 2) the weight of one cube using the density
- 3) the number of cubes in one mole of KF
- 4) Avogadro's Number by multiplying the number of cubes by 4, the number of formula units (KF) per cube.

Solving for Avogadro's Number:

1) The volume of the cube is found by cubing the length of the edge. From the crystallization of KF, it is seen that the length of the edge of cube is twice the spacing between ion centers. This is due to the fact that three ions make up an edge, and, as such, 2 spacings between the ions exist.

Therefore,

$$\text{length of edge} = 2.665 \times 10^{-8} \text{ cm} \times 2 = 5.33 \times 10^{-8} \text{ cm}$$

$$\text{volume of cube} = (5.33 \times 10^{-8})^3 = 1.51 \times 10^{-22} \text{ cm}^3.$$

2) Using the density, one knows that 1 cm³ weighs 2.48 g. Therefore, the weight of one cube is equal to the density times the volume of a cube.

$$\begin{aligned} \text{weight of one cube} &= 2.48 \text{ g/cm}^3 \times 1.51 \times 10^{-22} \text{ cm}^3 \\ &= 3.76 \times 10^{-22} \text{ g/cube} \end{aligned}$$

3) One mole of KF weighs the sum of the weights of one mole of K⁺ and one mole of F⁻. (MW of K = 39.10, MW of F = 19.00.)

$$\text{MW of KF} = 39.10 + 19.00 = 58.10 \text{ g/mole}$$

The number of cubes in one mole is then found by dividing 58.10 g/mole by the weight of one cube.

$$\begin{aligned} \text{no. of cubes in one mole} &= \frac{58.10 \text{ g/mole}}{3.76 \times 10^{-22} \text{ g/cube}} \\ &= 1.55 \times 10^{23} \text{ cubes/mole.} \end{aligned}$$

4) One finds the number of KF particles in one mole (Avogadro's Number) by multiplying 1.55 × 10²³ cubes/mole by 4 formula units/cube, because there are 4 KF formula units in each cube.



$$\begin{aligned} \text{Avogadro's Number} &= 4 \frac{\text{formula units}}{\text{cube}} \times 1.55 \times 10^{23} \frac{\text{cubes}}{\text{mole}} \\ &= 6.2 \times 10^{23} \text{ formula units/cube.} \end{aligned}$$

● PROBLEM 223

Iron may crystallize in the face-centered cubic system. If the radius of an Fe atom is 1.26 Å, (a) Determine the length of the unit cell. (b) Calculate the density of Fe if its atomic weight is 55.85.

Solution: Picture the face of the cube with an iron

face-centered lattice



atom at each corner and one in the center of each face as shown in the figure.

The diagonal BC will be equal to the length L 4 radii of Fe atoms. Hence,

$$BC = 4 \times 1.26 \text{ \AA} = 5.04 \text{ \AA}$$

Using the Pythagorean Theorem, one knows that the squares of the two sides of a right triangle added together equal the square of the hypotenuse. Thus,

$$\overline{AB}^2 + \overline{AC}^2 = \overline{BC}^2$$

Because the sides of a square are equal - $\overline{AB} = \overline{AC}$. From this it can be seen that $2 \overline{AB}^2 = \overline{BC}^2$. \overline{AB} is defined as the length of the unit cell. Using

$$2 \overline{AB}^2 = \overline{BC}^2,$$

it can be seen that

$$\overline{AB}^2 = \frac{\overline{BC}^2}{2} \quad \text{and} \quad AB = \sqrt{\frac{\overline{BC}^2}{2}} = \frac{BC}{\sqrt{2}}$$

Since we have already found BC, this equation can be solved. ($BC = 5.04 \text{ \AA}$.)

$$AB = \frac{\overline{BC}}{\sqrt{2}} = \frac{5.04 \text{ \AA}}{1.42} = 3.55 \text{ \AA}$$

The length of the unit cell is 3.55 \AA .

(b) The density is defined as the mass in grams of a unit cell in one unit volume. In this problem we are given the mass of one mole of Fe and we have calculated the length of one side of the unit cube. To find the density one must calculate the number of atoms in one unit cell, the weight of one atom of Fe and the volume of one unit cube first.

The following rules may be used to determine the number of unit particles (atoms) associated with the

face-centered lattice:

1) An atom at a corner contributes $1/8$ of its volume to each of 8 adjacent cubes.

2) An atom in the face of a cube contributes $1/2$ of its volume to each of 2 adjacent cubes.

In this lattice we have 8 corner atoms and 6 face atoms. To calculate the total number of atoms contributing to the unit cube one can use the above rules.

$$\begin{aligned}\text{Total number of atoms} &= 8 \text{ corners} \times 1/8 \text{ atom/corner} \\ &\quad + 6 \text{ faces} \times 1/2 \text{ atom/face} \\ &= 4 \text{ atoms.}\end{aligned}$$

The unit cube therefore contains the equivalent of 4 atoms.

To find the weight of 1 atom one divides the molecular weight by Avogadro's Number (6.02×10^{23}), because there are Avogadro's Number of atoms contained in a mole of atoms.

$$\text{weight of 1 atom} = \frac{55.85 \text{ g}}{6.02 \times 10^{23}} = 9.28 \times 10^{-23} \text{ g}$$

Since there are 4 atoms in this unit cube this figure is multiplied by 4.

$$\text{weight of 4 atoms} = \frac{4 \times (55.85 \text{ g})}{6.02 \times 10^{23}} = \text{weight of unit cube}$$

The volume of the cube is found by cubing the length of a side of the square because the volume is equal to the length \times width \times height of a rectangular solid. In a cube all three of these quantities are equal.


$$\text{The volume of the cube} = (3.55 \times 10^{-8} \text{ cm})^3.$$

($1 \text{ \AA} = 10^{-8} \text{ cm}$). We now have the quantities necessary to find the density. cc = cubic centimeters = cm^3 .

$$\begin{aligned}\text{density} &= \frac{\text{weight of unit cube}}{\text{volume of unit cube in cc}} \\ &= \frac{4 \times (55.85 \text{ g})}{6.02 \times 10^{23}} \\ &\quad (3.55 \times 10^{-8} \text{ cm})^3 \\ &= 8.30 \text{ g/cc}\end{aligned}$$

The density of Fe is therefore 8.30 grams per cubic centimeter.

Metallic gold crystallizes in the face-centered cubic lattice. The length of the cubic unit cell, a , is 4.070 Å. (a) What is the closest distance between gold atoms? (b) What is the density of gold?



face-centered cubic

Solution: An ordered array of atoms, ions, or molecules is called a lattice. Every lattice is a three-dimensional stacking of identical building blocks called unit cells. The most symmetrical crystals have cubic lattices. There are three kinds of elementary cubic lattices: the simple cubic, the face-centered cubic, and the body-centered cubic. The length of the cube edge is designated by the symbol a . In a face-centered lattice, an atom would be located at each corner of the unit cell and at the center of each of the six faces.

(a) In the gold crystal above, the closest distance from one corner atom to another corner atom is a , 4.070 Å. The distance from any corner atom to an atom at the center of a face is one-half the diagonal of that face. Using the fact that the diagonal of any square is $\sqrt{2} \times$ length of one side, the diagonal is then $a\sqrt{2}$. Half the diagonal is $\frac{1}{2} a\sqrt{2}$ or

$$\frac{1}{2} (4.070 \text{ Å}) \sqrt{2} = 2.878 \text{ Å}$$

This distance is the closest distance between atoms, since this distance is shorter than the distance between adjacent corners.

(b) The density is computed by first counting the number of gold atoms that can occupy one unit cell. This number is $1/8$ times the number of occupied corners in the unit cell plus $\frac{1}{2}$ times the number of occupied face-centers. Since a cube has 8 corners and 6 faces, the number is

$$1/8 \cdot 8 + \frac{1}{2} \cdot 6 = 4$$

The volume of the cubic unit cell is a^3 . The

density if the mass of 4 gold atoms (4 atomic Wt./Avogadro's #) divided by the volume of the unit cell, a^3 . Thus, the formula for the density of a gold crystal lattice is


$$\text{Density} = \frac{(4 \text{ atoms})(\text{Atomic Wt. of gold})}{a^3 (\text{Avogadro's number})}$$

Substituting into the equation the Atomic Weight of gold, 197.0 g/mole, Avogadro's number, 6.023×10^{23} atoms/mole, and converting 4.070 Å to 4.070×10^{-8} cm, then

$$\begin{aligned} \text{Density} &= \frac{(4 \text{ atoms}) (197.0 \text{ g/mole})}{(4.070 \times 10^{-8} \text{ cm})^3 (6.023 \times 10^{23} \text{ atoms/mole})} \\ &= 19.4 \text{ g/cm}^3. \end{aligned}$$

• PROBLEM 225

Krypton crystallizes in the face-centered cubic system, with the edge of the unit cell 5.59 Å. Calculate the density of solid krypton. Assume 1 Å = 10^{-8} cm.



face-centered lattice

Solution: To determine the density of krypton in g/cc, one can use the given information in the following way:

- 1) determine the number of atoms in one cube
- 2) determine the number of cubes in one cubic centimeter
- 3) determine the weight of one cube
- 4) determine the weight of one cubic centimeter (i.e. the density).

Determining the density:

- 1) In a face centered cube, there are 14 atoms contributing to the volume. 8 are corner atoms and 6 are

face atoms. Corner atoms contribute 1/8 of their volume to the cube and face atoms contribute 1/2 of their volume. Thus, the number of atoms making up the cube can be found.

$$\begin{aligned} \text{no. of atoms} &= (1/8 \text{ atom/corner} \times 8 \text{ corners}) \\ &+ (1/2 \text{ atom/face} \times 6 \text{ faces}) = 4 \text{ atoms.} \end{aligned}$$

2) $1 \text{ \AA} = 10^{-8} \text{ cm}$. Thus, the side of the cube is equal to 5.59 \AA or $5.59 \times 10^{-8} \text{ cm}$. The volume of a cube is equal to the length of the side cubed.

$$\text{volume of one cube} = (5.59 \times 10^{-8} \text{ cm})^3 = 1.75 \times 10^{-22} \text{ cc}$$

The number of cubes in 1 cc is found by dividing 1 cc by the volume of one cube.

$$\text{no. of cubes in 1 cc} = \frac{1 \text{ cc}}{1.75 \times 10^{-22} \text{ cc}} = 5.71 \times 10^{21}$$

3) The weight of one mole of krypton is 84. 6.02×10^{23} atoms (Avogadro's number) of it must weigh 84 g. Therefore, the weight of one atom is found by dividing 84 g by 6.02×10^{23} atoms. There are 4 atoms in one cube, thus the weight of one cube is equal to the weight of one atom times 4.

$$\text{weight of 1 atom} = \frac{84 \text{ g}}{6.02 \times 10^{23} \text{ atoms}}$$

$$\begin{aligned} \text{weight of 4 atoms} &= 4 \text{ atoms} \times \frac{84 \text{ g}}{6.02 \times 10^{23} \text{ atoms}} \\ &= 5.58 \times 10^{-22} \text{ g.} \end{aligned}$$


4) The density is now found by multiplying the weight of 1 cube by the number of cubes in 1 cc.

$$\text{density} = \text{no. of cubes in 1 cc} \times \text{weight of 1 cube}$$

$$\text{density} = 5.71 \times 10^{21} / \text{cc} \times 5.58 \times 10^{-22} \text{ g} = 3.19 \text{ g/cc.}$$

• PROBLEM 226

It is known that AgCl has the same structure as NaCl. X-ray measurements of AgCl show a unit-cell edge length of .55491 nm. The density of the crystal is found to be 5.561 g/cm³. Assuming 10^{-7} cm/nm , find the percentage of the sites that would appear empty. Assume that lattice vacancies are the only defects.



The Structure of Silver Chloride

Solution: The key to solving this problem is to compute the theoretical density of the unit cell from its X-ray determined cubic edge length. Then, compare this value with the actual density to find the percent of the sites that would appear to be empty.

$$\text{Theoretical density} = \frac{\text{mass of 1 unit cell}}{\text{volume of 1 unit cell}}$$

Each unit cell possesses 4 ions of each Ag^+ and Cl^- . This is determined from the fact that each corner ion contributes $1/8$ of its volume to each of 8 adjacent unit cells, each face ion contributes $1/2$ of its volume to the two adjacent cells, each edge ion contributes $1/4$ of its volume to each of 4 adjacent cells and each enclosed ion contributes its entire volume. As seen in the accompanying figure, there are 8 corner ions, 12 edge ions, 6 face ions and 1 totally enclosed ion. The molecular weight of AgCl is 143.32. Thus, the mass of 1 mole of unit cells is 4 times 143.32 g/mole of unit cells or 573.28 g/mole of unit cells. Since the unit cell is a cube, volume = (cubic-edge length)³. Cubic edge length = .55491 nm or $.55491 \times 10^{-7}$ cm. Thus, volume becomes equal to $(.55491 \times 10^{-7})^3 = 1.708 \times 10^{-22} \text{ cm}^3$. As such, theoretical density =

$$\frac{573.28 \text{ g/mole of unit cells}}{\frac{6.022 \times 10^{23} \text{ unit cells/mole}}{1.708 \times 10^{-22} \text{ cm}^3}} = 5.571 \text{ g/cm}^3$$

The given density is 5.561 g/cm^3 . Thus, the actual figure is


$$\frac{5.571 - 5.561}{5.571} \times 100\% = .2\% \quad \text{too small. This is}$$

the percent of lattice that is vacant.

• PROBLEM 227

Find the ionic radius of I^- , given that the unit-cell edge length of LiI is measured to be .6240 nm. Assume

that the large, negative ions are in actual contact with the diagonal.



The Structure of Lithium Iodide

Solution: To solve this problem, determine the length of the face diagonal in LiI space lattice, which is the pattern of the arrangement of atoms or molecules in a crystal. Because LiI has the same form of the cubic unit cell as NaCl, its space lattice will possess 3 I^- ions on the face diagonal. Thus if one knows the length of the face diagonal, one can divide by 4 to deduce the radius of I^- . The face diagonal can be found (from the Pythagorean Theorem) by multiplying $(\sqrt{2})$ times (cube-edge length). Thus, the face diagonal = $(\sqrt{2})$ (.6240) = .8823 nm. However, face diagonal = 4 I^- radii.


Thus, I^- radius = $\frac{.8823}{4} = .2206$ nm.

• PROBLEM 228

Consider a unit cell of sodium chloride in the accompanying figure. (a) What fraction of a Cl^- at each corner of the cube is within the unit cell? (b) What fraction of a Cl^- appears in each face? (c) What fraction of each Na^+ is within the unit cell? (Note that one complete Na^+ is in the center.) (d) Add all the fractions for each ion within the unit cell. What is the ratio of Na^+ to Cl^- ?

Solution: The following rules may be used to determine the number of unit particles (atoms) associated with each type of cubic unit cell:

- 1) An atom at a corner contributes $1/8$ of its volume to each of 8 adjacent cubes.
- 2) An atom on an edge of a cube contributes $1/4$ of its volume to each of 4 adjacent cubes.
- 3) An atom in the face of a cube contributes $1/2$ of



Crystal lattice of sodium chloride

its volume to each of 2 adjacent cubes.

4) An atom completely within a cube contributes all of its volume to the unit cell.

(a) According to rule 1 an atom at a corner contributes $1/8$ of its volume to the unit cell.

(b) From rule 3 one can see that an atom on the face of the unit cell contributes $1/2$ of its volume to the cell.

(c) From rule 2 one can see that atoms on the edge of the unit cell contribute $1/4$ of their volume to the unit cell. Therefore the sodium ions on the edge of the cell contribute $1/4$ of their volume and the one in the center of the cell contributes its entire volume.

(d) There are 14 Cl^- ions contributing to the cell. 8 are at corners and 6 are on faces - using parts a + b of this problem one can calculate the number of atoms that the Cl^- ions contribute to the unit cell.

$$\begin{aligned} \text{total number of } \text{Cl}^- \text{ ions} &= 8 \text{ corners} \times 1/8 \text{ atom/corner} \\ &\quad + 6 \text{ faces} \times 1/2 \text{ atom/face} \\ &= 4 \text{ atoms of } \text{Cl}^- \end{aligned}$$


The Na^+ ions can be dealt with in a similar manner. There are 13 Na^+ ions, 12 on edges and 1 in the center - using part c of this problem one can find the number of Na^+ ions contributing to the unit cell.

$$\begin{aligned} \text{total number of } \text{Na}^+ \text{ ions} &= 12 \text{ edges} \times 1/4 \text{ atom/edge} \\ &\quad + 1 \text{ atom in center} \\ &= 4 \text{ atoms of } \text{Na}^+ \end{aligned}$$

There are 4 atoms of Cl^- and 4 atoms of Na^+ contributing to the unit cell. The ratio will therefore be - $\text{Na}^+ : \text{Cl}^-$ is 4 : 4 or 1 : 1.

Na^+ and Cl^- contribute equally to the unit cell.

Find the distance between a plane passing through the centers of one close-packed layer of spheres and another plane passing through the centers of an adjacent close-packed layer. You may assume that all spheres are identical with a radius of .200 nm.



Solution: The small fraction of a space lattice, which is the pattern of points that describes the arrangement of atoms or molecules in a crystal, is a unit cell. The unit cells contain only points that locate atomic or molecular centers. Since atoms are space-filling entities, their structures can be described as resulting from the packing together of representative spheres. The most efficient packing together of equal spheres is called closest packing. In this problem, you are asked for a distance between two planes, which is just the spacing between the layers of spheres. A face-centered cube, a type of unit cell, is a close packed structure. From the accompanying figure, you can see that the body diagonal of a face-centered cube is perpendicular to the close-packed stacking layers. Thus, if you stand it on a corner, the close packed layers will be parallel to the floor. To see this more clearly, you can use models. From this, you can relate the body diagonal to the spacing between layers. Namely, body diagonal = 3 times the

spacing. Thus, you need to find the length of the body diagonal.

$$\text{Body diagonal} = \sqrt{3} \text{ (edge length).}$$


$$\text{Edge length} = 1/\sqrt{2} \text{ (face diagonal).}$$

The face diagonal = 4 times the radii of the spheres, as seen from the space lattice. You are told the radius of a sphere is .200 nm. Thus, face diagonal = $4(.200 \text{ nm}) = .800 \text{ nm}$. Thus, edge length = $(1/\sqrt{2})(.800) = .5657 \text{ nm}$ and body diagonal = $(\sqrt{3})(.5657) = .9798 \text{ nm}$. Recall, body diagonal is 3 times the spacing, which means the distance between the planes is $\frac{.9798 \text{ nm}}{3} = .327 \text{ nm}$.

PHASE DIAGRAMS


• PROBLEM 230

The diagram below is an example of a phase diagram for a pure substance. To what phases do the regions A, B, and C correspond?



Solution: Following the 1 atm constant pressure line from left to right, we are proceeding from low values of the temperature to high values. Therefore, we will intersect the three regions in the order solid-liquid-vapor. The regions A, B, C hence correspond to the solid, liquid, and vapor phases, respectively. Point a denotes the normal freezing (melting) point of the substance and the point b denotes the normal boiling point.

Draw a labelled phase diagram for a substance Z which has the following properties; normal boiling point = 220°C , normal freezing point 80°C , and triple point 60°C and .20 atm. Predict the freezing and boiling, if the pressure were .80 atm?



Solution: To draw this diagram you want to understand all the terms involved. The relation between solid, liquid, and gaseous states as a function of the given temperature and pressure can be summarized on a graph known as a phase diagram. From the given experimental observations, you can draw the diagram. The lines which separate the states in a diagram represent an equilibrium between the phases. The intersection of the three lines is called the triple point, where all three phases are in equilibrium with each other. By normal boiling and melting points you mean those readings taken at 1 atm. Thus, the phase diagram can be written as shown in the accompanying figure.

From the diagram you see that if the atm was .80, the b.p. and f.p. would drop, respectively, to 215°C and 85°C .

CHAPTER 7

PROPERTIES OF LIQUIDS

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 213 to 250 for step-by-step solutions to problems.

Liquids differ from solids in that the molecular structure is random, and the liquid will normally assume the shape of the vessel or container in which it is placed. Liquids are characterized by the following properties:

Density: The mass per unit volume occupied by a liquid is an important property used to characterize a liquid. Liquids are nearly incompressible and require extreme pressures to make a significant difference in the density. Also, only small differences in the density occur with moderate changes in the temperature; hence the density of a liquid is frequently considered to be a constant, except when very precise values are required. Water has a density of 1 gram per cubic centimeter; common hydrocarbons, such as those in gasoline, have densities of approximately 0.7 g/cc and float on water, while chloroform (CHCl_3), a common organic solvent, has a density of 1.5 g/cc and sinks in water.

Viscosity: The viscosity is the internal resistance of a liquid to shear stresses. Although not discussed in the problems of this text, it is an important characteristic that reflects the ability of a liquid to flow. Low viscosity liquids like water and gasoline flow and are pumped easily. Higher viscosity liquids like molasses flow very slowly (hence the cliché “slower than cold molasses”). Supercooled liquids, like glass, are so viscous that they do not flow at all and are usually considered solids.

Vapor Pressure: Liquids are in equilibrium with the gas or vapor above the liquid. The pressure of the gas above a pure liquid is called the vapor pressure or sometimes, for emphasis, the pure component vapor pressure. The vapor pressure is a strong function of the temperature.

Surface Tension: Liquids are characterized by attractive forces between the molecules. The strength of these attractive forces is characterized by the surface tension. At the surface of a liquid, the attractive forces tend to pull the surface atoms or molecules back into the liquid. This force can be measured as a force per unit length (N/m) of the liquid film.

The density of irregularly shaped solids is frequently measured by submerging them in a liquid to determine their volume. The solid volume is equal to the volume of liquid displaced. Since the determination of the mass is usually straightforward (it can be weighed on a balance), the density of irregular solids can be easily calculated once the volume is known.

Freezing Point Depression and Boiling Point Elevation

The freezing point is lowered and the boiling point raised for liquids when a solute is dissolved. The freezing point depression and boiling point elevation are proportional to the number of moles dissolved per unit mass of solvent. The change in freezing or boiling point is calculated from the simple equation

$$\Delta T_f = k_f (m) \quad 7-1$$

$$\Delta T_b = k_b (m) \quad 7-2$$

where k_f and k_b are the freezing and boiling point constants respectively, and m is the molality (moles of solute/1000 g solvent) of the solution. The freezing and boiling point constants are determined experimentally and are useful because they are a characteristic of the solvent and usually not a characteristic of the solute. Freezing point depressions are frequently used to determine the molecular weights of solutes since the mass of solute and freezing point depression are easily measured.

Equations 7-1 and 7-2 can be written with the molecular weight shown explicitly as

$$\Delta T_f = k_f \frac{(\text{mass of solute}/1000 \text{ g solvent})}{\text{M.W.}} \quad 7-3$$

$$\Delta t_b = k_b \frac{(\text{mass of solute}/1000 \text{ g solvent})}{\text{M.W.}} \quad 7-4$$

The key to solving the problems involving freezing point depression and boiling point elevation is the use of these equations to solve for the unknown quantity — ΔT , k_f or k_b , or the molecular weight, (M.W.).

Vapor Pressure—Raoult's Law

The vapor pressure is a characteristic of a liquid and is a strong function of temperature. It is precisely defined as the equilibrium gas (or vapor) pressure above a pure liquid at a specified temperature with no other species present. Except at extreme pressures, it is an excellent approximation to assume that the vapor pressure is independent of the total pressure, i.e., unaffected by the presence of other species in the gas phase.

There are empirical equations that permit the vapor pressure to be calculated, but these equations contain approximations and, for precise work, measurements should be used.

Raoult's Law predicts the partial pressure above a liquid which contains a mixture of compounds. It is correct only for ideal liquid mixtures, i.e., liquid mixtures where the attractive forces are the same between like and unlike molecules. It is a good approximation for mixtures of isomers (such as normal pentane, isopentane, and neopentane) or adjacent members of homologous series (such as benzene, toluene, and xylene).

Raoult's Law is written as follows:

$$p_i = x_i P_i^\circ \quad 7-5$$

where p is the partial pressure in the gas phase, x is the mole fraction in the liquid phase, and P° is the pure component vapor pressure (or just vapor pressure). The subscript i indicates that this equation can be applied to each species in the solution, i.e., $i = 1, 2, 3, \dots$. If two of the terms in Equation 7-5 are known, the third can be easily calculated. If Raoult's Law holds for one species in a solution, it will hold for all the species. Departures from Raoult's Law are common in nature, but are not discussed in this text.

One key to working the problems involving Raoult's Law is to observe that the total pressure is the sum of the partial pressures. From the ideal gas law, the mole fractions in the vapor phase are simply the respective partial pressures divided by the total pressure (sum of the partial pressures). As well, the mole fractions of the various solutes must add up to 1.

$$Y_i = P_i/P \quad 7-6$$

where Y is the vapor phase mole fraction and P is the total pressure.

Clausius Clapeyron Equation of Vaporization

The Clausius Clapeyron equation is useful in calculating changes in vapor pressure with temperature. It contains several approximations and is derived by integrating the Clapeyron equation, which has a firm basis in theory.

$$\frac{dP^\circ}{dT} = \frac{\Delta H}{T\Delta V} \quad 7-7$$

where ΔH and ΔV are the enthalpy and volume changes associated with vaporization of the liquid. If it is assumed that ΔV is simply the vapor phase volume ($\Delta V = RT/P$), and that ΔH is a constant, Equation 7-7 can be integrated to obtain the the Clausius Clapeyron equation.

$$\ln (P^\circ_2/P^\circ_1) = -\Delta H^{vap}/R (1/T_2 - 1/T_1) \quad 7-8$$

The assumptions are valid over limited temperature ranges for many species.

Note: Several problems (see Problems 263, 267, 268) in this text show the Clausius Clapeyron equation written as

$$\log (P^\circ_2/P^\circ_1) = \Delta H/19.15 (1/T_1 - 1/T_2) \quad 7-9$$

This form of the equation may be confusing. The factor, 19.15, is the product of the gas constant, R , in joules/mole $^\circ\text{K}$ (8.31) and the conversion factor from natural to common logarithms (\log_e to \log_{10}), i.e., $8.31 \times 2.303 = 19.15$. To use this form of the equation, ΔH must be expressed in joules/mole and base

10 logarithms must be used.

It is possible, from Equation 7-7, to calculate changes in vapor pressure from the enthalpy of vaporization or to calculate the enthalpy (or heat) of vaporization from measurements of the vapor pressure at more than one temperature.

Osmotic pressure (π), for very dilute solutions, can be represented by the equation

$$\pi = nRT/V \quad 7-10$$

where n is the number of moles of solute in a volume, V , of solution. It is important to note that this equation does not apply except for very dilute solutions. With Equation 7-9 it is possible to determine the molecular weight of a species by measuring the osmotic pressure produced by a known mass of solute.

Surface tension is usually determined by measuring the force required to pull a ring of known diameter from a solution or by measuring the height that a liquid "climbs" the wall of a wetted vessel.

If a ring is pulled from a liquid, two liquid films (inside and outside) form between the ring and the liquid. The force required to pull the ring from solution divided by the length of surface ($2\pi D$) is the surface tension. When surface tension is measured by determining the height that the liquid climbs the wall of a wetted vessel, the same principle is used to determine the surface tension. The equation that relates surface tension to the height of the liquid is

$$\gamma = 1/2 r \cdot h \rho g \quad 7-11$$

where γ is the surface tension, ρ the density of the liquid, r the radius of the tube containing the liquid, h the height that the liquid film climbs above the free surface of the liquid, and g is the constant downward acceleration of gravity.

Step-by-Step Solutions to Problems in this Chapter, "Properties of Liquids"

DENSITY

• PROBLEM 232

What volume of a block of wood (density = 0.80 g/cm^3), which weighs 1.0 kg , will be above the water surface when the block is afloat? (Note: A floating object displaces its own weight of water.)

Solution: Since a floating object displaces its own weight in water, 1 kg of water is displaced by this block of wood. One can find the volume of the block of wood above the water by solving for the volume of the block and subtracting the volume of 1 kg of water from it. One uses the density to solve for the volume.

$$\text{density} = \frac{\text{weight}}{\text{volume}} = \frac{g}{\text{cm}^3}$$

Therefore:

$$\text{volume} = \frac{\text{weight}}{\text{density}}$$

Solving for the volume of the wood: $1 \text{ kg} = 1000 \text{ g}$.

$$\text{volume} = \frac{1000 \text{ g}}{.80 \text{ g/cm}^3} = 1.25 \times 10^3 \text{ cm}^3$$

Solving for the volume of the water displaced:

By definition the density of water is 1.0 g/cm^3 .

$$\text{volume} = \frac{1000 \text{ g}}{1.0 \text{ g/cm}^3} = 1.00 \times 10^3 \text{ cm}^3$$

volume of wood above water = volume of wood - volume of water

$$\begin{aligned}\text{volume of wood above water} &= 1250 \text{ cm}^3 - 1000 \text{ cm}^3 \\ &= 250 \text{ cm}^3\end{aligned}$$

• PROBLEM 233

A chemist dropped a 200 g object into a tank of water. It displaced 60 ml. of the water when it sunk to the bottom of the tank. In a similar experiment, it displaced only 50 g of an oil into which it was dropped. Calculate the density of the object and the oil.

Solution: The density (ρ) of a substance is defined as its mass divided by its volume.

$$\rho = \frac{\text{mass}}{\text{volume}}$$

Thus, to solve for the densities of the object and the oil, one must first calculate their respective masses and volumes. The mass of the object is 200 g, but the volume is not given. An object dropped in any liquid displaces a volume of liquid equal to the volume of the object. The object displaces 60 ml of water; therefore, the volume of the object is 60 ml. Solving for the density of the object:

$$\rho = \frac{\text{mass}}{\text{volume}} = \frac{200 \text{ g}}{60 \text{ ml}} = 3.33 \text{ g/ml.}$$

One is given that the object displaces 50 g of the oil, and from the water experiment, it is known that the volume of the object is 60 ml. Since the object displaces the same volume of liquid as it occupies, 60 ml of the oil weighs 50 g. Solving for the density of the oil:

$$\rho = \frac{\text{mass}}{\text{volume}} = \frac{50 \text{ g}}{60 \text{ ml}} = .833 \text{ g/ml.}$$

• PROBLEM 234

Assuming that the density of water is .9971 g/cm³ at 25°C and that of ice at 0° is .917 g/cm³, what percent of a water jug at 25°C should be left empty so that, if the water freezes, it will just fill the jug?

Solution: Density = $\frac{\text{mass}}{\text{volume}}$.

When water freezes, the mass is constant, but its density decreases. This means, therefore, that an increase in volume occurs. To solve this problem, you must

determine the change in volume of water from 25° to 0°. Because you are given the densities of the substance at each temperature, you can say for 1 mole of H₂O (MW = 18)

$$\frac{\text{mass}}{\text{volume}} = \frac{18 \text{ g/mole}}{y} = \frac{0.997 \text{ g}}{\text{cm}^3} \text{ for water at } 25^\circ\text{C and}$$

$$\frac{\text{mass}}{\text{volume}} = \frac{18 \text{ g/mole}}{x} = \frac{0.917 \text{ g}}{\text{cm}^3} \text{ for ice at } 0^\circ\text{C.}$$

Rewriting, you have .997 y = 18 and .917 x = 18. Both expressions equal 18, which means .997 y = .917 x, or

$$\frac{x}{y} = \frac{.997}{.917} = 1.087.$$

This means, then, that at 0°C the volume is 1.087 times greater than the original at 25°C. If you let B equal the fraction of water to be put in the jug, then you can say 1.087 B = 1, or B = .920. In other words, only 92% of the jug should be filled at 25° to obtain a completely filled jug at 0°. This means that 8 % is empty space.

• PROBLEM 235

The molecular diameter of an N₂ molecule, as deduced from the Van der Waals b parameter, is 3.15×10^{-8} cm. The density of liquid nitrogen is 0.8081 g/cm³. On a hard-sphere model, what fraction of the liquid volume appears to be empty space?

Solution: To find the fraction of the liquid volume that appears to be empty space, one must subtract the volume of spheres from the total volume of the liquid. One can find the volume of the spheres that constitute one mole of N₂ liquid by multiplying the volume of one sphere by 6.02×10^{23} , the number of spheres in one mole. The volume of a sphere is equal to $\frac{4}{3} \pi r^3$ where r is the radius of the sphere. The diameter of the N₂ sphere is given as 3.15×10^{-8} cm, thus the radius is half of length or 1.58×10^{-8} cm. Solving for the volume of 1 sphere:

$$\begin{aligned} \text{volume of 1 sphere} &= \frac{4}{3} \pi (1.58 \times 10^{-8} \text{ cm})^3 \\ &= 1.652 \times 10^{-23} \text{ cm}^3/\text{sphere} \end{aligned}$$

Volume of 1 mole of spheres

$$\begin{aligned} &= (6.02 \times 10^{23} \text{ spheres/mole}) \times (1.652 \times 10^{-23} \text{ cm}^3/\text{sphere}) \\ &= 9.945 \text{ cm}^3/\text{mole} \end{aligned}$$

One can find the total volume of 1 mole of liquid N₂ by dividing the molecular weight by the density. (MW = 28)

$$\text{total volume} = \frac{28 \text{ g/mole}}{0.8081 \text{ g/cm}^3} = 34.65 \text{ cm}^3/\text{mole}$$

The volume of the empty space is equal to the volume of the spheres subtracted from the total volume.

$$\begin{aligned}\text{volume of empty space per mole} &= 34.65 \text{ cm}^3 - 9.945 \text{ cm}^3 \\ &= 24.70 \text{ cm}^3\end{aligned}$$

The percent of the volume taken up by empty space is equal to the volume of the space divided by the total volume and multiplied by 100.

$$\begin{aligned}\% \text{ of empty space} &= \frac{24.70 \text{ cm}^3}{34.65 \text{ cm}^3} \times 100 \\ &= 71.30 \%\end{aligned}$$

FREEZING POINT DEPRESSION AND BOILING POINT ELEVATION

• PROBLEM 236

The freezing point constant of toluene is 3.33°C per mole per 1000 g. Calculate the freezing point of a solution prepared by dissolving 0.4 mole of solute in 500 g of toluene. The freezing point of toluene is -95.0°C .

Solution: The freezing point constant is defined as the number of degrees the freezing point will be lowered per 1000 g of solvent per mole of solute present. The freezing point depression is related to this constant by the following equation.

$$\text{freezing pt depression} = \text{molality of solute} \times \text{freezing pt constant}$$

The molality is defined as the number of moles per 1000 g of solvent. Here, one is given that 0.4 moles of solute are added to 500 g of solvent, therefore there will be 0.8 moles in 1000 g.

$$\frac{0.4 \text{ moles}}{500 \text{ g}} = \frac{0.8 \text{ moles}}{1000 \text{ g}}$$

The molality of the solute is thus 0.8 m. One can now find the freezing point depression. The freezing point constant for toluene is 3.33° .

$$\begin{aligned}\text{freezing point depression} &= \text{molality} \times 3.33^\circ \\ &= 0.8 \times 3.33^\circ = 2.66^\circ\end{aligned}$$

The freezing point of toluene is thus lowered by 2.66° .
 freezing point of solution = $(-95^{\circ}\text{C}) - 2.66^{\circ} = -97.66^{\circ}\text{C}$.

• PROBLEM 237

By how much will 50 grams of water have its freezing point depressed if you add 30 grams (molecular weight 80) of glucose to it?

Solution: The addition of any substance to water will alter its boiling or freezing point. To determine the amount of change, you must know the concentration of solute (the substance dissolved) in the solvent (water). This information is required because the freezing point depression, ΔT_f , equals the molal freezing point constant, k_f , times the molality (m).

$$\Delta T_f = k_f (m).$$

The concept of molality refers to the number of moles of solute per 1 kilogram of solvent.

The solute, glucose, weighs 30 grams. Therefore, the number of moles of glucose is $30 \text{ g} / 180 \text{ g/mole} = 1/6$ moles. You have 50 grams of water. However, molality refers to moles per thousand grams. As such, a conversion is required; namely $50/1000$. Hence, the molality is

$$\frac{\frac{1}{6} \text{ moles}}{\frac{50}{1000} \text{ g}} = 3.33 \text{ m}$$

Recall that the amount of depression of the freezing point is defined as $\Delta T_f = (k_f m)$. k_f is given for water as $-1.86^{\circ} \text{ deg mole}^{-1}$. You calculated m. The temperature depression is thus $\Delta T_f = k_f \times m = -1.86^{\circ} \times 3.33 \text{ m} = -6.2^{\circ}$.

• PROBLEM 238

What is the freezing point of a solution of 92 g of alcohol ($\text{C}_2\text{H}_5\text{OH}$) and 500 g of H_2O ?

Solution: The freezing point is dependent on the number of solute particles. One mole of a substance dissolved in 1000 g of water lowers the freezing point 1.86° . One uses the following equation to find the freezing point depression:

$$\text{freezing pt. depression} = \text{molality of solute} \times 1.86^{\circ}.$$

Now, one must find the molality of the alcohol. The molality is defined as the number of moles in 1000 g of

H₂O. In this solution, there are 92 g of alcohol present. The number of moles is found by dividing 92 g by the molecular weight of the alcohol. MW of C₂H₅OH = 46.

$$\text{no. of moles} = \frac{92.0 \text{ g}}{46 \text{ g/mole}} = 2 \text{ moles}$$

There are 2 moles in 500 g of water.

In 1000 g of H₂O, there would be twice this amount or 4 moles. Therefore, the molality of the alcohol is 4 m.

One can now find the freezing point depression.

$$\begin{aligned}\text{freezing pt. depression} &= \text{molality} \times 1.86^\circ \\ &= 4 \times 1.86^\circ = 7.44^\circ.\end{aligned}$$

The freezing point of H₂O is 0°C.

$$\begin{aligned}\text{freezing pt of solution} &= \text{freezing pt of H}_2\text{O} - \\ &\quad \text{freezing pt of depression} \\ &= 0^\circ - 7.44^\circ = -7.44^\circ.\end{aligned}$$

• PROBLEM 239

Calculate the composition (molality) of an alcohol - water mixture which will not freeze above a temperature of - 10°C (+14°F). (MW of alcohol = 46; Freezing point constant for water (K_f) = 1.86°.)

Solution: For dilute solutions, the number of degrees that the freezing point is lowered by adding a solute to a solvent is equal to the molality of the solute times the freezing point constant (K_f) of the solvent.

$$\text{freezing point depression} = \text{Molality} \times K_f$$

Because the solvent in this case is water, the K_f for water is used in solving the above equation. The freezing point depression is found by subtracting the new freezing point of the water from the original one. The original freezing point of water is 0°C.

$$\text{freezing pt. depression} = 0 - (-10) = 10^\circ\text{C}.$$

Since the freezing point depression and the freezing point constant are known, one can now solve for the molality.

$$\text{molality} = \frac{\text{freezing point depression}}{\text{freezing point constant}}$$

$$\text{molality} = \frac{10^\circ}{1.86^\circ} = 5.4 \text{ molal}$$

Molality is defined as the number of moles of solute present per kilogram of solvent. Here, the solute is the alcohol and the solvent is water. The number of grams of the alcohol present in 1000 g of water can be found by multiplying the molality by the molecular weight of the alcohol.

$$46 \text{ g/mole} \times 5.4 \text{ moles/kg of H}_2\text{O} = 250 \text{ g/kg H}_2\text{O}$$

Therefore, if 250 g of this alcohol are added to 1000 g of water, the freezing point of the water will be lowered by 10°C.

• PROBLEM 240

The molal freezing point constant for a certain liquid is 0.500°C. 26.4 g of a solute dissolved in 250 g of this liquid yields a solution which has a freezing point 0.125° below that of the pure liquid. Calculate the molecular weight of this solute.

Solution: A mole of a substance in 1000 g of H₂O gives a definite and known lowering of the freezing point. By determining the freezing point of a solution of known concentration, one can calculate the molecular weight of the dissolved substance. A general formula may be developed for this kind of calculation:

$$\text{molecular weight} = \frac{\text{grams solute}}{\text{kg of solvent}} \times \frac{K_f}{\Delta T_f}$$

where K_f is the freezing point constant and ΔT_f is change in the freezing point. Here, 26.4 g of a solute is dissolved in 250 g of liquid. One can find the number of grams of solute using the following ratio: (Note: there are 1000 g in 1 kg.) Let X = number of g of solute in 1000 g

$$\frac{26.4 \text{ g}}{250 \text{ g}} = \frac{X}{1000 \text{ g}}$$

$$X = \frac{1000 \text{ g} \times 26.4 \text{ g}}{250 \text{ g}} = 105.6 \text{ g}$$

One can now solve for the molecular weight.

$$\begin{aligned} \text{molecular weight} &= \frac{105.6 \text{ g}}{1 \text{ kg}} \times \frac{0.500^\circ\text{C}}{0.125^\circ} \\ &= 422.4 \text{ g} \end{aligned}$$

Liquid naphthalene normally freezes at 80.2°C. When 1 mole of solute is dissolved in 1000 g of naphthalene, the freezing point of the solution is 73.2°C. When 6.0 g of sulfur is dissolved in 250 g of naphthalene, the freezing point is 79.5°C. What is the molecular weight of sulfur?

Solution: In order to determine the molecular weight of sulfur, we must determine how many moles of sulfur corresponds to 6.0 g.

When 1 mole of solute is added to 1000 g of naphthalene, the normal freezing point is lowered by 7°C, from 80.2°C to 73.2°C. This corresponds to a concentration of 1 molal (1 mole solute/1000 g solvent = 1 molal). The ratio of sulfur to naphthalene in the sulfur-naphthalene solution is 6.0 g sulfur/250 g naphthalene = 0.024. Thus, adding 6.0 g of sulfur to 250 g of naphthalene is equivalent to adding 24 g of sulfur to 1000 g of naphthalene (because 24 g sulfur/1000 g naphthalene is also 0.024).

The observed freezing point depression in the sulfur-naphthalene solution is 0.7°C (from 80.2°C to 79.5°C). This is the freezing point depression one would obtain by adding 24 g of sulfur to 1000 g of naphthalene. But this is one-tenth the lowering one would obtain by adding 1 mole of sulfur (the lowering would then be 7.0°C), hence 24 g of sulfur must correspond to one-tenth of a mole.

Thus, the apparent molecular weight of sulfur is

$$\frac{24 \text{ g}}{0.1 \text{ mole}} = 240 \text{ g/mole.}$$

A chemist wishes to determine the molecular weight and molecular formula of fructose (a sugar). He places .946 g of it in 150 g of H₂O (water) and finds that the freezing point of water is depressed to -0.0651°C. Determine the molecular weight and formula of fructose, assuming that the simplest formula of fructose is (CH₂)_nO.

Solution: To answer this question, you must know the quantitative relationship for the depression of the freezing point, ΔT_f . This relation can be expressed as $\Delta T_f = k_f \times m$, where ΔT_f = the actual depression, k_f = molal freezing point constant (-1.86 deg mol⁻¹ for water) and m = molality of the solution. Molality is defined as moles of solute per 1 kg of solvent.

In this problem, water is the solvent and the fructose compound is the solute. As such, you can express

the freezing point relationship as

$$\Delta T_f = k_f \times \frac{\text{grams solute}}{\text{molecular weight}} \times \frac{\text{grams solvent}}{\text{grams solvent}} \times 1000, \quad \text{where}$$

the numerator is the number of moles of solute. You possess all of the unknowns, except the molecular weight, which can be calculated. In other words, solve for the molecular weight by substitution. Hence.

$$-0.0651^\circ\text{C} = -1.86^\circ\text{C/mole} \times \frac{.946 \text{ g}}{\text{mol. wt.}} \times 1000 \text{ g/kg}$$

$$\text{mol wt.} = 180 \text{ g/mole.}$$

To determine the molecular formula, you must figure out what formula has a molecule weight of 180 g, and yet is still a multiple of $(\text{CH}_2)\text{O}$.

Thus, one determines the weight of 1 $(\text{CH}_2)\text{O}$ and divides this into 180 g. (MW of $(\text{CH}_2)\text{O} = 30$.)

$$\text{no. of } (\text{CH}_2)\text{O} \text{ in fructose} = \frac{180 \text{ g}}{30 \text{ g}/(\text{CH}_2)\text{O}} = 6 (\text{CH}_2)\text{O}$$

Therefore, the formula for fructose is $6 \times (\text{CH}_2)\text{O}$ or $\text{C}_6\text{H}_{12}\text{O}_6$.

• PROBLEM 243

A chemist discovered a new compound and wished to determine its molecular weight. He placed 2.00 grams of the unknown compound in 10 grams of water and found that the normal freezing point of the water was depressed to -1.22°C . What is the molecular weight of the new compound?

Solution: To answer this question, you must know the quantitative relationship for the depression of the freezing point, ΔT_f . This relationship can be expressed as $\Delta T_f = k_f \times m$, where ΔT_f is the actual depression, k_f = molal freezing point constant ($-1.86 \text{ deg mole}^{-1}$ for water), m = molality of the solution. Molality is defined as the number of moles of solute per 1 kilogram of solvent. In this problem, water is the solvent and the unknown compound is the solute. You can express the freezing point relationship as

$$\Delta T_f = k_f \times \frac{\text{g.solute/mol.wt.}}{\text{g.solvent}} \times 1000,$$

where the numerator is the number of moles of solute. You have all of the unknowns, except the molecular weight. Solving for the molecular weight of the unknown:

$$- 1.22^{\circ} = - 1.86^{\circ} \times \frac{2/\text{mol.wt.}}{10} \times 1000.$$

$$\text{mol. wt.} = 305 \text{ g/mole.}$$

• PROBLEM 244

Calculate the approximate freezing point of a solution of 162 g of HBr in 500 g H₂O, assuming that the acid is 90% ionized.

Solution: The freezing point of a solution is a colligative property, which means that it depends upon the number of solute particles present in the solution. The depression in the freezing point is related to the number of particles by the equation:

$$\text{freezing pt. depression} = K_f \times \text{molality of solute.}$$


where K_f is the freezing point constant. The freezing point constant is defined as the number of degrees the freezing point is lowered when 1 mole of solute is added to 1 kg of solvent. K_f for H₂O is 1.86°. One now must determine the molality of the solution. The molality of unionized HBr is found by first determining the number of moles present. (MW of HBr = 81.) The number of moles present equals the weight divided by the molecular weight.

$$\text{no. of moles} = \frac{162 \text{ g}}{81 \text{ g/mole}} = 2 \text{ moles}$$

One can now find the molality by dividing the number of moles by the number of kilograms of solvent (1000 g = 1 kg)

$$\text{molality} = \frac{2 \text{ moles}}{0.5 \text{ kg}} = 4 \text{ m}$$

The molality of the unionized HBr is 4.0 m, but one is given that the compound is 90% ionized. This means that the total number of particles in the solution will increase. One must therefore calculate the effective molality. HBr ionizes as shown:



One HBr forms, one H⁺ and one Br⁻. Because the HBr is 90% ionized, only 10% of the original HBr is left and .90 × molality of H⁺ and Br⁻ ions are formed. The effective molality of the solution is the sum of the concentrations of the three components.

$$\text{HBr} \quad .10 \times 4\text{m} = .4 \text{ m}$$

$$\text{H}^+ \quad .90 \times 4\text{m} = 3.6 \text{ m}$$

$$\text{Br}^- \quad .90 \times 4\text{m} = \underline{3.6 \text{ m}}$$

$$\text{effective molality} = 7.6 \text{ m}$$

One can now find the amount that the freezing point has lowered.

$$\text{freezing pt. depression} = K_f \times \text{eff. molality}$$

$$\text{freezing pt. depression} = 1.86^\circ \times 7.36 = 13.7^\circ$$

The freezing point of water is lowered by 13.7° . It was originally 0°C , it is now -13.7°C .

• PROBLEM 245

What percent of the AB particles are dissociated by water if the freezing point of a (.0100 m) AB solution is -0.0193°C ? The freezing point lowering constant of water is $-1.86^\circ\text{C}/\text{mole}$.

Solution: To answer this question, you must determine the number of particles dissociated and divide that number by the initial number of particles available for dissociation. To accomplish this, you must be able to relate quantitatively the freezing point depression with the concentration of particles in solution. From the equation, $\Delta T_f = k_f m$, where ΔT_f = the depression of the freezing point, k_f = the freezing point depression constant, and m = molality of the solution, one can determine the concentration of the particles. From the data given, the total concentration of particles,

$$m = \frac{\Delta T_f}{k_f} = \frac{-0.0193^\circ\text{C}}{-1.86^\circ\text{C}/\text{m}} = .0104 \text{ m}$$

With this in mind, let x be the concentration of AB particles that dissociate. As such, you have x molal for A^+ and x molal for B^- , since $AB \rightarrow A^+ + B^-$. You are told that you start with a molal solution of .0100. Therefore, the number of particles undissociated is $.01 - x$. As such, the total concentration of particles is $x + x + (.0100 - x)$, which is equal to $.0100 + x$. Before, however, you said it was $.0104 \text{ m}$ from the equation $\Delta T_f = k_f m$. It follows then, that you can equate the two. Thus, $.0104 \text{ m} = (.0100 + x)\text{m}$. Solving for x , you obtain, $x = .0004 \text{ m}$.

This number, $x = .0004 \text{ m}$, represents the molal concentrations of each A^+ and B^- . Therefore, it must be the number of particles that dissociated. You started with $.0100 \text{ m}$ concentration of particles. Therefore, the percent that dissociated is

$$\frac{.0004}{.0100} = .04 = 4\%$$

What is the approximate boiling point at standard pressure of a solution prepared by dissolving 234 g of NaCl in 500 g of H₂O?

Solution: The boiling point of a solution is a colligative property. This means that it depends on the ratio of number of solute particles to number of solvent particles or molality. Molality is defined as the number of moles of solute divided by the number of kilograms of solvent.

When water is the solvent the boiling point constant, K_b , is 0.52°. This means that when one mole of solute is in 1 kg of H₂O the boiling point of water is raised 0.52°.

$$\text{boiling pt elevation} = K_b \times \text{molality of solute}$$

One can find the boiling point elevation once the molality of the NaCl is found. To determine the molality one must first determine the number of moles present. This is done by dividing the number of grams present by the molecular weight. (MW of NaCl = 58.5.)


$$\text{no. of moles} = \frac{234 \text{ g}}{58.5} = 4.0 \text{ moles}$$

There are 4.0 moles of NaCl in 500 g, or 0.5 kg, of H₂O. The molality can now be found.

$$\text{molality} = \frac{\text{moles of solute}}{\text{kg of solvent}}$$

$$\text{molality} = \frac{4.0 \text{ moles}}{0.5 \text{ kg}} = 8.0 \text{ m}$$

NaCl is an electrolyte which means that when it is placed in H₂O it dissociates into its ions.



This means that for every NaCl present, there are two ions formed. There are thus twice as many moles of particles as molecules of NaCl present. The effective molality is, therefore, twice the true molality.

$$\text{effective molality} = 2 \times 8.0 \text{ m} = 16.0 \text{ m}$$

One can now solve for the boiling point elevation.

$$\text{boiling point elevation} = 0.52^\circ \times 16.0 \text{ m} = 8.32^\circ$$

The boiling point of the water is raised 8.32°. It was originally 100°C, therefore the new boiling point is 100°C + 8.32° = 108.32°C.

• PROBLEM 247

Ethanol boils at 78.5°C. If 10 g of sucrose ($C_{12}H_{22}O_{11}$) is dissolved in 150 g of ethanol, at what temperature will the solution boil? Assume $K_b = 1.20^\circ\text{C/M}$ for the alcohol.

Solution: When a non-volatile solute, such as sucrose, is dissolved in a solvent, such as ethanol, it will raise the boiling point of the solvent. The boiling point elevation can be found by using the equation

$$\Delta T_b = K_b \eta,$$

where ΔT_b is the boiling point elevation, K_b is the elevation constant, and η is the molality of the solution. You want to determine ΔT_b and are given K_b . Thus, you need to determine the molality of the solution. Molality is defined as the number of moles of solute per 1 kg of solvent, i.e. moles solute/1 kg solvent.

Solving for molarity of sucrose: (MW of $C_{12}H_{22}O_{11} = 342.3$).

$$\begin{aligned} \text{Moles of solute} &= \frac{\text{grams solute}}{\text{molecular wt. solute}} \\ &= \frac{10 \text{ g}}{342.3 \text{ g/mole}} = 2.921 \times 10^{-2} \text{ moles.} \end{aligned}$$

You have 150 g of solvent, ethanol. But molality is per 1 kg, so that you must multiply 150 g by 1 kg/1000 g.

$$\text{Molality} = \frac{.02921 \text{ mole}}{\frac{150}{1000} \text{ kg}} = .195 \text{ M}$$

Thus, the elevation of the boiling point is

$$\Delta T_b = K_b \eta = (1.20^\circ\text{C/M})(.195 \text{ M}) = .23^\circ\text{C}.$$

Thus, the boiling point of the solution is

$$78.5^\circ\text{C} + .23^\circ\text{C} = 78.73^\circ\text{C}.$$

• PROBLEM 248


Give the approximate boiling points at sea level for the following: (a) 2 molal HBr; (b) Suspension of 100 g of powdered glass in one liter of water; (c) 1.2×10^{24} sucrose molecules/liter; (d) 0.5 molal BaCl_2 .

Solution: The boiling point is a colligative property and, therefore, depends upon the number of particles present in 1 kg of solvent. The boiling point constant for water is 0.52°C . This means that for each mole of particles dissolved in 1 kg of H_2O the boiling point will be elevated by 0.52°C .

$$\text{boiling pt. elevation} = 0.52^{\circ}\text{C} \times \text{molality of solute.}$$

The molality is defined as the number of moles in one kg. of solvent.

(a) 2 molal HBr. HBr is a strong acid and will thus ionize completely when diluted with H_2O to form H^+ and Br^- ions.



Therefore, 2 particles will be formed by each HBr that ionizes. The effective molality of the solution is then twice the molality of the HBr.

$$\text{effective molality} = 2 \times 2 \text{ molal} = 4 \text{ molal.}$$

One can now find the boiling point elevation.

$$\text{boiling pt. elevation} = 0.52^{\circ} \times 4 \text{ molal} = 2.1^{\circ}$$

The normal boiling point of H_2O is 100°C the new boiling point is $100^{\circ} + 2.1^{\circ}$ or 102.1°C .

(b) Suspension of 100 g of powdered glass. These particles will not dissolve to their molecular components in H_2O , thus the boiling point of the water will not be changed.

(c) 1.2×10^{24} sucrose molecules/liter. One liter of H_2O weighs 1 kg, therefore, 1.2×10^{24} sucrose molecules are dissolved in 1 kg of H_2O . The number of moles present is equal to the number of molecules present divided by Avogadro's number, the number of particles in one mole.


$$\text{no. of moles} = \frac{1.2 \times 10^{24} \text{ molecules}}{6.02 \times 10^{23} \text{ molecules/mole}} = 2 \text{ moles}$$

Hence, there are 2 moles of sucrose in 1 kg of water, thus the solution is 2 molal. Sucrose does not ionize in water, therefore, the true molality is equal to the effective molality. Solving for the boiling point elevation:

$$\text{boiling pt. elevation} = 0.52^{\circ} \times 2 \text{ molal} = 1.04^{\circ}$$

The boiling point of this solution is $100^{\circ} + 1.04^{\circ}$ or 101.04° .

(d) 0.5 molal BaCl_2 . BaCl_2 is a strong electrolyte and will completely ionize in H_2O .



3 ions will be formed for each BaCl_2 present. The effective molality will therefore be 3 times the molality of unionized BaCl_2 .

$$\text{effective molality} = 3 \times 0.5 \text{ molal} = 1.5 \text{ molal}$$

One can now solve for the boiling point elevation.

$$\text{boiling pt. elevation} = 1.5 \text{ molal} \times 0.52^\circ = 0.78^\circ$$

The boiling point of H_2O , 100° , will be raised 0.78° to be 100.78°C .

• PROBLEM 249

The normal boiling point of benzene is 80.10°C . When 1 mole of a solute is dissolved in 1000 g of benzene, the boiling point of the resulting solution is 82.73°C . When 1.2 g of elemental sulfur is dissolved in 50 g of benzene, the boiling point of the solution is 80.36°C . What is the molecular weight of sulfur?

Solution: In order to determine the molecular weight of sulfur, we must determine how many moles of sulfur correspond to 1.2 g.

When 1 mole of solute is added to 1000 g of benzene, the normal boiling point is raised by 2.63°C , from 80.10°C to 82.73°C . This corresponds to a concentration of 1 molal (1 mole solute/1000 g solvent = 1 molal). The ratio of sulfur to benzene in the sulfur-benzene solution is $1.2 \text{ g sulfur}/50 \text{ g benzene} = 0.024$. Thus, adding 1.2 g of sulfur to 50 g of benzene is equivalent to adding 24 g of sulfur to 1000 g of benzene (because $24 \text{ g sulfur}/1000 \text{ g benzene}$ is also 0.024).

The observed boiling point elevation in the sulfur-benzene solution is 0.26°C (from 80.10°C to 80.36°C). This is the boiling point elevation one would obtain by adding 24 g of sulfur to 1000 g of benzene. But this is about one-tenth the rise one would obtain by adding 1 mole of sulfur (the rise would then be 2.63°C), hence 24 g of sulfur must correspond to about one-tenth of a mole.

Thus, the apparent molecular weight of sulfur is

$$\frac{24 \text{ g}}{0.1 \text{ mole}} = 240 \text{ g/mole.}$$

• PROBLEM 250

It is known that at 4.6 mm Hg, the boiling point of water is lowered to 0°C . But at 4.6 mm Hg, the freezing point is still 0°C . Explain (a) why the freezing point was not

altered substantially by a change in pressure, and (b) the paradox of water freezing and boiling at the same temperature.

Solution: To solve this problem, you want to consider what would alter the temperature of any phase change and see if the condition exists in this case. The temperature of any phase transition is affected by pressure only because of the accompanying volume change. Since the difference in density between solid and liquid is usually small, as compared to liquid and gas, a pressure change will not alter to any significant degree the freezing point, since the volume change will be small. Thus, the boiling point was altered, but not the freezing point. The paradox of both the boiling point and freezing point being at the same point can be explained by a discussion of the triple point. At this point, some molecules are going from liquid to solid and others from solid to liquid, and similarly for the gas-liquid and liquid-solid transformations. An equilibrium is established when an equal number of molecules are going each way in all three processes. Thus, at the triple point, you have a set of conditions in which all three states may exist in equilibrium with each other. The freezing and boiling points, when at the same temperature, suggest such an equilibrium. Thus, the existence of one temperature for both boiling and freezing point can occur at a given pressure.

• PROBLEM 251

A G.T.O. has a 22 gal. cooling system. Suppose you fill it with a 50-50 solution by volume of $(\text{CH}_2\text{OH})_2$, ethylene glycol, and water. At what temperature would freezing become a problem? Assume the specific gravity of ethylene glycol is 1.115 and the freezing point depression constant of water is $1.86^\circ\text{C}/\text{mole}$. You might have placed in methanol (CH_3OH) instead of $(\text{CH}_2\text{OH})_2$. If the current cost of ethylene glycol is 12 cents/lb and the cost of methanol is 8 cents/lb, how much money would you save by using CH_3OH ? And yet, ethylene glycol is the more desirable antifreeze. Why? density = .79 g/ml for CH_3OH and 3.785 liter = 1 gallon

Solution: To determine the freezing point of the water-ethylene glycol solution, you employ the following equation

$$\Delta T_f = \frac{1000g K_f (N_2)}{M_1 (N_1 + N_2)}$$

where ΔT_f is the change in the freezing point of the solvent, water, K_f the freezing point depression constant of the solvent, M_1 is the molecular weight of the solvent, N_1 the number of moles of solvent and N_2 the number of moles of solute.

You are given K_f for water and the molecular weight

is 18; hence, you can solve for ΔT_f after determining N_1 and N_2 .

You can calculate the number of moles of water and of ethylene glycol present by using their densities and volumes. Use the following steps:

1) Convert 22 gal (the volume of the tank) to liters.
volume = 22 gal \times 3.785 l/gal = 83.3 l

2) You are told that the mixture is 50-50 by volume or one half water and one half ethylene glycol. Hence there are 83.3/2 l or 41.68 l of each.

3) Use the density of each compound to find the number of grams of each present. (density of H_2O = 1.0 and 1000 cm^3 = 1 l)

$$\begin{aligned}\text{mass of } H_2O &= 1.0 \text{ g/cm}^3 \times 1000 \text{ cm}^3/\text{l} \times 41.68 \text{ l} \\ &= 41.68 \times 10^3 \text{ g } H_2O\end{aligned}$$

$$\begin{aligned}\text{mass of } (CH_2OH)_2 &= 1.115 \text{ g/cm}^3 \times 1000 \text{ cm}^3/\text{l} \times 41.68 \text{ l} \\ &= 46.47 \times 10^3 \text{ g of } (CH_2OH)_2\end{aligned}$$

4) The number of moles of each is found by dividing its mass by its molecular weight. (MW of H_2O = 18, MW of $(CH_2OH)_2$ = 62).

$$\text{moles of } H_2O = \frac{41.68 \times 10^3 \text{ g}}{18 \text{ g/mole}} = 2.32 \times 10^3 \text{ moles}$$

$$\text{moles of } (CH_2OH)_2 = \frac{46.47 \times 10^3 \text{ g}}{62 \text{ g/mole}} = 7.50 \times 10^2 \text{ moles}$$

You can now solve for ΔT_f by substituting into the above equation

$$\begin{aligned}\Delta T_f &= \frac{1000 \text{ g } (1.86 \text{ }^\circ\text{C/mole}) (7.50 \times 10^2 \text{ mole})}{(18 \text{ g/mole}) (2.32 \times 10^3 + 7.50 \times 10^2 \text{ moles})} \\ &= \frac{1.40 \times 10^6}{5.53 \times 10^4} = 25.2 \text{ }^\circ\text{C}\end{aligned}$$

The normal freezing point of water is 0°C , after adding the ethylene glycol, the freezing point becomes $0^\circ - 25.2^\circ\text{C}$ or -25.2°C .

Determine the amount of money it is possible to save by using a methanol-water mixture by calculating how much 41.68 l (half of the volume of the tank) of methanol costs and subtracting this value from the cost of 41.68 l of ethylene glycol. To calculate the cost of each

1) 1 lb = 454 g. Since the costs are given in cents/lb, determine the weight in pounds of 41.68 l of each. Earlier the weight of 41.68 l of ethylene glycol was found to be $46.47 \times 10^3 \text{ g}$. Converting this to pounds,

$$\text{weight of } (\text{CH}_2\text{OH})_2 = 46.47 \times 10^3 \text{ g} \times \frac{1 \text{ lb}}{454 \text{ g}} = 102.4 \text{ lb}$$

$$\begin{aligned}\text{weight of } (\text{CH}_3\text{OH}) &= \text{density} \times \text{volume} \\ &= .79 \text{ g/ml} \times 10^3 \text{ ml/l} \times 41.68 \text{ l} \\ &= 33.07 \times 10^3 \text{ g}\end{aligned}$$

Converting to pounds

$$\text{weight of } \text{CH}_3\text{OH} = 33.07 \times 10^3 \text{ g} \times \frac{1 \text{ lb}}{454 \text{ g}} = 72.8 \text{ lb.}$$

The cost of each is found by multiplying the price per pound by the weight of each.

$$\text{cost of } (\text{CH}_2\text{OH})_2 = 102.4 \text{ lb} \times \$12/\text{lb} = \$12.29$$

$$\text{cost of } (\text{CH}_3\text{OH}) = 72.8 \text{ lb} \times \$0.08/\text{lb} = \$5.82$$

The difference in cost is $\$12.29 - \$5.82 = \$6.47$. Hence, you save \$6.47 by using methanol.

You have found that a 50-50 mixture of water and ethylene glycol protects the car to a temperature of -25.2°C . The most desirable antifreeze will protect the car at the lowest temperature, hence, to determine whether ethylene glycol is more desirable than methanol, calculate the lowest temperature at which the car will be protected by methanol. This is accomplished by using the above equation for freezing point depression.

$$\Delta T_f = \frac{1000 \text{ g } (K_f) (N_2)}{M_1 (N_2 + N_1)}$$

Previously, one has determined K_f , N_1 and M_1 to be $1.86^\circ\text{C}/\text{mole}$, 2.32×10^3 moles and $18 \text{ g}/\text{mole}$, respectively. Thus, to solve for ΔT_f determine N_2 first. As shown, the weight of the methanol is $33.07 \times 10^3 \text{ g}$. Find N_2 by dividing this weight by the molecular weight of methanol, 32.

$$\begin{aligned}\text{No. of moles of methanol } (N_2) &= \frac{33.07 \times 10^3 \text{ g}}{32 \text{ g/mole}} \\ &= 1.03 \times 10^3 \text{ moles.}\end{aligned}$$

Solving for ΔT_f :

$$\begin{aligned}\Delta T_f &= \frac{1000 \text{ g } (1.86^\circ\text{C}/\text{mole}) (1.03 \times 10^3 \text{ moles})}{(18 \text{ g/mole}) (1.03 \times 10^3 + 2.32 \times 10^3 \text{ moles})} \\ &= 31.8^\circ\text{C.}\end{aligned}$$

The methanol will protect the car at temperatures as low as 0°C - 31.8°C or -31.8°C . This is fine while the car is not

in motion and is cold, but when the car is driven the engine becomes very warm. Methanol has a boiling point of 60°C and will evaporate while the engine is operating, whereas the ethylene glycol has a much higher boiling point and will not evaporate. Thus, if methanol is used as an anti-freeze it will have to be replaced after each use of the car while the ethylene glycol will last all winter. In the end it is far more expensive and troublesome to use methanol.

RAOULT'S LAW AND VAPOR PRESSURE

• PROBLEM 252

You have two 1-liter containers connected to each other by a valve which is closed. In one container, you have liquid water in equilibrium with water vapor at 25°C . The other container contains a vacuum. Suddenly, you open the valve. Discuss the changes that take place, assuming temperature is constant with regard to (a) the vapor pressure, (b) the concentration of the water molecules in the vapor, (c) the number of molecules in the vapor state.

Solution: The vapor pressure is the pressure exerted by the gas molecules when they are in equilibrium with the liquid. When the valve is opened some of the gas molecules will move to the empty container. At this point the pressure will be less than the equilibrium pressure because the concentration of the gas molecules will be lowered. Very quickly, though, the equilibrium will be attained again by the action of more liquid molecules vaporizing. Therefore, the vapor pressure and the concentration of the gaseous molecules of the system remains essentially unchanged.

Since the concentration of the gaseous molecules remains unchanged when the volume of the system is doubled, the number of molecules must also be doubled. This is true because concentration is an expression of the number of molecules per unit volume.

• PROBLEM 253

A chemist decides to find the vapor pressure of water by the gas saturation method. 100 liters of N_2 gas is passed through 65.44 g of water. After passage of the gas, 63.13 g remained. The temperature of the H_2O (water) is 25°C . Find the vapor pressure of water at this temperature.

Solution: In the gas saturation method, a dry, unreactive gas such as nitrogen or air is bubbled through a specific amount

of liquid maintained at constant temperature. After the gas has been bubbled away, the loss in weight of the liquid is determined. The weight loss is the number of grams of liquid in the vapor state. There exists an equation that relates the volume, pressure, weight loss, and molecular weight of the liquid.

$P = \frac{g}{MV} RT$, where P = vapor pressure, g = grams of vapor, M = molecular weight of liquid, R = universal gas constant ($.0821 \frac{\text{liter-atm}}{\text{mole}^\circ\text{K}}$), V = volume, and T = temperature in Kelvin (Celsius plus 273°)

The pressure will be expressed in mm, so that you use the conversion factor of $\frac{760 \text{ mm}}{\text{atm}}$. $g = 65.44 - 63.13 = 2.31$ grams or the weight of the liquid in the vapor state.

The molecular weight of water = 18.02 g/mole . Thus,

$$P = \frac{gRT}{MV} = \frac{(2.31 \text{ grams}) (.0821 \frac{\text{liter atm}}{\text{mole}^\circ\text{K}}) (298^\circ\text{K}) (\frac{760 \text{ mm}}{1 \text{ atm}})}{(18.02 \text{ g/mole}) (200 \text{ liters})}$$

$$= 23.8 \text{ mm} = \text{vapor pressure of } \text{H}_2\text{O}.$$

• PROBLEM 254

When a swimmer leaves cold water on a warm, breezy day, he experiences a cooling effect. Why?

Solution: The equilibrium vapor pressure, is defined as the pressure exerted by the gas of that substance in equilibrium with the liquid state of the same substance. When the swimmer leaves the cold water, he is, coated with water. As the water evaporates a layer of gaseous water molecules form around the swimmer's body.

This vapor attains an equilibrium with the atmosphere which hinders further vaporization of the water. However, when the breeze comes, it blows away the vapor above the water, and thereby keeps the partial pressure of the vapor low above the skin. This, then, causes increased evaporation so that the partial pressure can be reestablished. This is true because the concentration and vapor pressure of the gaseous water molecules must be below a certain point for evaporation to occur. Once this point has been reached no more liquid molecules of water can go into the gaseous state.

For the water to evaporate, it must go from a liquid to a gaseous phase, a process which requires heat. The water removes heat from the swimmer's body. Thus, the swimmer feels a cooling effect.

The vapor pressures of pure benzene and toluene at 60°C are 385 and 139 Torr, respectively. Calculate (a) the partial pressures of benzene and toluene, (b) the total vapor pressure of the solution, and (c) the mole fraction of toluene in the vapor above a solution with 0.60 mole fraction toluene.

Solution: The vapor pressure of benzene over solutions of benzene and toluene is directly proportional to the mole fraction of benzene in the solution. The vapor pressure of pure benzene is the proportionality constant. This is analogous to the vapor pressure of toluene. This is known as Raoult's law. It may be written as

$$P_1 = X_1 P_1^\circ$$

$$P_2 = X_2 P_2^\circ$$

where 1 and 2 refer to components 1 and 2, P_1 and P_2 represent the partial vapor pressure above the solution, P_1° and P_2° are the vapor pressures of pure components, and X_1 and X_2 are their mole fractions. Solutions are called ideal if they obey Raoult's law.

The mole fraction of a component in the vapor is equal to its pressure fraction in the vapor. The total vapor pressure is the sum of the vapor's component partial pressures.

To solve this problem one must

- 1) calculate the partial pressures of benzene and toluene using Raoult's law
- 2) find the total vapor pressure of the solution by adding the partial pressures
- 3) find the mole fraction of toluene in the vapor.

One knows the mole fraction of toluene in the solution is 0.60 and, thus, one also knows the mole fraction of benzene is (1-0.60) or 0.40. Using Raoult's law:

$$P_{\text{benzene}}^\circ = 385 \text{ Torr} \quad P_{\text{toluene}}^\circ = 139$$

$$\text{a) } P_{\text{benzene}} = (0.40)(385 \text{ Torr}) = 154.0 \text{ Torr}$$

$$P_{\text{toluene}} = (0.60)(139 \text{ Torr}) = 83.4 \text{ Torr}$$

$$\text{b) } P_{\text{total}} = 154.0 + 83.4 = 237.4 \text{ Torr}$$

$$\text{c) The mole fraction of toluene in the vapor =}$$

$$X_{\text{toluene, vap}} = \frac{P_{\text{toluene}}}{P_{\text{toluene}} + P_{\text{benzene}}} = \frac{83.4}{237.4} = 0.351.$$

• PROBLEM 256

At 90°C, benzene has a vapor pressure of 1022 Torr, and toluene has a vapor pressure of 406 Torr. From this information, calculate the composition of the benzene-toluene solution that will boil at 1 atm pressure and 90°C, assuming that the solution is ideal.

Solution: A solution boils when the sum of the partial pressures of the components becomes equal to the applied pressure (i.e. total pressure). To solve this problem, one must realize that the applied pressure is atmospheric pressure, 760 Torr.

One can determine the partial pressure of benzene (P_a°) and of toluene (P_b°) by using Raoult's law. Raoult's law states that the partial pressure p of a gas is equal to its vapor pressure p° times its mole fraction X in the solution.

$$p = p^\circ X$$

Dalton's law of partial pressure states that the sum of the partial pressures of a system is equal to the total pressure of the system. For this particular benzene-toluene solution to boil the total pressure of the system must equal atmospheric pressure 760 Torr. If one lets the mole fraction of benzene equal X_a , then the mole fraction toluene is equal to $1 - X_a$. One determines the partial pressures by substituting into Raoult's law

$$P_a = P_a^\circ (X_a)$$

$$P_b = P_b^\circ (1 - X_a).$$

Then, substituting into Dalton's law:

$$760 \text{ Torr} = P_a + P_b$$

$$760 \text{ Torr} = 1022 X_a + 406 (1 - X_a)$$

$$= 1022 X_a + 406 - 406 X_a$$

$$X_a = 0.574$$

The mole fraction of benzene in the liquid is 0.574 and the mole fraction of toluene is $(1 - 0.574) = 0.426$.

• PROBLEM 257

A chemist dissolves 300 g of urea in 1000 g of water. Urea is NH_2CONH_2 . Assuming the solution obeys Raoult's law, determine the following: a) The vapor pressure of the solvent at 0° and 100°C and b) The boiling and freezing point of the solution. The vapor pressure of pure water is 4.6 mm and 760 mm at 0°C and 100°C, respectively and the $K_f = 1.86^\circ\text{C}/\text{mole}$ and $K_b = .52^\circ\text{C}/\text{mole}$.

Solution: To solve (a), you must employ Raoult's law, which states $P_{H_2O} = P^{\circ}_{H_2O} X_{H_2O}$, where P_{H_2O} = the partial pressure of water (the solvent), $P^{\circ}_{H_2O}$ = the vapor pressure of water when pure, and X_{H_2O} is the mole fraction of H_2O in the solution. The mole fraction is equal to $\frac{N_{H_2O}}{N_{\text{solute}} + N_{H_2O}}$,

where N_{H_2O} = moles of H_2O , and N_{solute} = moles of solute.

You are told the $P^{\circ}_{H_2O}$ for H_2O at the temperatures in question. Thus, to find the vapor pressures of the solution, you must calculate X_{H_2O} and substitute into Raoult's law.

To calculate the concentration of urea, remember moles = $\frac{\text{grams}}{\text{molecular weight}}$. Since you have 300 grams of urea and the molecular weight of urea is 60.06 g, you have $\frac{300}{60.06}$ or 4.995 moles of urea. Similarly, for water (MW=18), you have $\frac{1000 \text{ g}}{18 \text{ g/mole}} = 55.55$ moles of water. Thus, the mole fraction of water = $\frac{55.55}{4.995 + 55.55} = .917$. Therefore, at 0° ,

$$(P_{H_2O} = P^{\circ}_{H_2O} X_{H_2O} = (4.6 \text{ mm})(.917) = 4.2 \text{ mm}) \text{ at } 760^{\circ},$$

$(P_{H_2O} = P^{\circ}_{H_2O} X_{H_2O} = (760)(.917) = 697 \text{ mm})$ To solve (b), you use the equations $\Delta T_f = K_f m$, where ΔT_f = freezing point depression, K_f = molal freezing pt. depression constant, and m = molality and $\Delta T_b = K_b m$, where ΔT_b = boiling point elevation, K_b = molal boiling point depression constant. If you find ΔT_f , you can calculate the freezing point, since the normal freezing point is decreased by this amount to give you the new freezing point. Boiling point works in the same way, except that you add to the normal boiling point. To calculate ΔT_f and ΔT_b , you must know molality. Molality = moles solute per 1 kilogram solvent. You have 4.995 moles of urea and

1000 g of water or solvent, so that $\frac{4.995 \text{ mole}}{1 \text{ kg}} \approx 5 \text{ molal}$ solution,

Thus, $\Delta T_f = (1.86)(5) = 9.3^{\circ}\text{C}$ and $\Delta T_b = (.52)(5) = 2.6^{\circ}\text{C}$. Therefore, the freezing point is now $0.0^{\circ}\text{C} - 9.3^{\circ}\text{C} = -9.3^{\circ}\text{C}$ and the boiling point is $100^{\circ}\text{C} + 2.6^{\circ}\text{C} = 102.6^{\circ}\text{C}$.

• PROBLEM 258

A solution of 20.0 g of a non-volatile solute in 100 g of benzene at 30°C has a vapor pressure 13.4 torr lower than the vapor pressure of pure benzene. What is the mole

fraction of solute? Vapor pressure of benzene at 30°C = 121.8 torr.

Solution: At constant temperature, the lowering of the vapor pressure by a non-volatile solute is proportional to the concentration of the solute in the solution (the mole fraction). This is called Raoult's Law. The mole fraction of a solute is defined as the number of moles of solute divided by the sum of the number of moles of solute and of solvent.

$$\text{mole fraction of solute} = \frac{\text{moles solute}}{\text{moles solute} + \text{moles solvent}}$$

Raoult's Law is used to solve this problem. It can be stated:

$$P = P^{\circ} X_2,$$

where P° is the vapor pressure of the pure solvent, P is the vapor pressure of the solution, and X_2 is the mole fraction of the solute.

Using Raoult's Law for this solution:

$$13.4 \text{ torr} = (121.8 \text{ torr}) (X_2)$$

$$0.111 = X_2$$

The mole fraction of the solute is 0.111.

• PROBLEM 259

The vapor pressure of benzene at 75°C is 640 torr. A solution of 3.68 g of a solute in 53.0 g benzene has a vapor pressure of 615 torr. Calculate the molecular weight of the solute. (MW of benzene = 78.0.)

Solution: At constant temperature, the lowering of the vapor pressure by a non-volatile solute is proportional to the concentration of the solute in the solution.

$$\text{Hence, } P^{\circ} - P = P^{\circ} X_2,$$

where P° is the original pressure, P is the final pressure and X_2 is the mole fraction of the solute. Here, one solves for the mole fraction of the solute, and from that, one can determine the molecular weight.

$$X_2 = \text{mole fraction}$$

$$P^{\circ} - P = P^{\circ} X_2$$

$$P^{\circ} = 640 \text{ torr}$$

$$P = 615 \text{ torr}$$

$$\begin{aligned}
 640 \text{ torr} - 615 \text{ torr} &= (640 \text{ torr}) \cdot X_2 \\
 25 \text{ torr} &= (640 \text{ torr}) \cdot X_2 \\
 \frac{25 \text{ torr}}{640 \text{ torr}} &= X_2
 \end{aligned}$$

$$\text{mole fraction} = .039$$

Mole fraction is defined as the number of moles of each component divided by the sum of the number of moles in the solution.

mole fraction of solute =

$$= \frac{\text{no. of moles of solute}}{\text{no. of moles of solute} + \text{no. of moles of benzene}}$$

To find the number of moles of solute, one must first know the number of moles of benzene present. This is found by dividing the number of grams of benzene by the molecular weight of benzene. (MW = 78.)

$$\text{no. of moles of benzene} = \frac{\text{no. of grams}}{\text{MW}}$$

$$\text{no. of moles} = \frac{53.0 \text{ g}}{78.0 \text{ g/mole}} = .679 \text{ moles}$$

One can now solve for the number of moles of solute. Let X = moles of solute

$$\text{mole fraction of solute} = \frac{\text{moles solute}}{\text{moles benzene} + \text{moles solute}}$$

$$.039 = \frac{X}{.679 + X}$$

$$.039 (.679 + X) = X$$

$$.265 + .039X = X$$

$$.265 = .961 X$$

$$.275 = X$$

Thus there are .275 moles of solute present. One is told that there are 3.68 g of this solute, therefore, there are .275 moles in 3.68 g. The molecular weight is found by dividing 3.68 g by 0.275 moles.

$$\text{molecular weight} = \frac{3.68 \text{ g}}{0.275 \text{ mole}} = 133.8 \text{ g/mole.}$$

• PROBLEM 260

Water at 30°C has a vapor pressure of 31.82 mm Hg. When 25.0 g of ethylene glycol is added to 1000 g of water, the vapor pressure is lowered to 31.59 mm Hg. Determine the molecular weight of ethylene glycol.

Solution: This problem is an application of Raoult's law.

Let p^0 denote the vapor pressure of pure water, p the vapor pressure of the ethylene glycol - water solution, n_w the number of moles of water in the solution, and n_e the number of moles of ethylene glycol. Then from Raoult's law,

$$p = p^0 \frac{n_w}{n_e + n_w} .$$

Multiplying by minus one and adding p^0 to both sides of this equation gives

$$\begin{aligned} p^0 - p &= p^0 - p^0 \frac{n_w}{n_e + n_w} \\ &= p^0 \left(1 - \frac{n_w}{n_e + n_w} \right) \\ &= p^0 \left(\frac{n_e + n_w}{n_e + n_w} - \frac{n_w}{n_e + n_w} \right) , \end{aligned}$$

$$\text{or,} \quad p^0 - p = p^0 \frac{n_e}{n_e + n_w}$$

Since the number of moles is equal to the number of grams divided by the molecular weight, this equation becomes

$$p^0 - p = p^0 \frac{g_e/M_e}{g_e/M_e + g_w/M_w} ,$$

where g_e denotes the number of grams of ethylene glycol, g_w the number of grams of water, M_e the molecular weight of ethylene glycol (which we are trying to determine), and M_w the molecular weight of water.

For this problem, $p^0 = 31.82$ mm, $p = 31.59$ mm, $g_e = 25.0$ g, $g_w = 1000$ g, and the molecular weight of water, $M_w = 18$. Substituting into the last equation, we obtain

$$31.82 \text{ mm} - 31.59 \text{ mm} = 31.82 \text{ mm} \left(\frac{25.0 \text{ g}/M_e}{1000 \text{ g}/18 + 25.0 \text{ g}/M_e} \right) ,$$

$$\text{or,} \quad 0.23 \text{ mm} = 31.82 \text{ mm} \left(\frac{25.0 \text{ g}/M_e}{1000 \text{ g}/18 + 25.0 \text{ g}/M_e} \right) .$$

Solving for M_e ,

$$\frac{0.23 \text{ mm}}{31.82 \text{ mm}} \left(\frac{1000 \text{ g}}{18} + \frac{25.0 \text{ g}}{M_e} \right) = \frac{25.0 \text{ g}}{M_e},$$

$$\begin{aligned} \frac{0.23 \text{ mm}}{31.82 \text{ mm}} \times \frac{1000 \text{ g}}{18} &= \frac{25.0 \text{ g}}{M_e} - \frac{0.23 \text{ mm}}{31.82 \text{ mm}} \times \frac{25.0 \text{ g}}{M_e} \\ &= \frac{1}{M_e} \times 25.0 \text{ g} \left(1 - \frac{0.23 \text{ mm}}{31.82 \text{ mm}} \right), \end{aligned}$$

$$\text{or, } M_e = \frac{31.82 \text{ mm}}{0.23 \text{ mm}} \times \frac{18}{1000 \text{ g}} \times 25.0 \text{ g} \left(1 - \frac{0.23 \text{ mm}}{31.82 \text{ mm}} \right) = 62.$$

Thus, the molecular weight of ethylene glycol is 62.

CLAUSIUS-CLAPEYRON EQUATION OF VAPORIZATION

• PROBLEM 261

If the vapor pressure of ethyl alcohol, C_2H_5OH , is 0.132 atm at 34.9°C , and 0.526 atm at 63.5°C what do you predict it will be at 19.0°C ?

Solution: Equilibrium vapor pressure is the pressure exerted by a vapor when the vapor is in equilibrium with its liquid. The magnitude of the equilibrium vapor pressure depends (1) on the nature of the liquid and (2) on its temperature. The vapor pressure is related to temperature by the following equation

$$\log p = \frac{-\Delta H}{19.15 T} + C$$

where p is the vapor pressure, T is the absolute temperature, ΔH is the heat required to transform one mole of liquid to the ideal gas state, and C is a constant dependent on the liquid and units used for expressing pressure. One is given p and T for two trials, thus ΔH for ethyl alcohol can be found.

For $p = 0.132 \text{ atm}$

$$T = 34.9^\circ\text{C} + 273 = 307.9^\circ\text{K}$$

$$\text{a) } \log 0.132 = \frac{-\Delta H}{19.15 (307.9^\circ\text{K})} + C$$

For $p = 0.526 \text{ atm}$

$$T = 63.5^\circ\text{C} + 273 = 336.5^\circ\text{K}$$

$$b) \quad \log 0.526 \text{ atm} = \frac{-\Delta H}{19.15 (336.5^\circ\text{K})} + C$$

One can solve for ΔH by subtracting equation a from equation b.

$$\log 0.526 = -\Delta H/19.15(336.5^\circ\text{K}) + C$$

$$- [\log 0.132 = -\Delta H/19.15(307.9^\circ\text{K}) + C]$$

$$\log 0.526 - \log 0.132 = -\Delta H/19.15(336.5^\circ\text{K}) + \Delta H/19.15(307.9^\circ\text{K})$$

$$- .279 - (-.879) = -\Delta H/6444 + \Delta H/5896$$

$$(6444)(5896)(.60) = (-\Delta H/6444 + \Delta H/5896)(6444)(5896)$$

$$2.28 \times 10^7 = -5896\Delta H + 6444\Delta H$$

$$2.28 \times 10^7 = 548\Delta H$$

$$\frac{2.28 \times 10^7}{548} = \Delta H$$

$$4.16 \times 10^4 \text{ J/mole} = \Delta H$$

After solving for ΔH one can find the vapor pressure at 19.0°C by a similar method.

$$\text{For } p = 0.132 \text{ atm}$$

$$T = 34.9^\circ\text{C} + 273 = 307.9^\circ\text{K}$$

$$\Delta H = 4.16 \times 10^4 \text{ J/mole}$$

$$c) \quad \log 0.132 = \frac{-4.16 \times 10^4 \text{ J/mole}}{19.15 (307.9^\circ\text{K})} + C$$

$$\log 0.132 = -7.055 + C$$

$$\text{For } p = ?$$

$$T = 19.0^\circ\text{C} + 273 = 292^\circ\text{K}$$

$$\Delta H = 4.16 \times 10^4 \text{ J/mole}$$

$$d) \quad \log p = \frac{-4.16 \times 10^4 \text{ J/mole}}{(19.15)(292^\circ\text{K})} + C = -7.439 + C$$

Subtracting equation c from equation d:

$$\log p = -7.439 + C$$

$$- [\log 0.132 = -7.055 + C]$$

$$\log p - \log 0.132 = - .384$$

$$\log p = - .384 + \log 0.132$$

$$\log p = - .384 + (-.879)$$

$$\log p = -1.263$$

$$p = .0545 \text{ atm.}$$

• PROBLEM 262

If the vapor pressure of methyl alcohol, CH_3OH , is 0.0526 atm at 5.0°C and 0.132 atm at 21.2°C , what do you predict the normal boiling point will be?

Solution: The normal boiling point of a liquid is the temperature at which the vapor pressure of the liquid is equal to the prevailing atmospheric pressure. Atmospheric pressure is defined as 1 atm. The pressure is related to the temperature by the equation

$$\ln p = \frac{-\Delta H}{RT} + C$$

where p is the vapor pressure, ΔH is the heat required to transform one mole of liquid to the ideal-gas state; R is the gas constant, $8.314 \text{ J/mole } ^\circ\text{K}$; T is the absolute temperature; and C is a constant dependent on the liquid. One can solve for ΔH by using the values obtained for p at 5°C and 21.2°C .

For $p = 0.0526 \text{ atm}$

$$T = 5.0^\circ\text{C} + 273 = 278^\circ\text{K}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln (0.0526) = \frac{-\Delta H}{(8.314 \text{ J/mole } ^\circ\text{K})(278^\circ\text{K})} + C$$

a) $-2.945 = -\Delta H/2.311 \times 10^3 \text{ J/mole} + C$

For $p = 0.132 \text{ atm}$

$$T = 21.2^\circ\text{C} + 273 = 294.2^\circ\text{K}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln (0.132) = \frac{-\Delta H}{(8.314 \text{ J/mole } ^\circ\text{K})(294.2^\circ\text{K})} + C$$

b) $-2.025 = -\Delta H/2.446 \times 10^3 \text{ J/mole} + C$

One can solve for ΔH by subtracting equation a from equation b.

$$-2.025 = -\Delta H/2.446 \times 10^3 \text{ J/mole} + C$$

$$-[-2.945 = -\Delta H/2.311 \times 10^3 \text{ J/mole} + C]$$

$$.920 = -\Delta H/2.446 \times 10^3 \text{ J/mole} + \Delta H/2.311 \times 10^3 \text{ J/mole}$$

$$(2.446 \times 10^3 \text{ J/mole}) (2.311 \times 10^3 \text{ J/mole}) (.920) =$$

$$(-\Delta H/2446 \times 10^3 \text{ J/mole} + \Delta H/2.311 \times 10^3 \text{ J/mole}) (2.446 \times 10^3 \text{ J/mole}) \\ \times (2.311 \times 10^3 \text{ J/mole})$$

$$5.200 \times 10^6 \text{ J}^2/\text{mole}^2 = -(2.311 \times 10^3 \text{ J/mole}) \Delta H \\ + (2.446 \times 10^3 \text{ J/mole}) \Delta H$$

$$5.200 \times 10^6 \text{ J}^2/\text{mole}^2 = (.135 \times 10^3 \text{ J/mole}) \Delta H$$

$$\frac{5.200 \times 10^6 \text{ J}^2/\text{mole}^2}{.135 \times 10^3 \text{ J/mole}} = \Delta H$$

$$3.85 \times 10^4 \text{ J/mole} = \Delta H$$

Using a similar method the normal boiling point can be found now that one knows ΔH .

For $p = 0.0526 \text{ atm}$

$$T = 5.0^\circ\text{C} + 273 = 278^\circ\text{K}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\Delta H = 3.85 \times 10^4 \text{ J/mole}$$

$$\ln 0.0526 = \frac{-3.85 \times 10^4 \text{ J/mole}}{(8.314 \text{ J/mole } ^\circ\text{K})(278^\circ\text{K})} + C$$

c) $-2.945 = -16.657 + C$

For $p = 1.0 \text{ atm}$

$$T = ?$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\Delta H = 3.85 \times 10^4 \text{ J/mole}$$

$$\ln 1.0 = \frac{-3.85 \times 10^4 \text{ J/mole}}{(8.314 \text{ J/mole } ^\circ\text{K})T} + C$$

d) $0 = -\frac{4.63 \times 10^3 ^\circ\text{K}}{T} + C$

Subtracting equation d from equation c one can obtain T.

$$-2.945 = -16.657 + C$$

$$- \left(0 = -\frac{4.63 \times 10^3 ^\circ\text{K}}{T} + C \right)$$

$$-2.945 = -16.657 + \frac{4.63 \times 10^3 ^\circ\text{K}}{T}$$

$$13.712 = \frac{4.63 \times 10^3 ^\circ\text{K}}{T}$$

$$T = \frac{4.63 \times 10^3 \text{ } ^\circ\text{K}}{13.712} = 337.7^\circ\text{K}$$

$$\text{Boiling point in } ^\circ\text{C} = 337.7^\circ\text{K} - 273 = 64.7^\circ\text{C}.$$

• PROBLEM 263

Find the vapor pressure of CCl_4 (carbon tetrachloride) at 38°C , if $H^1 = 32,000 \text{ J/mole}$ and at 23°C , the vapor pressure is .132 atm.

Solution: To solve this problem, employ the Clausius-Clapeyron equation, which permits evaluation of vapor pressure in terms of ΔH^1 (the heat required to transform one mole of liquid to the ideal-gas state), the original vapor pressure and the two temperatures. The equation states:

$$\log \frac{p_1}{p_2} = \frac{\Delta H^1}{19.15} \left(\frac{1}{T_2} - \frac{1}{T_1} \right),$$

where p_1 and p_2 are vapor pressures and T_1 and T_2 are the temperatures in Kelvin. Let p_2 be the vapor pressure of CCl_4 at 38°C . Substitute the known values for solution,

$$\log \frac{.132}{p_2} = \frac{32,000}{19.15} \left(\frac{1}{311} - \frac{1}{296} \right).$$

$$\text{Solve for } p_2, \text{ to obtain } p_2 = .250 \text{ atm.}$$

• PROBLEM 264

Two substances A and B have the same 0.132-atm vapor pressure at 15°C , but A has a heat of vaporization of 25,000 J/mole, whereas B, 35,000 J/mole. What will be the respective normal boiling points?

Solution: The vapor pressure, heat of vaporization and absolute temperature are related in the Clausius-Clapeyron equation. This equation is written:

$$\ln \frac{p_1}{p_2} = \frac{\Delta H}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

where p_1 is the initial pressure, p_2 is the final pressure, ΔH is the heat of vaporization when T_2 is the boiling point, R is the gas constant, $8.314 \text{ J/mole } ^\circ\text{K}$, T_1 is the initial temperature and T_2 is the final temperature. To solve for the boiling point of A, note that at the boiling point of a liquid the vapor pressure is equal to the atmospheric pressure. Atmospheric pressure is 1 atm.

$$p_1 = 0.132 \text{ atm} \quad T_1 = 15^\circ\text{C} + 273 = 288^\circ\text{K}$$

$$p_2 = 1 \text{ atm} \quad T_2 = ?$$

$$\Delta H = 25,000 \text{ J/mole} \quad R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln \frac{0.132}{1} = \frac{25,000 \text{ J/mole}}{8.314 \text{ J/mole } ^\circ\text{K}} \left(\frac{1}{T_2} - \frac{1}{288^\circ\text{K}} \right)$$

$$- 2.025 = 3.00 \times 10^3 \text{ } ^\circ\text{K} (1/T_2 - 3.47 \times 10^{-3}/^\circ\text{K})$$

$$- 6.75 \times 10^{-4}/^\circ\text{K} = 1/T - 3.47 \times 10^{-3}/^\circ\text{K}$$

$$2.795 \times 10^{-3}/^\circ\text{K} = 1/T$$

$$357.78 = T$$

The boiling point of A in $^\circ\text{C}$ is

$$357.78 - 273 = 84.78^\circ\text{C}$$

Solving for the boiling point of B:

$$p_1 = 0.132 \text{ atm} \quad T_1 = 15^\circ\text{C} + 273 = 288^\circ\text{K}$$

$$p_2 = 1.0 \text{ atm} \quad T_2 = ?$$

$$\Delta H = 35,000 \text{ J/mole}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln \frac{0.132}{1.0} = \frac{35,000 \text{ J/mole}}{8.314 \text{ J/mole } ^\circ\text{K}} \left(\frac{1}{T_2} - \frac{1}{288^\circ\text{K}} \right)$$

$$- 2.025 = 4.21 \times 10^3 \text{ } ^\circ\text{K} (1/T_2 - 3.47 \times 10^{-3}/^\circ\text{K})$$

$$- 4.81 \times 10^{-4}/^\circ\text{K} = 1/T_2 - 3.47 \times 10^{-3}/^\circ\text{K}$$

$$2.989 \times 10^{-3}/^\circ\text{K} = 1/T_2$$

$$334.56^\circ\text{K} = T_2$$

The boiling point of B is equal to 334.56°K or $334.56 - 273 = 61.56^\circ\text{C}$.

• PROBLEM 265

At a 4000-m altitude the atmospheric pressure is about 0.605 atm. What boiling point would you expect for water under these conditions?

Solution: The pressure and temperature of a liquid that vaporizes are related in the Clausius-Clapeyron equation. This equation is

$$\ln \frac{P_1}{P_2} = \frac{\Delta H}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

where P_1 is the initial pressure, P_2 is the final pressure, ΔH is the heat of vaporization, R is the gas constant, $8.314 \text{ J/mole } ^\circ\text{K}$, T_2 is the boiling point at P_2 and T_1 is the boiling point at P_1 . The heat of vaporization of H_2O is

40,600 J/mole. One can assume for purposes of calculation that P_1 is 1 atm. The boiling point of H_2O at 1 atm is 373°K. Solving for the boiling point at 0.605 atm:

$$P_1 = 1 \text{ atm} \quad T_1 = 373^\circ\text{K}$$

$$P_2 = 0.605 \text{ atm} \quad T_2 = ?$$

$$\Delta H = 40,600 \text{ J/mole}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln \frac{1}{0.605} = \frac{40,600 \text{ J/mole}}{8.314 \text{ J/mole } ^\circ\text{K}} \left(\frac{1}{T_2} - \frac{1}{373^\circ\text{K}} \right)$$

$$.5025 = 4.88 \times 10^3 \text{ } ^\circ\text{K} \left(\frac{1}{T_2} - 2.681 \times 10^{-3}/^\circ\text{K} \right)$$

$$1.0297 \times 10^{-4}/^\circ\text{K} = \frac{1}{T_2} - 2.681 \times 10^{-3}/^\circ\text{K}$$

$$2.784 \times 10^{-3}/^\circ\text{K} = \frac{1}{T_2}$$

$$359.20^\circ\text{K} = T_2$$

The boiling point in $^\circ\text{C}$ is $359.20^\circ\text{K} - 273$ or 86.2°C .

• PROBLEM 266

Liquid nitrogen is an excellent bath for keeping temperatures around 77°K, its normal boiling point. What pressure would you need to maintain over the liquid nitrogen if you wanted to set the bath temperature at 85°K? Heat of vaporization is about 5560 J/mole.

Solution: One uses the Clausius-Clapeyron equation to solve for the final pressure. This equation is written

$$\ln \frac{P_1}{P_2} = \frac{\Delta H}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

where P_1 is the initial pressure, P_2 is the final pressure at 85°K, ΔH is the heat of vaporization, R is the gas constant, 8.314 J/mole-°K, T_1 is the initial temperature and T_2 is the final temperature. The pressure of a liquid at its boiling point is equal to the atmospheric pressure, which is 1 atm. Solving for the final pressure:

$$P_1 = 1 \text{ atm} \quad T_1 = 77^\circ\text{K}$$

$$P_2 = ? \quad T_2 = 85^\circ\text{K}$$

$$\Delta H = 5560 \text{ J/mole}$$

$$R = 8.314 \text{ J/mole } ^\circ\text{K}$$

$$\ln \frac{1}{P_2} = \frac{5560 \text{ J/mole}}{8.314 \text{ J/mole } ^\circ\text{K}} \left(\frac{1}{85^\circ\text{K}} - \frac{1}{77^\circ\text{K}} \right)$$

$$-\ln P_2 = 6.687 \times 10^2 ^\circ\text{K} (1.176 \times 10^{-2} ^\circ\text{K} - 1.298 \times 10^{-2} ^\circ\text{K})$$

$$-\ln P_2 = 6.687 \times 10^2 ^\circ\text{K} (-1.22 \times 10^{-3} / ^\circ\text{K})$$

$$-\ln P_2 = -8.1581 \times 10^{-1}$$

$$P_2 = 2.26 \text{ atm.}$$

• PROBLEM 267

If the $\Delta H'$ of water is 40,600 J/mole at 100°C , what is the boiling point of water at a pressure of .750 atm.

Solution: The boiling point is defined as the temperature at which the vapor pressure of a liquid equals that of the surrounding atmosphere. The normal or standard boiling point is that temperature where the vapor pressure of a gas equals one atmosphere. For water, this is 100°C .

In this problem, therefore, you have two pressures for water, one temperature, and $\Delta H'$, which equals the heat required to transform one mole of liquid to the ideal-gas state. These parameters are used to solve for the non-standard boiling point in the Clausius-Clapeyron equation, which states

$$\log \frac{P_1}{P_2} = \frac{\Delta H'}{19.15} \left(\frac{1}{T_2} - \frac{1}{T_1} \right),$$

where p_1 and p_2 are vapor pressures and T_1 and T_2 are their respective temperatures in degrees Kelvin. If you let T_2 be the boiling point of water at .750 atm, substituting into the equation:

$$\log \frac{1.00}{.750} = \frac{40,600}{19.15} \left(\frac{1}{T_2} - \frac{1}{373} \right)$$

Solving for T_2 , you obtain 365°K or 92°C , which is the boiling point of water at .750 atm.

• PROBLEM 268

If the vapor pressure of CCl_4 (carbon tetrachloride) is .132 atm at 23°C and .526 atm at 58°C , what is the $\Delta H'$ in this temperature range?

Solution: Molecules in a vapor exert a pressure that is characteristic of its liquid state. The pressure exerted by this vapor, when in equilibrium with the liquid, is called the equilibrium vapor pressure. The equilibrium vapor pressure can be expressed quantitatively as

$$\log p = \frac{-\Delta H'}{19.15 T} + C$$

where p = vapor pressure, T = temperature in degrees Kelvin, $\Delta H'$ = heat required to transform one mole of liquid to the ideal-gas state, and C = a constant that is characteristic of the liquid. C can be eliminated from the expression by denoting that at two temperature ranges, T_1 and T_2 , you have:

$$\log p_1 - \log p_2 = \frac{-\Delta H'}{19.15 T_1} + \frac{\Delta H'}{19.15 T_2} \quad \text{or}$$

$$\log \frac{p_1}{p_2} = \frac{\Delta H'}{19.15} \left(\frac{1}{T_2} - \frac{1}{T_1} \right).$$

In this problem, you are asked to find $\Delta H'$.

Let $p_1 = .132$ atm at $T_1 = 296^\circ\text{K}$ and $p_2 = .526$ atm at $T_2 = 331^\circ\text{K}$, then substitute these values in the above equation. Therefore,

$$\log \frac{.132}{.526} = \frac{\Delta H'}{19.15} \left(\frac{1}{331} - \frac{1}{296} \right)$$

Solving for $\Delta H'$, you obtain

$$\Delta H' = 32,000 \text{ J/mole.}$$

OSMOTIC PRESSURE

• PROBLEM 269

A sugar solution was prepared by dissolving 9.0 g of sugar in 500 g of water. At 27°C , the osmotic pressure was measured as 2.46 atm. Determine the molecular weight of the sugar.

Solution: The molecular weight of the sugar is found by determining the concentration, C , of sugar from the equation for osmotic pressure,

$$\pi = CRT,$$

where π is the osmotic pressure, R = universal gas constant = 0.08206 liter-atm/mole- $^\circ\text{K}$, and T is the absolute temperature.

The osmotic pressure is measured as $\pi = 2.46$ atm and the absolute temperature is $T = 27^\circ\text{C} + 273 = 300^\circ\text{K}$, hence

$$\pi = CRT,$$

$$2.46 \text{ atm} = C \times 0.08206 \text{ liter-atm/mole}^\circ\text{K} \times 300^\circ\text{K},$$

$$\text{or } C = \frac{2.46 \text{ atm}}{0.08206 \text{ liter-atm/mole}^\circ\text{K} \times 300^\circ\text{K}}$$

$$= 0.10 \text{ mole/liter.}$$

If we assume that the volume occupied by the sugar molecules in so small a concentration can be neglected, then in 1 liter of solution there is approximately 1 liter of water, or 1000 g of water, and

$$C = 0.10 \text{ mole/1000 g.}$$

Therefore, there is 0.10 mole of sugar dissolved in 1000 g of water.

9.0 g of sugar dissolved in 500 g of water is equivalent to 18.0 g of sugar dissolved in 1000 g of water (9.0 g/500 g = 18.0 g/1000 g). But since $C = 0.10 \text{ mole/1000 g}$, the 18.0 g of sugar must correspond to 0.10 mole of sugar. Therefore, the molecular weight of the sugar is

$$18 \text{ g}/0.1 \text{ mole} = 180 \text{ g/mole.}$$

• PROBLEM 270

A chemist dissolves 10 g of an unknown protein in a liter of water at 25°C. The osmotic pressure is found to be 9.25 mmHg. What is the protein's molecular weight. Based upon the number of moles in 10 g of protein, what would the freezing point depression and boiling point elevation be? Assume $R = \text{Universal Gas Constant} = .0821 \text{ liter-atm/mole}^\circ\text{K}$, $k_f = 1.86^\circ\text{C/m}$, and $k_b = .52^\circ\text{C/m}$.

Solution: The osmotic pressure (π) can be related to the molar concentration (C), Universal Gas Constant (R) and temperature (T) in degrees Kelvin of a solution via the formula $\pi = CRT$. After solving for C one can determine the molecular weight of the protein. Recall, $C = \text{moles/liter} = N/V$, where N = moles and V = volume in liters. Hence, $\pi = N/V RT$. But $N = \text{moles} = \text{No. of grams/molecular wt} = g/\text{M.W.}$ Rewriting,

$$\pi = \left(\frac{g}{\text{M.W.}} \right) RT \quad \text{or} \quad \text{M.W.} = \frac{gRT}{\pi V}$$

Substituting for the known values,

$$\text{M.W.} = \frac{(10 \text{ g})(.0821)(298^\circ\text{K})}{9.25 \text{ mmHg} (1)} \times \frac{760 \text{ mm}}{\text{atm}} = 20,100 \frac{\text{g}}{\text{mole.}}$$

Note, you assumed a volume of 1 liter and multiplied by the conversion factor 760 mm/atm, since R is in atm. and π in mmHg. Now that you know that the molecular weight is 20,100, the number of moles in the 10 g of protein is $10/20,100 = 4.98 \times 10^{-4} \text{ moles}$. To find the freezing point depression,

ΔT_f , and boiling point elevation, ΔT_b , you must find the molality, since $\Delta T_b = k_b m$ and $\Delta T_f = k_f m$, where k_f = molal freezing point depression constant, k_b = molal boiling point elevation constant and m = molality. Molality = moles of solute per 1 kg solvent. Thus, molality = .000498 mole/1 kg, since in 1 liter of water you have 1000 ml, which weighs 1 kg. Therefore,

$$\Delta T_f = (1.86^\circ\text{C})(.000498) = 9.3 \times 10^{-4} \text{ } ^\circ\text{C} \quad \text{and}$$

$$\Delta T_b = .52^\circ\text{C}(.000498) = 2.59 \times 10^{-4} \text{ } ^\circ\text{C}.$$

SURFACE TENSION

• PROBLEM 271

What would be the surface tension of a liquid (density 0.876 g/cm^3) which rises 2.0 cm in a capillary where water at 20°C rises 4.5 cm?

Solution: Surface tension refers specifically to the force within a liquid that acts parallel to the surface and tends to stretch the surface out. The equation that relates surface tension to the density of a liquid is

$$\gamma = 1/2 r h \rho g$$

where γ is the surface tension, r is the radius of the tube, h is the height the liquid rises, ρ is the density of the liquid and g is the downward acceleration of gravity.

This equation can be rewritten

$$r g = \frac{2\gamma}{h\rho}$$

Because r and g are constant for a given capillary tube

$$\frac{2\gamma_{\text{H}_2\text{O}}}{h_{\text{H}_2\text{O}} \rho_{\text{H}_2\text{O}}} = \frac{2\gamma_{\text{liq}}}{h_{\text{liq}} \rho_{\text{liq}}}$$

One can find $\gamma_{\text{H}_2\text{O}}$ and $\rho_{\text{H}_2\text{O}}$ from standard tables.

$$\gamma_{\text{H}_2\text{O}} = 72.62 \text{ dyn/cm}, \rho_{\text{H}_2\text{O}} = 1 \text{ g/cm}^3.$$

Solving for γ_{liq} :

$$\frac{2 (72.62 \text{ dyn/cm})}{(4.5 \text{ cm}) (1 \text{ g/cm}^3)} = \frac{2 \gamma_{\text{liq}}}{(2.0 \text{ cm}) (0.876 \text{ g/cm}^3)}$$

$$\gamma_{liq} = \frac{2(72.62 \text{ dyn/cm})(2.0 \text{ cm})(0.876 \text{ g/cm}^3)}{2(4.5 \text{ cm})(1 \text{ g/cm}^3)}$$

$$\gamma_{liq} = 28.27 \text{ dyn/cm}$$

• PROBLEM 272

Given that the surface tension of water is 72.62 dyn/cm at 20°C, how high should water rise in a capillary that is 1.0 mm in diameter?

Solution: The surface tension is related to the radius of the capillary tube by the equation:

$$\gamma = \frac{1}{2} r h \rho g$$

where γ is the surface tension, r is the radius of the capillary tube, h is the height the water rises, ρ is the density and g is the downward force of gravity. The density of H_2O is 1 g/cm³, the force of gravity is 980 cm/s² and the surface tension is given as 72.62 dyn/cm. The radius is found by dividing the diameter of the tube by 2. (1 mm = 10⁻¹ cm). Solving for h :

$$72.62 \frac{\text{dyn}}{\text{cm}} = 72.62 \frac{\text{g}}{\text{sec}^2}$$

$$72.62 \text{ g/sec}^2 = \frac{1}{2}(10^{-1} / 2 \text{ cm})(h)(1 \text{ g/cm}^3)(980 \text{ cm/sec}^2)$$

$$h = \frac{72.62 \text{ g/sec}^2 (2)(2)}{(10^{-1} \text{ cm})(1 \text{ g/cm}^3)(980 \text{ cm/sec}^2)}$$

$$h = 2.96 \text{ cm.}$$

CHAPTER 8



SOLUTION CHEMISTRY

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 251 to 275 for step-by-step solutions to problems.

It is important to understand the different ways in which the concentration of a solvent in a solution can be expressed. The different expressions of concentrations used in this chapter follow.

- a) Formality: the number of gram formula weights per liter of solution (abbreviated F)
- b) Molarity: the number of gram molecular weights per liter of solution (abbreviated M)
- c) Normality: the number of gram equivalent weights per liter of solution (abbreviated N)

On occasion, each of these measures may be referred to as 1000 g of solvent rather than one liter of solution. They are then called weight-formality, weight-molarity, and weight-normality respectively. The weight-molarity is the only one of the three in common usage and is also called molality. Therefore, a fourth definition should be added to the list.

- d) Molality: the number of gram molecular weights per 1000 g of solvent (abbreviated m)

The differences in (a), (b), and (c) are sufficiently subtle to justify additional explanation. The "formula weight" may differ from the "molecular weight" if hydration is present in the solute but, of course, loses its identity when dissolved in an aqueous solution. For example, the formula weight of

$\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ is 256.43 but, in an aqueous solution, the water of hydration loses its identity and only magnesium (Mg^{+2}) and nitrate (NO_3^-) ions can be traced to the solute. Formula weights and molecular weights also differ for complicated molecules, such as polymers, where the formula is known but the number of monomer units in an individual molecule, and hence the size of the molecule, are uncertain.

Normality is related to the reactivity of a species. In the example given above, if 76.9 g of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ is dissolved in water to form one liter of solution, the formality and molarity are both 0.3 (76.9/256.43). The salt completely ionizes to form magnesium (Mg^{+2}) ions and nitrate (NO_3^-) ions. The concentration of magnesium ions (Mg^{+2}) is 0.3M, and of nitrate ions (NO_3^-) is 0.6M. However, the normality of both the magnesium ions (Mg^{+2}) and the nitrate ions (NO_3^-) is 0.6N. For magnesium, the normality is twice the molarity because each ion has a charge of +2. While it is generally true, for ions, that the normality is equal to the molarity times the charge on the ion, it is important to examine the reaction context before specifying the normality.

In calculating the volumes of solutions required to mix a new solution of intermediate concentration or to neutralize species from separate solutions, it is important to calculate the amount (i.e., moles) of solute in each solution. In the problems in this text, non-ideal volume effects are, in general, neglected, and you may assume the volumes are additive. The procedure is best illustrated by an example.

Determine the volume of 0.1N H_2SO_4 and 0.1M H_2SO_4 needed to neutralize 100 ml of 0.5M NaOH. It is important to understand the meaning of molarity and normality.

In 100 ml of 0.5M NaOH, there exists 0.05 moles of OH^- ions. Therefore, 0.05 moles of H^+ ions will be required for neutralization. In 0.1M H_2SO_4 , there are 0.2 moles of H^+ per liter available for reaction, while in 0.1N H_2SO_4 , there are only 0.1 moles of H^+ per liter available for reaction. For the 0.1N solution, 500 ml (0.5 liters) will contain 0.05 moles of H^+ , while for the 0.1M (0.2N) solution, 250 ml (0.25 liters) will contain the required 0.05 moles of H^+ .

Step-by-Step Solutions to Problems in this Chapter, "Solution Chemistry"

DENSITY AND FORMALITY

• PROBLEM 273

In over 90% of cases, the concentration of ethanol (C_2H_5OH , density = 0.80 g/ml) in blood necessary to produce intoxication is 0.0030 g/ml. A concentration of 0.0070 g/ml is fatal. What volume of 80 proof (40% ethanol by volume) Scotch whiskey must an intoxicated person consume before the concentration of ethanol in his blood reaches a fatal level? Assume that all the alcohol goes directly to the blood and that the blood volume of a person is 7.0 liters.

Solution: The difference between a fatal concentration of ethanol and an intoxicating concentration of ethanol is $0.0070 \text{ g/ml} - 0.0030 \text{ g/ml} = 0.0040 \text{ g/ml}$. We must determine the total volume of ethanol in the blood which corresponds to a concentration of 0.0040 g/ml and then calculate the volume of Scotch whiskey which will provide this concentration.

The total mass of ethanol in blood needed to raise the concentration from an intoxicating to a fatal level is equal to concentration of ethanol \times volume of blood = $0.0040 \text{ g/ml} \times 7 \text{ liters} = 0.0040 \text{ g/ml} \times 7000 \text{ ml} = 28 \text{ g}$ of ethanol. Dividing this mass by the density, we obtain the corresponding volume of ethanol, or $28 \text{ g} / 0.80 \text{ g/ml} = 35 \text{ ml}$ of ethanol.

The amount of Scotch whiskey that must be consumed must provide 35 ml of ethanol. But the Scotch whiskey is only 40% ethanol, or 0.40 ml ethanol/ml Scotch. Let v denote the volume of Scotch in ml. Then

$$\left(\begin{array}{c} \text{ratio of} \\ \text{ethanol to Scotch} \end{array} \right) \times \text{volume of Scotch} = \text{volume of ethanol}$$

$$0.40 \text{ ml ethanol/ml Scotch} \times v \text{ ml Scotch} = 35 \text{ ml ethanol}$$

or, $v = \frac{35 \text{ ml ethanol}}{0.40 \text{ ml ethanol/ml Scotch}} = 88 \text{ ml Scotch.}$

Thus, under our assumptions, an intoxicated person must drink 88 ml of 80 proof Scotch whiskey (about 3 ounces) before ethanol reaches a fatal level in his blood.

• PROBLEM 274

What are the mole fractions of solute and solvent in a solution prepared by dissolving 98 g H_2SO_4 (M.W. 98) in 162 g H_2O (M.W. 18)?

Solution: The mole fraction of solute is defined as the moles of solute divided by the sum of the number of moles of the solute and the number of moles of the solvent.

$$\text{Mole fraction of solute} = \frac{\text{moles of solute}}{\text{moles of solute} + \text{moles of solvent}}$$

The mole fraction of the solvent is defined similarly.

$$\text{Mole fraction of solvent} = \frac{\text{moles of solvent}}{\text{moles of solute} + \text{moles of solvent}}$$

Here, the solute is H_2SO_4 and the solvent is H_2O . One is given the amount of H_2SO_4 and H_2O in grams. Therefore, these quantities must be converted to moles. This can be done by dividing the number of grams available by the molecular weight.

$$\text{No. of moles} = \frac{\text{no. of grams}}{\text{MW}}$$

$$\text{No. of moles of } \text{H}_2\text{SO}_4 = \frac{98 \text{ g}}{98 \text{ g/mole}} = 1 \text{ mole}$$

$$\text{No. of moles of } \text{H}_2\text{O} = \frac{162 \text{ g}}{18 \text{ g/mole}} = 9 \text{ moles}$$

Now that the number of moles of both solvent and solute are known, the mole fraction can be found.

$$\text{Mole fraction of } \text{H}_2\text{SO}_4 = \frac{1 \text{ mole}}{1 \text{ mole} + 9 \text{ moles}} = 0.1$$

$$\text{Mole fraction of } \text{H}_2\text{O} = \frac{9 \text{ moles}}{1 \text{ mole} + 9 \text{ moles}} = 0.9.$$

• PROBLEM 275

A wine has an acetic acid (CH_3COOH , 60 g/formula weight) content of 0.66% by weight. If the density of the wine is 1.11 g/ml, what is the formality of the acid?

Solution: This problem involves the correct interpretation of percent by weight. 0.66% by weight means 0.66×10^{-2} g acid per 100 g wine or 6.6×10^{-2} g acid per 1000 g wine. To convert 6.6×10^{-2} g acid to formula weight we divide by 60 g/formula weight and obtain 6.6×10^{-2} g/60 g/formula weight = 1.1×10^{-3} formula weight. To convert 1000 g wine to volume, we divide by the density of the wine and obtain 1000 g/1.11 g/ml = 900 ml = 0.90 liter. The formality is then

$$\begin{aligned}\text{formality} &= \frac{\text{formula weights}}{\text{volume (liters)}} = \frac{1.10 \times 10^{-3} \text{ formula weight}}{0.90 \text{ liter}} \\ &= 1.2 \times 10^{-3} \text{ formal} = 1.2 \times 10^{-3} \text{ F.}\end{aligned}$$

MOLALITY

• PROBLEM 276

2.3 g of ethanol ($\text{C}_2\text{H}_5\text{OH}$, molecular weight = 46 g/mole) is added to 500 g of water. Determine the molality of the resulting solution.

Solution: This problem is a calculation of the molality of an aqueous solution. Molality is equal to the number of moles of solute per kilogram of solvent. In this case the solvent is water and the solute is ethanol.

Since 1 kg of water corresponds to twice the amount of water given in the problem it is desirable to calculate the amount of ethanol that would be added to 1 kg of water and still maintain the same concentration of 2.3 g ethanol per 500 g of water. For $2 \times 500 \text{ g} = 1000 \text{ g} = 1 \text{ kg}$ of water, we require $2 \times 2.3 \text{ g} = 4.6 \text{ g}$ of ethanol.

We now know that our solution of 2.3 g ethanol in 500 g water corresponds to 4.6 g ethanol per 1 kg water, and all that remains to be done is to calculate the number of moles of ethanol in 4.6 g. We do this by dividing by the molecular weight to obtain $4.6 \text{ g}/46 \text{ g/mole} = 0.1 \text{ mole ethanol}$.

The concentration of the solution is then
0.1 mole ethanol/1 kg water = 0.1 molal.

What is the molality of a solution in which 49 g of H_2SO_4 (MW 98) is dissolved in 250 grams of water?

Solution: Molality is defined as the number of moles of solute per kilogram of solvent.

$$\text{Molality} = \frac{\text{moles of solute}}{\text{no. of kg. of solvent}}$$

Here, the solute is H_2SO_4 and the solvent is water. In this problem, one is given the number of grams of solute and the number of grams of solvent. One must calculate the number of moles of solute and the number of kilograms of solvent.

The number of moles of solute (H_2SO_4) is found by dividing the number of grams available by the molecular weight.

$$\text{no. of moles} = \frac{\text{no. of grams}}{\text{MW}}$$

$$\text{no. of moles of } \text{H}_2\text{SO}_4 = \frac{49 \text{ g}}{98 \text{ g/mole}} = 0.5 \text{ moles}$$

Grams can be converted to kilograms by multiplying the number of grams by the conversion factor 1 kg/1000 gm. For the water, $250 \times 1 \text{ kg}/1000 \text{ gm} = .250 \text{ kg}$. The molality can now be found.

$$\text{Molality} = \frac{\text{no. of moles of } \text{H}_2\text{SO}_4}{\text{no. of kg. of } \text{H}_2\text{O}}$$

$$\text{Molality} = \frac{0.5 \text{ moles}}{.250 \text{ kg}} = 2.0 \text{ moles/kg.}$$

Determine the mass of water to which 293 g of NaCl (formula weight = 58.5 g/mole) is added to obtain a 0.25 molal solution.

Solution: To solve this problem we use the relationship

$$\text{molality} = \frac{\text{number of moles of solute}}{\text{kg of solvent}} \quad \text{or,}$$

$$\text{molality} = \frac{\text{moles of NaCl}}{\text{kg of water}} .$$

The number of moles of NaCl in 293 g is determined by dividing 293 g by the formula weight of NaCl, or
 $\text{moles NaCl} = \frac{\text{mass NaCl}}{\text{formula weight NaCl}} = \frac{293 \text{ g}}{58.5 \text{ g/mole}} = 5.0 \text{ moles NaCl}$. Then, solving for the kg of water,

$$\text{kg of water} = \frac{\text{moles of NaCl}}{\text{molality}} = \frac{5.0 \text{ moles NaCl}}{0.25 \text{ molal}} = 20 \text{ kg H}_2\text{O}.$$

MOLARITY

• PROBLEM 279

Calculate the molarity of a solution containing 10.0 grams of sulfuric acid in 500 ml of solution. (MW of H_2SO_4 = 98.1.)

Solution: The molarity of a compound in a solution is defined as the number of moles of the compound in one liter of the solution. In this problem, one is told that there are 10.0 grams of H_2SO_4 present. One should first calculate the number of moles that 10.0 g represents. This can be done by dividing 10.0 g by the molecular weight of H_2SO_4 .

$$\text{number of moles} = \frac{\text{amount present in grams}}{\text{molecular weight}}$$

$$\text{number of moles of H}_2\text{SO}_4 = \frac{10.0 \text{ g}}{98.1 \text{ g/mole}} = 0.102 \text{ moles}$$

Since molarity is defined as the number of moles in one liter of solution, and since, one is told that there is 0.102 moles in 500 ml ($\frac{1}{2}$ of a liter), one should multiply the number of moles present by 2. This determines the number of moles in H_2SO_4 present in 1000 ml.

$$\text{Number of moles in 1000 ml} = 2 \times 0.102 = 0.204.$$

Because molarity is defined as the number of moles in 1 liter, the molarity (M) here is 0.204 M.

• PROBLEM 280

Calculate the weight in grams of sulfuric acid in 2.00 liters of 0.100 molar solution. (MW of H_2SO_4 = 98.1.)

Solution: Molarity is defined as the number of moles of a compound in a solution. In this problem, one is trying to

find the amount of H_2SO_4 in 2.00 liters of a solution that is 0.100 molar in H_2SO_4 . From the definition of molarity, one can see that in one liter of this solution there is 0.100 moles of H_2SO_4 . This means that in 2.00 liters of the same solution, there is twice that amount.

$$\begin{aligned}\text{number of moles of } \text{H}_2\text{SO}_4 \text{ in 2 liters} &= 2 \text{ liters} \times \\ &0.100 \text{ moles/liter} = 0.200 \text{ moles}\end{aligned}$$

Since one knows that the molecular weight of H_2SO_4 is 98.1 g/mole and that there are 0.200 moles present in this solution, one can find the number of grams present by multiplying the MW by the number of moles present.

$$\text{number of grams} = \text{MW} \times \text{number of moles present}$$

$$\begin{aligned}\text{number of grams of } \text{H}_2\text{SO}_4 &= 98.1 \text{ g/mole} \times 0.200 \text{ moles} \\ &= 19.6 \text{ g.}\end{aligned}$$

• PROBLEM 281

Hydrogen peroxide solution for hair bleaching is usually prepared by mixing 5.0 g of hydrogen peroxide (H_2O_2 , molecular weight = 34 g/mole) per 100 ml of solution. What is the molarity of this solution?

Solution: Before employing the definition

$$\text{molarity} = \frac{\text{number of moles}}{\text{volume (liters)}}$$

we must convert 5.0 g of H_2O_2 to the corresponding number of moles. To do this, we use the formula

$$\text{moles} = \text{mass/molecular weight.}$$

Then,

$$\begin{aligned}\text{moles} &= \frac{\text{mass}}{\text{molecular weight}} \\ &= \frac{5.0 \text{ g}}{34 \text{ g/mole}} = 0.15 \text{ moles } \text{H}_2\text{O}_2\end{aligned}$$

Converting 100 ml to liters, $100 \text{ ml} = 100 \text{ ml} \times \frac{1 \text{ liter}}{1000 \text{ ml}} = 0.10 \text{ liter}$. The molarity is then

$$\begin{aligned}\text{molarity} &= \frac{\text{number of moles}}{\text{volume (liters)}} = \frac{0.15 \text{ moles}}{0.10 \text{ liters}} \\ &= 1.5 \text{ moles/liter} = 1.5 \text{ molar} = 1.5 \text{ M.}\end{aligned}$$

• PROBLEM 282

Each liter of human blood serum contains about 3.4 g of sodium ions (Na^+ , ionic weight = 23 g/mole). What is the molarity of Na^+ in human blood serum?

Solution: We are given that the concentration of Na^+ is 3.4 g/liter. To convert this concentration to molarity, we need to convert 3.4 g to the corresponding number of moles. This is accomplished by dividing 3.4 g by the ionic weight of Na^+ (which is essentially the atomic weight, since the missing electron does not detract much from the weight) to obtain $3.4 \text{ g} / 23 \text{ g/mole} = 0.15 \text{ mole}$. The concentration is thus $0.15 \text{ mole/liter} = 0.15 \text{ molar} = 0.15 \text{ M}$.

• PROBLEM 283

A 10^6 liter tank of seawater contains 16,600 kg of chlorine (Cl^-), 9200 kg of sodium (Na^+) and 1180 kg of magnesium (Mg^{++}). Calculate the molarity of each. Is all the charge accounted for?

Solution: You are given the volume and weight of the materials in seawater and asked to calculate the molarity. Since molarity = no. of moles/liters, you need to calculate the number of moles of Cl^- , Mg^{++} and Na^+ present to solve this problem.

A mole = $\frac{\text{no. of grams}}{\text{atomic weight}}$. For Cl^- , then, the

$$\text{no. of moles} = \frac{1.66 \times 10^7 \text{ g}}{35.45 \text{ g/moles}} = 4.68 \times 10^5 \text{ moles}$$

$$\text{Therefore, its molarity is } \frac{4.68 \times 10^5}{10^6} = .468 \text{ M } \text{Cl}^-$$

$$\text{For } \text{Mg}^{++}, \text{ the no. of moles} = \frac{1.18 \times 10^6 \text{ g}}{24.3 \text{ m/g}}$$

$$\text{Its molarity is, thus, } \frac{\frac{1.18 \times 10^6 \text{ g}}{24.3 \text{ m/g}}}{10^6 \text{ liters}} = .048 \text{ M}$$

$$\text{For } \text{Na}^+, \text{ you have } \frac{9.200 \times 10^6 \text{ g}}{23 \text{ m/g}} = \text{no. of moles of } \text{Na}^+.$$

$$\text{Its molarity} = \frac{\frac{9.200 \times 10^6 \text{ g}}{23 \text{ m/g}}}{10^6} = .400 \text{ M}.$$

The total positive charge is $.400 + 2(.048) = .496$ M. The concentration of Mg^{++} must be doubled (in total positive charge) since each ion has 2 charges instead of one, as in Na^+ . The total negative charge stems only from Cl^- , which must equal $.468$ M. Thus, 0.496 moles/liter - $.468$ moles/liter = $.028$ moles/liter negative charge is unaccounted for, since all opposite charges must cancel each other out.

• PROBLEM 284

What mass of calcium bromide $CaBr_2$, is needed to prepare 150 ml of a 3.5 M solution? (M.W. of $CaBr_2 = 200.618$ g/mole)

Solution: M is the molarity of the solution and is defined as the number of moles of solute per liter. The solute is the substance being added to solution. A 1 M solution contains 1 mole of solute per liter (1000 ml) of solution.

We are asked to calculate the mass, however, no term for mass appears in the molarity equation. Therefore, a connection must be found between mass and another variable. The connection is the mole equation (moles = mass/M.W.). After substitution, the molarity equation reads

$$M = \frac{\frac{\text{grams (mass)}}{\text{M.W.}}}{\text{liters}}$$

$$\begin{aligned} \text{grams} &= \text{liters (M.W.) (M)} \\ &= .15\text{l (}200.618 \text{ g/mole) (}3.5 \text{ moles/l)} \\ &= 105.32 \text{ g of } CaBr_2. \end{aligned}$$

• PROBLEM 285

In the United States, alcohol is rated according to "proof", which is usually defined as twice the percent by volume of pure ethanol in solution, measured at 60°F. What is the molarity of ethanol in a "92 proof" solution of ethanol in water. The density of ethanol (C_2H_5OH , molecular weight = 46 g/mole) is 0.80 g/cm^3 at 60°F; the density of water (H_2O , molecular weight = 18 g/mole) is 1.0 gm/cm^3 at 60°F (15.5°C).

Solution: We will solve this problem by determining the number of moles of ethanol in one liter of solution.

A solution which is "92 proof" corresponds to 92/2 or 46% ethanol by volume. Consider one liter of solution. The volume of ethanol in this liter is $46\% \times 1 \text{ liter} = 0.46 \text{ liter}$. Converting this to cm^3 , $0.46 \text{ liter} = 0.46 \text{ liter} \times 1000 \text{ cm}^3/\text{liter} = 460 \text{ cm}^3$. To convert this to

the corresponding mass we multiply by the density (mass = volume \times density), obtaining $460 \text{ cm}^3 \times 0.80 \text{ g/cm}^3 = 460 \times 0.80 \text{ g}$ of ethanol. The number of moles in our one liter solution is obtained by dividing the number of grams by the molecular weight (moles = mass/molecular weight), or, moles = $460 \times 0.80 \text{ g}/46 \text{ g/mole} = 8.0 \text{ moles}$.

There are 8.0 moles of ethanol in one liter of solution, and the concentration is therefore

$$8.0 \text{ moles/l liter} = 8.0 \text{ molar.}$$

• PROBLEM 286

A student has 50.00 mg crystal of $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ (M.W. = 315) and wants to make a solution of .12 M OH^- . How much water must the student add to obtain such a solution?

Solution: Molarity is defined as moles/liter. You are given the desired molarity, .12 M, and can calculate the number of moles of OH^- that would have to be present for this molarity. With these two known values, the volume (liters) can be obtained. You have 50 mg of $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ or (1 g/1000mg) 50 mg/315 g/mole = 1.59×10^{-4} moles of $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$.

Now, $\text{Ba}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$ contains 2 moles of OH^- , which means that you have 3.18×10^{-4} moles of OH^- present in this crystal. Recalling, the stated definition of molarity, liters (volume) = moles/molarity. Therefore, the volume (liter) of water to be added =

$$= \frac{3.18 \times 10^{-4} \text{ moles}}{.120 \text{ moles/liter}} = 2.65 \times 10^{-3} \text{ liters}$$

or 2.65 ml total solution volume.

• PROBLEM 287

Determine the molarity of a 40.0% solution of HCl which has a density of 1.20 g/ml.

Solution: The molarity is defined as the number of moles of a compound present in 1 liter of solution. Here one is told that the density of this solution is 1.20 g/ml. This means that one ml of the solution weighs 1.20 g. The solution is 40.0% HCl, thus 40.0% of the 1.20 g is made up by HCl and 60% by H_2O . Here one finds the molarity of the HCl by: 1) determining the total weight of 1 liter of the solution, 2) calculating the weight of HCl present, and 3) finding the number of moles of HCl present in 1 liter (molarity).

Solving: 1) If 1 ml of this solution weight 1.20 g, 1000 ml (1 liter) is equal to 1000×1.20 g.

weight of solution = density \times 1000 ml

$$= 1.20 \text{ g/ml} \times 1000 \text{ ml} = 1200 \text{ g}$$

2) 40.0% of this weight is taken up by HCl.

weight of HCl = weight of solution \times .40

$$= 1200 \text{ g} \times .40 = 480 \text{ g.}$$

3) The molecular weight of HCl is 36.5 g/mole, thus the number of moles of HCl present in this 1 liter of solution is equal to the weight of the HCl present divided by its molecular weight.

no. of moles of HCl in 1 liter = $\frac{\text{weight of HCl}}{\text{MW}}$

$$= \frac{480 \text{ g}}{36.5 \text{ g/mole}} = 13.15 \text{ moles}$$

The solution is therefore 13.15 moles/liter in HCl or 13.15 M.

• PROBLEM 288

If 25 ml of .11 M sucrose is added to 10 ml of .52 M sucrose, what would be the resulting molarity?

Solution: In this problem one mixes two solutions of different concentrations and wishes to find the resulting molarity. The answer is obtained by first calculating the total number of moles present and then dividing this by the total volume.

Molarity is defined as moles \div volume. Therefore,

$$\text{moles} = \text{Molarity} \times \text{Volume}$$

For the .11 M solution,

$$\text{no. of moles} = .11 \text{ moles/liter} \times .025 \text{ liter}$$

$$= .00275 \text{ moles}$$

For the .52 M solution,

$$\text{no. of moles} = .52 \text{ moles/liter} \times .010 \text{ liter}$$

$$= .00520 \text{ moles}$$

$$\text{total no. of moles} = .00275 + .00520 = .00795$$

$$\text{total no. of liters} = .025 + .010 = .035$$

$$\text{Final Molarity} = \frac{.00795}{.035} = .23 \text{ M.}$$

• PROBLEM 289

A chemist wants to dilute 50 ml of 3.50 M H_2SO_4 to 2.00 M H_2SO_4 . To what volume must it be diluted?

Solution: Molarity is defined as the number of moles of solute per liter of solution. In other words,

$$\text{molarity} = \frac{\text{no. of moles of solute}}{\text{liters of solution}} .$$

In this problem you have a 50 ml solution of 3.50 M. In a liter there are 1000 ml. Thus, the number of liters in this solution is .05. Substituting, you have

$$3.50 \text{ M} = \frac{\text{no. of moles of } \text{H}_2\text{SO}_4}{.05 \text{ L}}$$

Solving for the number of moles, you obtain .175 moles of H_2SO_4 .

When you dilute this mixture, you will still have .175 moles of the solute, H_2SO_4 . You are only increasing the volume of the solvent, water. Therefore, in the diluted 2.0 M solution you wish to have

$$2.00 = \frac{.175}{\text{liters of solvent}} .$$

Solving for liters of solvent, you obtain .0875 liters or 87.5 ml. Thus, to dilute to a molarity of 2, the total volume must be 87.5 ml.

• PROBLEM 290

What will be the final concentration of a solution of 50 ml of .5M H_2SO_4 added to 75 ml of .25M H_2SO_4 (the final volume is 125 ml)?

Solution: Two facts must be known to answer this question:

(1) M = Molarity = the number of moles of solute per liter of solution and (2) that the number of moles of H_2SO_4 in the final solution is the sum of the number of moles of H_2SO_4 in each solution.

$$\text{molarity} = \frac{\text{no. of moles}}{\text{liters of solution}}$$

One can rearrange this equation to determine the number of moles present in each solution.

$$\text{no. of moles} = \text{molarity} \times \text{liters of solution}$$

The total number of moles in the final solution is equal to the sum of the number of moles of each of the two initial solutions.

$$\begin{aligned}\text{total number of moles} &= (0.05 \text{ l})(.5 \text{ moles/l}) + \\ &\quad (.075 \text{ l})(.25 \text{ moles/l}) \\ &= 0.025 \text{ moles} + .019 \text{ moles} \\ &= .044 \text{ moles.}\end{aligned}$$

To obtain the final concentration, you return to the definition of molarity, supplying the known values. Thus, molarity (final concentration) =


$$\frac{0.044 \text{ moles}}{.125 \text{ liter}} = .35 \text{ M H}_2\text{SO}_4$$

(It was given that the volume of the final solution equaled 125 ml = .125 liters.)

• PROBLEM 291

What will be the final concentration of I^- in a solution made by mixing 35.0 ml of 0.100 M NaHSO_3 with 10.0 ml of 0.100 M KIO_3 ? Assume volumes are additive.

Solution: The equation for this reaction is



I^- is formed in the following reaction:



One must first calculate the concentration of I_2 formed in reaction 1 before the amount of I^- can be found. There are 5 moles of HSO_3^- and 2 moles of IO_3^- needed to form 1 mole of I_2 . Molarity is defined as moles per liter. Thus the solution of NaHSO_3 contains

$$35 \text{ ml} \times \frac{0.100 \text{ moles}}{1000 \text{ ml}} = 3.50 \times 10^{-3} \text{ moles}$$

$$\text{There are } 10.0 \text{ ml} \times \frac{.100 \text{ moles}}{1000 \text{ ml}} = 1.00 \times 10^{-3} \text{ moles}$$

of KIO_3 present. In any given reaction one of the reactants is called the limiting reagent. This means that

when the reaction proceeds, all of the limiting reagent is used but there are other reactants left in their original state. The limiting reagent limits the amount of product formed. Two moles of IO_3^- are needed to form 1 mole of I_2 . Thus, if all of the KIO_3 present reacts

$$\frac{1.00 \times 10^{-3} \text{ moles}}{2} \quad \text{or } 5.00 \times 10^{-4} \text{ moles of } \text{I}_2$$

will be formed. 5 moles of NaHSO_3 are needed to form 1 mole of I_2 , if all of the NaHSO_3 present reacts,

$$\frac{3.50 \times 10^{-3} \text{ moles}}{5} \quad \text{or } 7.0 \times 10^{-4} \text{ moles of } \text{I}_2$$

will be formed. Therefore, when all of the KIO_3 reacts, only 5.0×10^{-4} moles of I_2 are formed and when all of the NaHSO_3 is reacted, 7.0×10^{-4} moles of I_2 are formed. Thus, KIO_3 is the limiting reagent and only 5.00×10^{-4} moles of I_2 are formed. When 5.00×10^{-4} moles of I_2 are formed, $5 \times 5.0 \times 10^{-4}$ moles, or 2.5×10^{-3} moles of NaHSO_3 are used. This leaves $(3.5 \times 10^{-3} - 2.5 \times 10^{-3})$ moles or 1.0×10^{-3} moles of NaHSO_3 left unreacted to react to form the I^- in equation 2. H_2O , HSO_3^- and I_2 react to form I^- . The number of moles present of each of these must be known to find the limiting reagent. This is done in a manner similar to that used for reaction 1. From equation 1, one knows that there is 1 mole of H_2O formed for each I_2 formed. Thus, there are 5.0×10^{-4} moles of H_2O present.

From equation 2 one notices that one mole of each H_2O , HSO_3^- , and I_2 are needed to form 2 moles of I^- . There are 5.0×10^{-4} moles of H_2O , 1.0×10^{-3} moles of HSO_3^- , and 5.0×10^{-4} moles of I_2 present. Thus, the I_2 is the limiting reagent because there is the least of it present to react. Therefore, $2 \times 5.0 \times 10^{-4}$ moles or 1.0×10^{-3} moles of I^- formed. If the volumes are additive, the final volume present is 45 ml. Thus, there are 1.0×10^{-3} moles of I^- present in the final volume of 45 ml of solution. Solving for the concentration

$$\frac{1.0 \times 10^{-3} \text{ moles}}{.045 \text{ liter}} = .002 \text{ M.}$$

• PROBLEM 292

What is the mole fraction of H_2SO_4 in a 7.0 molar solution of H_2SO_4 which has a density of 1.39 g/ml?

Solution: The mole fraction of H_2SO_4 is equal to the number of moles of H_2SO_4 divided by the sum of the number of moles of H_2SO_4 and of H_2O .

$$\text{mole fraction of } \text{H}_2\text{SO}_4 = \frac{\text{moles of } \text{H}_2\text{SO}_4}{\text{moles of } \text{H}_2\text{SO}_4 + \text{moles of } \text{H}_2\text{O}}$$

Since the solution is 7.0 molar in H_2SO_4 , you have 7 moles of H_2SO_4 per liter of solution. If one knows how much 7 moles of H_2SO_4 weighs, and how much one liter of the solution weighs, the weight of water can be determined by taking the difference between the quantities. The weight of one liter of the solution can be calculated by multiplying the density by the conversion factor 1000 ml/liter.

$$\begin{aligned} \text{weight of 1 liter of the solution} \\ = 1.39 \text{ g/ml} \times 1000 \text{ ml/liter} = 1390 \text{ g/l.} \end{aligned}$$

$$\begin{aligned} \text{weight of the water in a one liter solution} \\ = \text{weight of the total solution} \\ - \text{weight of 7 moles of } \text{H}_2\text{SO}_4 \end{aligned}$$

7 moles of H_2SO_4 weigh 7 times the molecular weight of H_2SO_4 .

$$\text{weight of 7 moles of } \text{H}_2\text{SO}_4 = 7 \times 98 \text{ g} = 686 \text{ g.}$$

$$\text{Hence, weight of the water} = 1390 \text{ g} - 686 \text{ g} = 704 \text{ g.}$$

The number of moles of H_2O is found by dividing its weight by the molecular weight of H_2O .

$$\text{no. of moles of } \text{H}_2\text{O} = \frac{704 \text{ g}}{18 \text{ g/mol}} = 39 \text{ moles}$$

The number of moles of both components of the system is now known, therefore, the mole fraction of H_2SO_4 can be obtained.

$$\text{mole fraction } \text{H}_2\text{SO}_4 = \frac{\text{moles of } \text{H}_2\text{SO}_4}{\text{moles of } \text{H}_2\text{SO}_4 + \text{moles of } \text{H}_2\text{O}}$$

$$\text{mole fraction of } \text{H}_2\text{SO}_4 = \frac{7}{7 + 39} = 0.15.$$

• PROBLEM 293

You have 100-proof (50 percent alcohol by volume) bonded Scotch whisky. Calculate its molarity, mole fraction, and molality. If the temperature were to drop to -10°C , could you still drink the Scotch? Assume density = .79 g/ml for ethyl alcohol ($\text{C}_2\text{H}_5\text{OH}$) and $K_f = 1.86^\circ\text{C/m}$ for water.

Solution: To find molarity, mole fraction, and molality, you need to know how many moles of alcohol are present in the whiskey. You can obtain this from a calculation using the density.

To find whether you can still drink the whisky at

- 10°C, requires you to know its freezing point, since you cannot drink something if it is frozen. You can determine the freezing point of the Scotch by determining the freezing point depression of the water containing the alcohol. This is found from $\Delta T_f = K_f m$, where ΔT_f = the depression, K_f = freezing point depression constant, and m = molality.

You now proceed as follows:

You are told that 100 proof means 50 percent alcohol by volume. This means that 1 liter of components possess 500 cm³ of ethyl alcohol (a liter = 1000 cm³). Density = mass/volume. You know the density and volume, so that mass of alcohol = density \times volume = .79(500) = 395 g.

Molecular wt. of the alcohol = 46.07 g. Thus, you have 395/46.07 = 8.57 moles of alcohol. Hence,

$$\text{molarity} = \frac{\text{moles of solute}}{\text{liters of solution}} = \frac{8.57}{1} = 8.57 \text{ M.}$$

$$\text{Molality} = \frac{\text{moles of solute}}{\text{kilograms of solvent}} .$$

To find the number of grams of solvent, you need to use the density again. In 1 liter of Scotch Whisky, you have 500 cm³ of water. Density of water = 1 g/cm³.

Thus, mass of water = 500 cm³(1 g/cm³) = 500 grams. Thus,

$$\text{Molality} = \frac{8.57}{\frac{500}{1000}} = 17.14 \text{ m}$$

You divided 500 by 1000 since molality is moles per 1000 g of solvent.

Mole fraction: (MW of H₂O = 18.) moles of H₂O = 500 g/18.0 g/mole.

Mole fraction of ethyl alcohol =

$$\frac{\text{moles ethyl alcohol}}{\text{moles ethyl alcohol} + \text{moles H}_2\text{O}} = \frac{8.57}{8.57 + \frac{500}{18}} = .2357.$$

Now, you answer whether the scotch can be consumed at - 10°C. You already calculated the molality. Thus, $\Delta T_f = K_f m = (1.86)(17.14) = 31.88^\circ\text{C}$. Thus, the freezing point of whisky is 0°C - 31.88°C = - 31.88°C. The temperature is only - 10°C, however. This means whisky is still a liquid and can be consumed.

NORMALITY

• PROBLEM 294

Calculate the normality of a solution containing 2.45 g of sulfuric acid in 2.00 liters of solution. (MW of $\text{H}_2\text{SO}_4 = 98.1$.)


Solution: Normality is defined by the following equation.

$$\text{Normality} = \frac{\text{grams of solute}}{\text{equivalent weight} \times \text{liters of solution}}$$

In this problem, one is given the grams of solute (H_2SO_4) present and the number of liters of solution it is dissolved in. One equivalent of a substance is the weight in which the acid contains one gram atom of replaceable hydrogen. This means that when the acid is dissolved in a solution, and it ionizes, that in one equivalent weight, one hydrogen atom is released. The equivalent weight for acids is defined as:

$$\text{equivalent weight} = \frac{\text{MW}}{\text{no. of replaceable H}}$$

When H_2SO_4 is dissolved in a solution, there are two replaceable H as shown in the following equation:



This means that in calculating the equivalent weight for H_2SO_4 , the molecular weight is divided by 2.

$$\begin{aligned}\text{equivalent weight of } \text{H}_2\text{SO}_4 &= \frac{98.1 \text{ g}}{2 \text{ equivalents}} \\ &= 49 \text{ g/equiv}\end{aligned}$$

The normality can now be calculated.

$$\text{Normality} = \frac{\text{grams of solute}}{\text{equiv. wt} \times \text{liters of soln}}$$

$$\begin{aligned}\text{Normality} &= \frac{2.45 \text{ g}}{49 \text{ g/equiv} \times 2.0 \text{ liters}} \\ &= 0.025 \text{ equiv/liter.}\end{aligned}$$

• PROBLEM 295

How many grams of sulfuric acid are contained in 3.00 liters of 0.500 N solution? (MW of H_2SO_4 = 98.1.)


Solution: The number of grams of sulfuric acid contained in this solution can be determined by using the definition of normality.

$$\text{Normality} = \frac{\text{grams of solute}}{\text{equivalent weight} \times \text{liters of solution}}$$

Here, one is given the normality (N) and the number of liters of solution. The equivalent weight for an acid is the molecular weight divided by the number of replaceable hydrogens,

$$\text{equivalent weight} = \frac{\text{MW}}{\text{no. of replaceable H}}$$

The number of replaceable hydrogens is determined by the number of hydrogens that will ionize when the acid is placed in solution. This number is 2 for H_2SO_4 , as shown in the following equation:



Once the number of replaceable hydrogen is known, the equivalent weight can be found.

$$\text{equivalent weight} = \frac{\text{MW}}{\text{no. of replaceable H}}$$

$$\text{equivalent weight of } \text{H}_2\text{SO}_4 = \frac{98.1 \text{ g}}{2 \text{ equiv}} = 49 \text{ g/equiv.}$$

At this point the number of grams of solute can be determined.


$$\text{grams of solute} = \text{Normality} \times \text{equiv weight} \times \text{liters of solution}$$

$$\begin{aligned} \text{grams of } \text{H}_2\text{SO}_4 &= 0.500 \text{ equiv/l} \times 49 \text{ g/equiv} \times 3.0 \text{ liter} \\ &= 73.5 \text{ g.} \end{aligned}$$

• PROBLEM 296

Calculate the normality of a solution 40 ml of which is required to neutralize 0.56 g of KOH.

Solution: KOH ionizes as shown in this equation.



This means that for each KOH molecule ionized, one OH^- ion is formed. Thus to neutralize the KOH one must have 1 H^+ ion for each OH^- ion. Thus, one must first determine the number of moles of KOH present in 0.56 g. This is done by dividing 0.56 g by the molecular weight of KOH (MW of KOH = 56)

number of moles of KOH = number of moles of OH^-

$$\text{number of moles of KOH} = \frac{0.56 \text{ g}}{\text{MW}} = \frac{0.56 \text{ g}}{56 \text{ g/mole}} = .01 \text{ mole}$$

Therefore, in the 40 ml of the solution there must be .01 moles of H^+ ions. The normality of an acid is defined as the number of equivalents of H^+ in 1 liter of solution. An equivalent may be defined as the weight of acid or base that produces 1 mole of H^+ or OH^- ions. In this problem, equivalents = moles. There are 1000 ml in 1 liter, thus in 40 ml there are

$$40 \text{ ml} \times \frac{1 \text{ liter}}{1000 \text{ ml}}$$

$$\text{number of liters} = 40 \text{ ml} \times \frac{1 \text{ liter}}{1000 \text{ ml}} = .04 \text{ liters.}$$

The normality can now be found.

$$\begin{aligned} \text{normality} &= \frac{\text{number of equivalents}}{\text{number of liters}} \\ &= \frac{.01 \text{ equivalents}}{.04 \text{ liters}} = .25 \text{ N.} \end{aligned}$$

• PROBLEM 297

If 20 ml of 0.5 N salt solution is diluted to 1 liter, what is the new concentration?

Solution: When considering normality one must always keep in mind that it is a concentration defined as the number of equivalents per liter. Since the number of equivalents does not change during dilution, equivalents before dilution = equivalents after dilution, or in other words,

$$N_1 V_1 = N_2 V_2 \qquad 1 \text{ liter} = 1000 \text{ ml}$$

where N_1 is the normality of the initial solution, V_1 the initial volume, N_2 the final normality and V_2 is final volume.

Solving for the final normality:

$$(0.5)(0.020\text{L}) = N_2 (1\text{L})$$

$$N_2 = \frac{(0.5)(0.020\%)}{1\text{ L}}$$

$$= 0.01\text{ N}$$

• PROBLEM 298

A sulfuric acid solution has a density of 1.8 g/ml and is 90% H_2SO_4 by weight. What weight of H_2SO_4 is present in 1000 ml of the solution? What is the molarity of the solution? the normality?

Solution: 1) Here one is asked to find the weight of H_2SO_4 in 1000 ml of the solution and told that the solution weighs 1.8 g/ml and that 90% of this weight is made up by H_2SO_4 . The total weight of the solution is 1000 ml times the weight of one ml.

Hence, weight of solution = 1000 ml \times 1.8 g/ml = 1800 g. H_2SO_4 makes up 90% of this weight.

$$\text{weight of } \text{H}_2\text{SO}_4 = .90 \times \text{weight of solution}$$


$$= .90 \times 1800\text{ g} = 1620\text{ g.}$$

2) The molarity is defined as the number of moles in one liter of solution. One has already found that 1620 g of H_2SO_4 is present in 1 liter of the solution. Thus, to calculate molarity, one should determine the number of moles present in 1620 g. This is done by dividing 1620 g by the molecular weight of H_2SO_4 . (MW of H_2SO_4 = 98.1.)

$$\text{moles of } \text{H}_2\text{SO}_4 \text{ present} = \frac{1620\text{ g}}{98.1\text{ g/mole}} = 16.5\text{ moles}$$

Since there is 1 liter of solution, the molarity of H_2SO_4 is 16.5 M.

3) The normality is defined as the number of moles of ionizable hydrogens per liter of solution. From the following equation one can see that there are 2 ionizable hydrogens for each molecule of H_2SO_4 .



Therefore if there are 16.5 moles of H_2SO_4 in one liter of the solution the normality is twice this amount.

$$\text{normality of an acid} = \text{no. of ionizable H} \times \text{molarity}$$

$$= 2 \times 16.5 = 33.0\text{ N.}$$

A solution is prepared by dissolving 464 g NaOH in water and then diluting to one liter. The density of the resulting solution is 1.37 g/ml. Express the concentration of NaOH as (a) percentage by weight, (b) molarity, (c) normality, (d) molality, (e) mole fraction.

Solution: (a) The percentage by weight of NaOH in this solution is found by dividing the weight of NaOH present, 464 g, by the weight of the solution and multiplying by 100. The weight of the solution is found by using the density of the solution. The density, 1.37 g/ml, tells the weight of 1 ml of solution, namely, 1.37 g. In one liter, there are 1000 ml, thus the weight of the solution is 1000 times the weight of one ml.

$$\text{weight of 1 liter} = 1000 \text{ ml} \times 1.37 \text{ g/ml} = 1370 \text{ g.}$$

The percentage by weight of the NaOH in the solution can now be found.


$$\begin{aligned} \text{percentage of NaOH} &= \frac{\text{weight of NaOH}}{\text{weight of solution}} \times 100 \\ &= \frac{464 \text{ g}}{1370 \text{ g}} \times 100 = 33.9 \%. \end{aligned}$$

(b) The molarity is defined as the number of moles in one liter of solution. The molarity in this case can be found by determining the number of moles in 464 g of NaOH, which is the amount of NaOH in one liter. The number of moles can be found by dividing 464 g by the molecular weight of NaOH. (MW of NaOH = 40.)

$$\text{no. of moles} = \frac{464 \text{ g}}{\text{MW}} = \frac{464 \text{ g}}{40 \text{ g/mole}} = 11.6 \text{ moles.}$$

The molarity of this solution is thus 11.6 M.

(c) The normality of a basic solution is the number of moles of ionizable OH^- ions in one liter of solution. There is one ionizable OH^- ion in each NaOH as shown by the equation:



Therefore, there are the same number of OH^- ions in the solution as NaOH molecules dissolved. Thus, the molarity equals the normality.

$$\text{normality} = \text{molarity} \times 1 \text{ ionizable OH}^-$$

$$\text{normality} = 11.6 \text{ N}$$

(d) The molality is defined as the number of moles present in 1 kg of solvent. One has already found that the solution weighs 1370 g or 1.37 kg and that there are 11.6 moles of NaOH present. Therefore the molality can now be found.

$$\begin{aligned}\text{molality} &= \frac{\text{no. of moles present}}{\text{no. of kg present}} \\ &= \frac{11.6 \text{ moles}}{1.37 \text{ kg} - .339(1.370 \text{ kg})} = 12.8 \text{ m}\end{aligned}$$

(e) The mole fraction is equal to the number of moles of each component divided by the total number of moles in the system. The components in this system are H_2O and NaOH. One already has found that there are 11.6 moles of NaOH present, but not the number of moles of H_2O . This can be found by determining the weight of the water and dividing it by its molecular weight. This solution weighs 1370 g, the NaOH weighs 464 g, thus the weight of the water is equal to the difference of these two figures,

$$\begin{aligned}\text{weight of } \text{H}_2\text{O} &= \text{weight of solution} - \text{weight of NaOH} \\ &= 1370 \text{ g} - 464 \text{ g} = 906 \text{ g}.\end{aligned}$$

One can now find the number of moles of H_2O present. (MW of H_2O = 18.)

$$\text{no. of moles} = \frac{906 \text{ g}}{18 \text{ g/mole}} = 50.3 \text{ moles}$$

The total number of moles in the system is the sum of the number of moles of H_2O and of the NaOH.

$$\begin{aligned}\text{no. of moles in system} &= \text{moles } \text{H}_2\text{O} + \text{moles NaOH} \\ &= 50.3 + 11.6 = 61.9 \text{ moles}\end{aligned}$$

One can now find the mole fractions.

$$\text{mole fraction} = \frac{\text{no. of moles of each component}}{\text{total no. of moles in system}}$$

$$\text{mole fraction of } \text{H}_2\text{O} = \frac{50.3}{61.9} = .81$$

$$\text{mole fraction of NaOH} = \frac{11.6}{61.9} = .19.$$

NEUTRALIZATION

• PROBLEM 300

A mixture consisting only of KOH and $\text{Ca}(\text{OH})_2$ is neutralized by acid. If it takes exactly 0.100 equivalents to neutralize 4.221 g of the mixture, what must have been its initial composition by weight?

Solution: One equivalent of an acid is the mass of acid required to furnish one mole of H_3O^+ ; one equivalent of a base is the mass of base required to furnish one mole of OH^- or to accept one mole of H_3O^+ . Here one must find the mixture of KOH and $\text{Ca}(\text{OH})_2$ that contains 0.100 equivalent of base. There is one equivalent for each OH^- in a molecule of base. Thus, there is 1 equivalent of base for each KOH. There are two for each $\text{Ca}(\text{OH})_2$. Since one is given that there are 0.100 equivalent in 4.221 g of the mixture, one should solve for the equivalent per gram for KOH and $\text{Ca}(\text{OH})_2$.

for KOH: MW = 56.1 equiv = 1

$$\frac{\text{equiv}}{\text{g}} = \frac{1}{56.1} = 1.78 \times 10^{-2} \text{ equiv/g}$$

For $\text{Ca}(\text{OH})_2$ MW = 74.06 equiv = 2

$$\frac{\text{equiv}}{\text{g}} = \frac{2}{74.06} = 2.70 \times 10^{-2} \text{ equiv/g.}$$

There must be 0.100 equiv present. Let

x = number of grams of KOH

$4.221 - x$ = number of grams of $\text{Ca}(\text{OH})_2$

$$0.100\text{equiv.} = \left(1.78 \times 10^{-2} \frac{\text{equiv}}{\text{g}}\right)(x) + \left(2.70 \times 10^{-2} \frac{\text{equiv}}{\text{g}}\right)(4.221 - x)$$

$$0.100\text{equiv.} = 0.0178x \frac{\text{equiv}}{\text{g}} + .114 \text{ equiv} - .0270x \frac{\text{equiv}}{\text{g}}$$

$$0.100\text{equiv.} = - .0092x \frac{\text{equiv}}{\text{g}} + .114 \text{ equiv}$$

$$-.014\text{equiv.} = - .0092x \frac{\text{equiv}}{\text{g}}$$

$$\frac{.014 \text{ equiv}}{.0092 \text{ equiv/g}} = x$$

$$1.52 \text{ g} = x$$


$$2.701 \text{ g} = 4.221 - x$$

The original mixture contains 1.52 g KOH and 2.701 g Ca(OH)_2 .

• PROBLEM 301

Calculate the volume of 0.3 N base necessary to neutralize 3 liters of 0.01 N nitric acid.

Solution: For neutralization to occur, there must be the same number of hydrogen ions as there are hydroxide ions. This is shown by the following equation



The H^+ ions come from the acid, the OH^- ions from the base. The number of H^+ and OH^- ions are equal to the number of equivalents.


Normality is defined as the number of equivalents of acid or base per liter of solution.

$$\text{Normality} = \frac{\text{equivalents}}{\text{liter}}$$

An equivalent is the number of grams of the acid or base multiplied by the number of replaceable hydrogens or hydroxides divided by the molecular weight of the acid or base.

$$\text{equivalent} = \frac{\text{grams of solute} \times \text{no. of replaceable H or OH}}{\text{MW of solute}}$$

The number of replaceable hydrogens or hydroxides is defined as the number which ionize when the compound is placed in solution. For nitric acid, there is one replaceable hydrogen



In brief, for neutralization there must be the same equivalent amount of base as there is acid. Because normality is defined as

$$\text{Normality} = \frac{\text{equivalents}}{\text{liters}}$$

equivalents are equal to the normality times the volume.

Thus, the normality of the base times its volume equals the normality of the acid times its volume.

$$N_{\text{base}} V_{\text{base}} = N_{\text{acid}} V_{\text{acid}}$$

where N is the normality and V is the volume.

$$\text{Here, } 0.3 \text{ equiv/liter} \times V_{\text{base}} = 0.01 \text{ equiv/liter} \times 3 \text{ liters}$$

$$V_{\text{base}} = \frac{0.01 \times 3 \text{ liters}}{0.3}$$

$$= 0.1 \text{ liter or } 100 \text{ mL}$$

• PROBLEM 302

What weight of $\text{Ca}(\text{OH})_2$, calcium hydroxide is needed to neutralize 28.0 g of HCl . What weight of phosphoric acid would this weight of $\text{Ca}(\text{OH})_2$ neutralize?

Solution: When we have neutralization problems, i.e., those where an acid reacts with a base, we must consider the meaning of equivalents. Equivalent weight may be defined as the molecular weight of a substance (grams/mole) divided by the number of protons (H^+) or hydroxyl (OH^-) ions available for reaction. In other words,

$$\text{Equivalent weight} = \frac{\text{molecular weight}}{\text{number of } \text{H}^+ \text{ or } \text{OH}^-}$$

In a neutralization reaction, the number of equivalents of acid equals the number of equivalents of base.

The number of equivalents of acid and base are equal, thus, if we can calculate the number of equivalents of 1 reactant, we automatically know the number of equivalents present of the other reactant. To find this quantity use the equation

$$\text{no. of equivalents} = \frac{\text{grams}}{\text{grams/equivalent}}$$

The equivalent weight of HCl is

$$\frac{\text{M.W.}}{\text{no. of } \text{H}^+(\text{OH}^-)} = \frac{36.5}{1} = 36.5 \quad \text{and } \text{Ca}(\text{OH})_2 \text{ is } \frac{74.1}{2} =$$

37 g/equiv.

$$\text{no. of equivalents HCl present} = \frac{28 \text{ g}}{36.5 \text{ g/equiv}} = .767$$

Therefore, there are also .767 equivalents of $\text{Ca}(\text{OH})_2$. To find the grams of $\text{Ca}(\text{OH})_2$ needed use

$$\text{no. of equiv} = \frac{\text{grams Ca(OH)}_2}{\text{grams/equiv}}$$

$$.767 = \frac{\text{grams}}{37}$$

grams Ca(OH)_2 = 28.4 need to neutralize 28 g of HCl .

The second half of the problem is answered in exactly the same way. We know then that there must also be .767 equiv of H_3PO_4 (phosphoric acid) present due to the fact .767 equiv of base Ca(OH)_2 is present. Recall, equivalents of an acid must equal the equivalents of a base for neutralization to occur. The M.W. of H_3PO_4 = 98.0 g, which yields a value of $\frac{98}{3} = 32.7$ grams per equiv. Therefore,

$$\text{No. of equiv} = \frac{\text{grams H}_3\text{PO}_4}{\text{grams/equiv}}$$

$$.767 = \frac{\text{grams}}{32.7}$$

grams H_3PO_4 = 25.1 g needed to neutralize 28.4 g of Ca(OH)_2 .

CHAPTER 9

EQUILIBRIUM

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 277 to 303 for step-by-step solutions to problems.

Most physical and chemical phenomena encountered in chemistry (e.g., reactions, solubility, ionization, and dissociation) proceed to some equilibrium state where concentrations and other properties do not change with time. The equilibrium state, which might well appear to be a state where nothing is happening, is a dynamic state where reactions or other changes occur in opposite directions at equal rates.

The key to working all equilibrium problems lies in the definition of the equilibrium constant and conversion of this definition to an equation with only one unknown variable. For the general reaction in Equation 9-1



(which can represent a classical reaction, ionization, solubility, dissociation, or any other equilibrium phenomenon) the equilibrium constant is defined as follows:

$$K_{eq} = (a_R^r a_S^s) / (a_B^b a_D^d) \quad 9-2$$

where a_i^i is the activity of the species, i , raised to the stoichiometric coefficient i . For gases at low pressures, the activity is equal to the partial pressure in atmospheres; for liquids and solutes in dilute solution, the activity is equal to the concentration in moles/liter. The activity of pure species is one.

In order to solve for the equilibrium composition, it is necessary to convert the activity (the partial pressure or concentration) of each of the species to an expression in a single variable. This can almost always be done by referring

to the stoichiometry of the reaction and writing the concentration of each species as a function of the fractional or absolute conversion of one species. An example will best illustrate the technique.

Consider the reaction in Problem 314:



Since all species in this reaction are gases at moderate temperatures, the equilibrium constant is

$$K_{eq} = p_{\text{HI}}^2 / (p_{\text{H}_2} p_{\text{I}_2}) = n_{\text{HI}}^2 / (n_{\text{H}_2} n_{\text{I}_2}) (P/n_T)^0 \quad 9-4$$

where p represents the partial pressure in atm, P the total pressure, and n the number of moles. (Note that for gases at low pressures, $p_i = (n_i/n_T)P$). In order to solve Equation 9-4 for the equilibrium composition, n_{HI} , n_{H_2} , n_{I_2} , and n_T must be written in terms of a single variable. A convenient variable is x , the fraction of the initial H_2 consumed at equilibrium. From Equation 9-3, the stoichiometric equation representing the reaction, the amounts of each species can be written in terms of the variable x .

$$\begin{aligned} n_{\text{H}_2} &= 1.0(1-x) \\ n_{\text{I}_2} &= 1.0(1-x) \\ n_{\text{HI}} &= 1.0(2x) \\ n_T &= 2 \end{aligned}$$

Equation 9-4 can now be written in terms of a single variable, x , and solved as follows;

$$K = (2x)^2 / (1-x)^2 \quad 9-5$$

$$K^{1/2} = 2x / (1-x) \quad 9-6$$

$$x = K^{1/2} / (2 + K^{1/2}) \quad 9-7$$

The equilibrium constant varies with temperature (to be discussed in Chapter 15), but is independent of pressure. However, the equilibrium composition will be changed by changes in the pressure if the number of moles changes during the reaction. In Equation 9-4, the zero exponent on pressure is a consequence of the fact that there is no change in the number of moles in the reactants and products. If the number of moles changes, the exponent on total pressure will be non-zero and the composition will change as pressure changes.

The Principle of LeChatelier provides a way to determine quickly the qualitative effect of temperature and pressure changes (or changes in other important parameters) on equilibrium composition. LaChatelier's principle, simply stated, says that "*a system will respond to offset any imposed change.*" Therefore, if the pressure is increased, the equilibrium composition will shift in a way that will lower the pressure. In gas phase reactions, this is equivalent to shifting in a way to reduce the total number of moles. If the moles of products of a reaction exceed the moles of the reactants, the equilibrium composition will shift toward the reactants and vice versa. If there is no change in the number of moles in a reaction (as in the HI synthesis represented by Equation 9-4), changes in the pressure will not affect the equilibrium composition (hence the zero exponent on total pressure).

Similarly, if the temperature is increased, the equilibrium will shift in a way to reduce the temperature. This means, for increases in temperature, the equilibrium composition will shift toward the reactants for exothermic reactions and toward the products for endothermic reactions. The reverse is true for decreases in temperature (Exothermic reactions produce heat, while endothermic reactions consume heat).

Step-by-Step Solutions to Problems in this Chapter, "Equilibrium"

THE EQUILIBRIUM CONSTANT

• PROBLEM 303

Determine the equilibrium constant for the reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ if the equilibrium concentrations are: H_2 , 0.9 moles/liter; I_2 , 0.4 mole/liter; HI , 0.6 mole/liter.

Solution: The equilibrium constant (K_{eq}) is defined as the product of the concentrations of the products divided by the product of the concentrations of the reactants. These concentrations are brought to the power of the stoichiometric coefficient of that component. For example for the reaction $2\text{A} + \text{B} \rightleftharpoons 3\text{C}$, the equilibrium constant can be expressed:

$$K_{eq} = \frac{[\text{C}]^3}{[\text{A}]^2 [\text{B}]}$$

where $[\]$ indicates concentration.

One can now express the equilibrium constant for the reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ as

$$K_{eq} = \frac{[\text{HI}]^2}{[\text{H}_2] [\text{I}_2]}$$

One is given the concentrations of the components for the system, thus one can solve for the equilibrium constant.

$$K_{eq} = \frac{[\text{HI}]^2}{[\text{H}_2] [\text{I}_2]}$$


$[\text{H}_2] = 0.9 \text{ mole/liter}$
 $[\text{I}_2] = 0.4 \text{ mole/liter}$
 $[\text{HI}] = 0.6 \text{ mole/liter}$

$$K_{eq} = \frac{[0.6 \text{ mole/liter}]^2}{[0.9 \text{ moles/liter}][0.4 \text{ moles/liter}]}$$

$$= \frac{0.36}{0.36} = 1.0$$


• PROBLEM 304

One of the two most important uses of ammonia is as a reagent in the first step of the Ostwald process, a synthetic route for the production of nitric acid. This first step proceeds according to the equation



What is the expression for the equilibrium constant of this reaction?

Solution: This problem is an exercise in writing the equilibrium constant of a reaction. In general, for a reaction in which reactants A, B, C, ... go to products W, X, Y, ... according to the equation



where a, b, c, w, x, y, \dots are the stoichiometric coefficients, the equilibrium constant is given by

$$K = \frac{[\text{W}]^w [\text{X}]^x [\text{Y}]^y \dots}{[\text{A}]^a [\text{B}]^b [\text{C}]^c \dots}$$

Hence, for the reaction



the equilibrium constant is given by

$$K = \frac{[\text{NO}]^4 [\text{H}_2\text{O}]^6}{[\text{NH}_3]^4 [\text{O}_2]^5}$$

• PROBLEM 305

Given the reaction $\text{A} + \text{B} \rightleftharpoons \text{C} + \text{D}$, find the equilibrium constant for this reaction if .7 moles of C are formed when 1 mole of A and 1 mole of B are initially present. For this same equilibrium, find the equilibrium composition when 1 mole each of A, B, C, and D are initially present.

Solution: In general, an equilibrium constant measures the ratio of the concentrations of products to reactants, each raised to the power of their respective coefficients in the chemical equation. Thus, for this reaction, the equilibrium constant, K, is equal to

$$K = \frac{[\text{C}][\text{D}]}{[\text{A}][\text{B}]}$$

You are asked to find K in the first part of this problem. Therefore,

you must evaluate this expression. To do this, use stoichiometry. From the equation for the chemical reaction, you see that all the quantities are in equimolar amounts; 1:1:1:1. It is given that .7 moles of C is produced. This means, therefore, that .7 mole of D must also be produced. If the equimolar quantities are to be maintained, then, .7 mole of each A and B must have been consumed. If you started with 1 mole of each, then .3 mole must be left. These mole amounts are the concentrations if you assume they are in a given amount of volume. Thus, substituting these values in

$$K = \frac{[C][D]}{[A][B]}$$

you obtain

$$K = \frac{(.7)^2}{(.3)^2} = 5.44$$

Therefore, you have calculated the equilibrium constant of this reaction to be 5.44.

The second part of this problem asks you to use this same equilibrium for a reaction that starts with 1 mole each of A, B, C, and D. To find its composition, determine the concentration of each species at equilibrium. This can be found from

$$K = \frac{[C][D]}{[A][B]} = 5.44$$

To find the composition, assume X moles/liter (concentration) react. Thus, at equilibrium, both C and D have a final concentration of $1 + x$, since you started with 1 mole/liter and had x moles/liter of each produced. This means, therefore, that A's and B's initial concentration must be reduced by x moles/liter to $1 - x$. Substituting,

$$5.44 = \frac{(1+x)^2}{(1-x)^2}$$

Solving,

$$x = .40 \text{ moles/liter}$$

Thus, the concentration of both A and B = $1 - .4 = .6M$ and of both C and D = $1 + .4 = 1.4M$.

• PROBLEM 306

At a certain temperature, K_{eq} for the reaction $3C_2H_2 \rightleftharpoons C_6H_6$ is 4. If the equilibrium concentration of C_2H_2 is 0.5 mole/liter, what is the concentration of C_6H_6 ?

Solution: The equilibrium constant (K_{eq}) for this reaction is stated:

$$K_{eq} = \frac{[C_6H_6]}{[C_2H_2]^3}$$

where [] indicate concentration. The $[C_2H_2]$ is brought to the third power because three moles of it react. Equilibrium is defined as the point where no more product is formed and no more reactant is dissipated; thus their concentrations remain constant. Here, one is given K_{eq} and $[C_2H_2]$ and asked to find $[C_6H_6]$. This can be

done by substituting the given into the equation for the equilibrium constant.


$$K_{eq} = \frac{[C_6H_6]}{[C_2H_2]^3} \quad K_{eq} = 4$$

$$4 = \frac{[C_6H_6]}{(0.5)^3} \quad [C_2H_2] = 0.5 \text{ moles/liter}$$

$$[C_6H_6] = (0.5)^3 \times 4 = 0.5 \text{ moles/liter}$$

• PROBLEM 307

The following reaction



was allowed to proceed to equilibrium. The contents of the two-liter reaction vessel were then subjected to analysis and found to contain 1.0 mole H_2S , 0.20 mole H_2 , and 0.80 mole S_2 . What is the equilibrium constant K for this reaction?

Solution: This problem involves substitution into the equilibrium constant expression for this reaction,

$$K = \frac{[H_2]^2 [S_2]}{[H_2S]^2}$$

The equilibrium concentration of the reactant and products are $[H_2S] = 1.0 \text{ mole}/2 \text{ liters} = 0.50M$, $[H_2] = 0.20 \text{ mole}/2 \text{ liters} = 0.10M$, and $[S_2] = 0.80 \text{ mole}/2 \text{ liters} = 0.40M$. Hence, the value of the equilibrium constant is

$$K = \frac{[H_2]^2 [S_2]}{[H_2S]^2} = \frac{(0.10)^2 (0.40)}{(0.50)^2} = 0.016$$

for this reaction.

• PROBLEM 308

You have the reaction $A_2 + B_2 \rightleftharpoons 2AB$ with AB initially at 5 liters, $27^\circ C$, and 25 atm. $K = 50$. Find (a) the initial concentration of AB , (b) the A_2 and B_2 concentrations at equilibrium and (c) the partial pressures of A_2 and B_2 and AB at equilibrium.

Solution: Use the equation of state, which indicates $PV = NRT$, where P = pressure, V = volume, N = moles, R = universal gas constant ($0.0821 \text{ liter-atm/mole-K}$), and T = temperature in degrees kelvin (Celsius plus 273°). You can use this equation to find N , the moles

of AB initially present.

$$(a) \quad PV = NRT \text{ or } N = \frac{PV}{RT} = \frac{(25)(5)}{(.0821)(300)} = 5.08 \text{ moles.}$$

Thus, the initial concentration = moles/liters = $5.08/5 = 1.02$.

$$(b) \quad K = 50 \text{ (given). By definition, } K = \frac{[AB]^2}{[A_2][B_2]}. \text{ At equilibrium,}$$

let x = moles/liter of AB that are dissociated. If its initial concentration is 1.02M, then the concentration is $(1.02-x)$ M at equilibrium. From the reaction, 1 mole each of A_2 and B_2 is produced by every 2 moles of AB which react. Thus, if x moles/liter of AB dissociate, $\frac{1}{2}x$ moles/liter of both A_2 and B_2 are produced. Recalling,

$$K = \frac{[AB]^2}{[A_2][B_2]} = 50, \text{ you now substitute to obtain } \frac{(1.02-x)^2}{(.5x)^2} = 50.$$

Solving for x , $x = .225$ mole/liter. Therefore, at equilibrium, $[A_2] = [B_2] = x/2 = .1125$ mole/liter and $[AB] = 1.02 - .225 = .795$ mole/liter.

(c) The partial pressures of each will be $P = \frac{N}{V} RT$ from the equation of state, where $\frac{N}{V}$ is the concentration at equilibrium. Thus,


$$P_{A_2} = .1125(.0821)(300) = 2.77 \text{ atm, } P_{A_2} = P_{B_2}, \text{ since these concentrations are the same at equilibrium,}$$

$$P_{AB} = .795(.0821)(300) = 19.5 \text{ atm.}$$

EQUILIBRIUM CALCULATIONS

• PROBLEM 309

At 986°C, you have the following equilibrium:



Initially, 49.3 mole percent CO_2 is mixed with 50.7 mole percent H_2 . At equilibrium, you find 21.4 mole percent CO_2 , 22.8 mole percent H_2 , and 27.9 mole percent of CO and H_2O . Find K . If you start with a mole percent ratio of 60:40, CO_2 to H_2 , find the equilibrium concentrations of both reactants and products.

Solution: An equilibrium constant K_{eq} measures the ratio of the concentrations of products to reactants, each raised to the power of their respective coefficients in the chemical equation. Thus, K_{eq} for this reaction =

$$\frac{[CO][H_2O]}{[CO_2][H_2]}.$$

If you assume each substance occupies the same volume in liters, the

concentration can be expressed in moles because concentration = moles/liter, i.e., liters cancel out of the equilibrium constant expression. Thus, to find K , you need to find the number of moles of each of the products and reactants and then to substitute into the equilibrium expression. You are told the final product mole percents. The reactants, then, at equilibrium, have mole percents that equal their initial amounts minus the amount that decomposed to produce the products.

Thus, at equilibrium, $[CO_2] = 49.3 - 21.4 = 27.9$, which was given. Similarly $[H_2] = 50.7 - 22.8 = 27.9$, which was given. Thus, by substitution into

$$K = \frac{[CO][H_2O]}{[CO_2][H_2]}, \text{ you obtain } K = \frac{(27.9)(27.9)}{(21.4)(22.8)} = 1.60,$$

which is the equilibrium constant for this reaction.

The second part follows. You begin with a 60:40 ratio of $CO_2:H_2$, which means that initially you have .600 moles CO_2 and .400 moles H_2 . To find the equilibrium concentrations, let x = moles of CO formed. Thus, x = moles H_2O formed since coefficients tell us they are formed in equimolar amounts. The fact that moles of a product form, means that x moles of a reactant must have decomposed. Thus, at equilibrium, you have $0.600-x$ moles of CO_2 and $0.400-x$ moles of H_2 . Recalling,

$$K = \frac{[CO][H_2O]}{[CO_2][H_2]},$$

you can now substitute these values to give $K = \frac{x^2}{(0.6-x)(0.4-x)}$. From previous part, $K = 1.60$, therefore,

$$1.60 = \frac{x^2}{(0.6-x)(0.4-x)}.$$

Solving, $x = 0.267$. Thus, $[CO] = [H_2O] = x = 0.267M$, $[CO_2] = 0.6-x = 0.333M$ and $[H_2] = 0.4-x = 0.133M$.

• PROBLEM 310

A chemist mixes nitric oxide and bromine to form nitrosyl bromide at $298^\circ K$, according to the equation $2NO(g) + Br_2(g) \rightleftharpoons 2NOBr(g)$. Assuming $K = 100$, what is the quantity of nitrosyl bromide formed, if the reactants are at an initial pressure of 1 atm? $R = 0.0821 \text{ liter-atm./mole}^\circ K$.

Solution: You are given the equilibrium constant for this reaction and asked to calculate the quantity of nitrosyl bromide produced. The first step is to write out the equilibrium expression and equate it with the given value. For the general reaction,

$$x A + y B \rightleftharpoons z C, \quad K \text{ is defined } \frac{[C]^z}{[B]^y [A]^x},$$

where the brackets represent concentrations. For this reaction,

$$K = \frac{[\text{NOBr}]^2}{[\text{NO}]^2 [\text{Br}_2]} = 100.$$

To find out how much NOBr is produced, you would have to know how many moles of NO and Br₂ were reacted. Once this is known, you can find the number of grams produced. You know that the equilibrium expression is based on concentration of reactants and products. Concentration is expressed in moles per liter. This means that if the volume of the NOBr and its concentration is known, you can find moles, since concentration x volume (liters) = moles. Let us represent the concentration as

$$\frac{N}{V} = \frac{\text{moles}}{\text{Volume}}.$$

Thus, the equilibrium expression becomes

$$K = \frac{(N_{\text{NOBr}}/V)^2}{(N_{\text{NO}}/V)^2 (N_{\text{Br}_2}/V)}.$$

Let x = moles of NOBr formed. Then, x moles of NO and x/2 moles of Br₂ are consumed, since the coefficients of the reaction show a 2:2:1 ratio among NOBr:NO:Br₂. The equilibrium expression becomes

$$100 = \frac{N_{\text{NOBr}}^2 V}{N_{\text{NO}}^2 N_{\text{Br}_2}} = \frac{x^2 V}{(2-x)^2 (1-.5x)}.$$

If x moles of NOBr form, and you started with 2 moles of NO, then, at equilibrium, you have left 2-x moles of NO. You started with only 1 mole of Br₂ and 1/2x moles of it form NOBr; thus you have 1-.5x moles left. Therefore, you need to determine only the volume to find the quantity NOBr formed. V can be found from the equation of state, PV = NRT, where P = pressure, V = volume, N = moles, R = universal gas constant, and T = temperature in kelvin (celsius plus 273°). You are told that the reactants are under a pressure of 1 atm. at 298°K. N = 3, since the coefficients inform you that a relative sum of 3 moles of reagents exist. You know R. Thus,

$$V = \frac{NRT}{P} = \frac{(3)(.0821)(298)}{1} = 73.4 \text{ liters}.$$


Now that V is known, the equilibrium expression becomes

$$\frac{x^2 (73.4)}{(2-x)^2 (1-0.5x)} = 100.$$

Solving for x, you obtain x = .923 moles = moles of NOBr formed. Molecular weight = 110. Grams produced = .923 x 110 = 101.53g.

• PROBLEM 311

Carbon dioxide is reduced by hydrogen according to the equation



One mole of CO₂ is reacted with one mole of H₂ at 825K and it

is found that, at equilibrium, 0.27 atm of CO is present. Determine the equilibrium constant, K , at 825K.

Solution: This problem makes use of the fact that since there are equal numbers of moles of gaseous products and reactants, the partial pressure of each component is equal to the number of moles present at equilibrium.

For each mole of CO_2 and each mole of H_2 reacted, one mole of CO and one mole of H_2O are produced. Thus, if at equilibrium there is 0.27 atm = 0.27 mole of CO , then there is 0.27 mole of H_2O , $1 - 0.27 = 0.73$ mole of CO_2 , and $1 - 0.27 = 0.73$ mole of H_2 . If we let v denote the volume (in liters) in which the reaction takes place, then:


$$[\text{CO}] = [\text{H}_2\text{O}] = 0.27 \text{ mole}/v, [\text{CO}_2] = [\text{H}_2] = 0.73 \text{ mole}/v.$$

Substituting these values into the expression for the equilibrium constant gives

$$K = \frac{[\text{CO}][\text{H}_2\text{O}]}{[\text{CO}_2][\text{H}_2]} = \frac{(0.27/v)(0.27/v)}{(0.73/v)(0.73/v)} = \frac{(0.27)^2}{(0.73)^2} = 0.137.$$

• PROBLEM 312

Two moles of gaseous NH_3 are introduced into a 1.0-liter vessel and allowed to undergo partial decomposition at high temperature according to the reaction



At equilibrium, 1.0 mole of $\text{NH}_3(\text{g})$ remains. What is the value of the equilibrium constant?

Solution: This problem involves substitution into the expression for the equilibrium constant for this reaction,

$$K = \frac{[\text{N}_2][\text{H}_2]^3}{[\text{NH}_3]^2}$$


Since 1.0 mole of NH_3 remains, $2.0 - 1.0 = 1.0$ mole of NH_3 was consumed. Also, since one mole of N_2 and three moles of H_2 are formed per 2 moles NH_3 consumed, at equilibrium, there are $3/2$ moles of H_2 and $1/2$ mole of N_2 . The equilibrium concentrations are therefore $[\text{NH}_3] = 1.0 \text{ mole}/1.0 \text{ liter} = 1.0\text{M}$, $[\text{N}_2] = (1/2 \text{ mole})/(1.0 \text{ liter}) = 0.5\text{M}$, and $[\text{H}_2] = (3/2 \text{ moles})/(1.0 \text{ liter}) = 1.5\text{M}$.

Substituting these into the expression for K gives

$$K = \frac{[N_2][H_2]^3}{[NH_3]^2} = \frac{(0.5)(1.5)^3}{(1)^2} = 1.6875$$

• PROBLEM 313

Four moles of PCl_5 are placed in a 2-liter flask. When the following equilibrium is established



the flask is found to contain 0.8 mole of Cl_2 . What is the equilibrium constant?

Solution: The equilibrium constant (K_{eq}) for this reaction is:


$$K_{eq} = \frac{[PCl_3][Cl_2]}{[PCl_5]}$$

where [] indicate concentrations. One is given that, at equilibrium, there are 0.8 mole of Cl_2 present. From the equation for the reaction, it follows that there is 0.8 mole of PCl_3 present also. This means that 0.8 moles of PCl_5 has reacted. Originally 4 moles of PCl_5 were present, thus, at equilibrium, $4.0 - 0.8$ moles or 3.2 moles are present. These components are present in a two-liter flask. Therefore, the concentrations will be expressed in moles/2 liters. One can now solve for the equilibrium constant.

$$\begin{aligned} K_{eq} &= \frac{[PCl_3][Cl_2]}{[PCl_5]} & [PCl_3] &= 0.8 \text{ moles/2liters} \\ & & [Cl_2] &= 0.8 \text{ moles/2liters} \\ & & [PCl_5] &= 3.2 \text{ moles/2liters} \\ K_{eq} &= \frac{(0.8/2)(0.8/2)}{(3.2/2)} = \frac{(0.4M)(0.4M)}{(1.6M)} \\ &= \frac{0.16M^2}{1.6M} = 0.1M \end{aligned}$$

• PROBLEM 314

For the formation of hydrogen iodide,



the value of the equilibrium constant at 700K is 54.7. If one mole of H_2 and one mole of I_2 are the only materials initially present, what will be the equilibrium concentrations of H_2 , I_2 , and HI .

Solution: This problem is an application of the expression for the

equilibrium constant

$$K = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$$

except that the concentrations are unknown. However, as we shall see, we do not need to know the concentrations.

Let x be the number of moles of H_2 that disappeared at equilibrium. Since one mole of I_2 is consumed and two moles of HI are formed per mole of H_2 consumed, then, at equilibrium, x moles of I_2 will have reacted, $2x$ moles of HI will have formed, and $1-x$ moles of H_2 and $1-x$ moles of I_2 will remain. If we let the volume of the vessel in which the reaction takes place be v , then $[\text{H}_2] = (1-x)/v$, $[\text{I}_2] = (1-x)/v$, and $[\text{HI}] = 2x/v$. The expression for the equilibrium constant is then

$$K = 54.7 = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]} = \frac{(2x/v)^2}{[(1-x)/v][(1-x)/v]} = \frac{4x^2}{(1-x)^2} \cdot \frac{v^2}{v^2} \\ = \frac{4x^2}{1-2x+x^2}$$

or

$$54.7 - 2(54.7)x + 54.7x^2 = 4x^2$$

$$50.7x^2 - 109.4x + 54.7 = 0$$

Using the quadratic formula,

$$x = \frac{109.4 \pm \sqrt{(-109.4)^2 - 4(50.7)(54.7)}}{2(50.7)}$$

or

$$x = 0.79, x = 1.37$$

Since x , the number of moles of H_2 or I_2 that reacted, cannot be greater than one, the answer $x = 1.37$ must be a nonphysical entity, and we retain only the answer $x = 0.79$. Hence, at equilibrium,


$$[\text{H}_2] = 1 - x = 1 - 0.79 = 0.21 \text{ mole}$$

$$[\text{I}_2] = 1 - x = 1 - 0.79 = 0.21 \text{ mole}$$

$$[\text{HI}] = 2x = 2(0.79) = 1.58 \text{ moles.}$$

• PROBLEM 315

For the reaction



the value of the equilibrium constant at 825°K is 0.137. If 5.0 moles of CO_2 , 5.0 moles of H_2 , 1.0 mole of CO , and 1.0 mole of H_2O are initially present, what is the composition of the equilibrium mixture?

Solution: This problem is an application of the expression for the equilibrium constant.

From the stoichiometry of the reaction, one mole of CO and one mole of H_2O are produced for one mole of CO_2 and one mole of H_2 that are reacted. Hence, if x moles of CO are produced at equilibrium, then x moles of H_2O are produced, x moles of CO_2 are consumed, and x moles of H_2 are consumed. Therefore, at equilibrium, there are $1+x$ moles of CO , $1+x$ moles of H_2O , $5-x$ moles of CO_2 , and $5-x$ moles of H_2 . If we let v denote the volume of the reaction vessel, the equilibrium concentrations are

$$[\text{CO}] = [\text{H}_2\text{O}] = (1+x)/v, \text{ and } [\text{CO}_2] = [\text{H}_2] = (5-x)/v.$$

Substituting these values into the expression for the equilibrium constant gives

$$K = 0.137 = \frac{[\text{CO}][\text{H}_2\text{O}]}{[\text{CO}_2][\text{H}_2]} = \frac{[(1+x)/v][(1+x)/v]}{[(5-x)/v][(5-x)/v]} = \frac{(1+x)^2 v^2}{(5-x)^2 v^2} = \frac{1+2x+x^2}{25-10x+x^2},$$

or

$$0.137(25) - 0.137(10)x + 0.137x^2 = 1+2x+x^2$$

$$0.863x^2 + 3.370x - 2.425 = 0.$$

Using the quadratic equation,

$$x = \frac{-3.370 \pm \sqrt{(3.370)^2 - 4(0.863)(-2.425)}}{2(0.863)}$$

or

$$x = 0.62, \quad x = -4.52.$$

The second of these is nonphysical and is therefore discarded.

The equilibrium concentrations are then

$$[\text{CO}] = 1+x = 1+0.62 = 1.62 \text{ moles}$$

$$[\text{H}_2\text{O}] = 1+x = 1+0.62 = 1.62 \text{ moles}$$

$$[\text{CO}_2] = 5-x = 5-0.62 = 4.38 \text{ moles}$$

$$[\text{H}_2] = 5-x = 5-0.62 = 4.38 \text{ moles.}$$

• PROBLEM 316

One mole of H_2 is to be reacted with one mole of I_2 . Assuming the equilibrium constant is 45.9, what will be the final concentrations of the chemical components in a 1 liter box at 490°C ?

Solution: The first thing to do with this type of problem is to write a balanced equation for the reaction. A reaction between H_2 and I_2 is one which produces HI . As such, you have $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightarrow 2\text{HI}(\text{g})$. You are given the equilibrium constant of this reaction. To use this information, you must know the meaning of such a constant. Given the general reaction: $x\text{A} + y\text{B} \rightleftharpoons z\text{C}$, the equilibrium constant, K , is defined as

$$\frac{[\text{C}]^z}{[\text{B}]^y [\text{A}]^x},$$

where the brackets serve to indicate concentrations. For this problem,

$$K = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]} .$$

You are asked to find the concentrations. Let x = the no. of moles of H_2 that react. It follows, then, that x is also equal to the number of moles of I_2 that react and $2x$ is equal to the number of moles of HI that form. Substituting you obtain

$$\frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]} = 45.9 .$$

You started with one mole of each H_2 and I_2 . Therefore, at equilibrium you must have $1-x$ moles of each species. Since the volume of the box is one liter, this means that the concentration of each substance is equal to the number of moles of each substance. You can now substitute these values in the equilibrium equation.

$$\frac{(2x)^2}{(1-x)(1-x)} = 45.9 .$$

Solving for x using the quadratic equation, you obtain $x = .772$. Therefore, at equilibrium you have


$$[\text{H}_2] = 1 - x = .228 \text{ mol/liter}$$

$$[\text{I}_2] = 1 - x = .228 \text{ mol/liter}$$

$$[\text{HI}] = 2x = 1.554 \text{ mol/liter} .$$

• PROBLEM 317

For the reaction



the value of the equilibrium constant at 700K is 0.0183. If 3.0 moles of HI are placed in a 5-liter vessel and allowed to decompose according to the above equation, what percentage of the original HI would remain undissociated at equilibrium?

Solution: This problem is an application of the equilibrium expression

$$K = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2} .$$

Since two moles of HI are involved in the production of one mole of H_2 and one mole of I_2 , if x moles of H_2 (and therefore x moles of I_2) are present at equilibrium, then $2x$ moles of HI have been consumed and $3-2x$ moles of HI remain. Therefore, at equilibrium, $[\text{H}_2] = [\text{I}_2] = x \text{ moles} / 5 \text{ liters}$ and $[\text{HI}] =$

$(3-2x) \text{ moles} / 5 \text{ liters} .$

Substituting these into the expression for K gives

$$K = 0.0183 = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2} = \frac{(x/5)(x/5)}{[(3-2x)/5]^2} = \frac{x^2}{(3-2x)^2} \times \frac{5^2}{5^2}$$

$$= \frac{x^2}{(3-2x)^2} = \frac{x^2}{9-12x+4x^2}$$

or

$$9(0.0183) - 12(0.0183)x + 4(0.0183)x^2 = x^2$$

$$0.9268x^2 + 0.2196x - 0.1647 = 0$$

Using the quadratic formula,

$$x = \frac{-0.2196 \pm \sqrt{(0.2196)^2 - 4(0.9268)(-0.1647)}}{2(0.9268)}$$

or

$$x = -0.56, \quad x = 0.32$$

Since negative moles are a nonphysical entity, the first answer is discarded and the second retained.

Thus, at equilibrium,

$$[H_2] = x = 0.32 \text{ mole}$$

$$[I_2] = x = 0.32 \text{ mole}$$

$$[HI] = 3-2x = 3-2(0.32) = 2.36 \text{ moles.}$$

$$\% = \frac{.64}{3.00} = .213 \times 100\% = 21.3\%$$

• PROBLEM 318

For the reaction $I_2(g) \rightleftharpoons 2I(g)$, $K = 3.76 \times 10^{-5}$ at $1000^\circ K$. Suppose you inject 1 mole of I_2 into a 2.00 liter-box at $1000^\circ K$. What will be the final equilibrium concentrations of I_2 and of I ?

Solution: Final equilibrium concentrations can be determined from the equilibrium constant expression. For this reaction, K , the equilibrium constant, equals

$$\frac{[I]^2}{[I_2]}$$

You are given this value as 3.76×10^{-5} at $1000^\circ K$. Equating, therefore, you obtain

$$3.76 \times 10^{-5} = \frac{[I]^2}{[I_2]}$$

You inject 1 mole of I_2 in a 2 liter box, which means the initial concentration is 1 mole/2 liter = 0.5M. Let x = the amount of I_2 that decomposes. Then, at equilibrium $[I_2] = .5 - x$, i.e., the initial amount minus the decomposed amount. Whatever decomposes yields I . For every mole of I_2 that decomposes, two moles of I are generated. This can be seen from the coefficients in the chemical equation. Thus, at equilibrium, $[I] = 2x$. Substituting these values in the equilibrium constant expression, you obtain

$$\frac{(2x)^2}{(.5-x)} = 3.76 \times 10^{-5}$$

Solving for x you obtain $x = 2.17 \times 10^{-5}$, using the quadratic equation. Thus, the concentrations become

$$\text{and} \quad I = 2x = 4.34 \times 10^{-5} \text{ M}$$

$$I_2 = .5 - x = .498 \text{ M}.$$

• PROBLEM 319

Given the equilibria $\text{H}_2\text{S} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HS}^-$ and $\text{HS}^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{S}^{2-}$, find the final concentration of S^{2-} , if the final concentrations of H_3O^+ and H_2S are 0.3 M and 0.1 M, respectively.

$$k_1 = 6.3 \times 10^{-8} \text{ for } \text{H}_2\text{S} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HS}^- \quad \text{and}$$

$$k_2 = 1 \times 10^{-14} \text{ for } \text{HS}^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{S}^{2-}.$$

Solution: This problem can be solved by writing the equilibrium constant expressions for the equilibria. These expressions give the ratio of the concentrations of products to reactants, each raised to the power of its coefficient in the equilibrium equation.

Therefore, for $\text{H}_2\text{S} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HS}^-$, we can write

$$k_1 = 6.3 \times 10^{-8} = \frac{[\text{H}_3\text{O}^+][\text{HS}^-]}{[\text{H}_2\text{S}]}$$

For $\text{HS}^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{S}^{2-}$,

$$k_2 = 1 \times 10^{-14} = \frac{[\text{H}_3\text{O}^+][\text{S}^{2-}]}{[\text{HS}^-]}$$

Note: water concentration is not included in the equilibrium expression since its concentration is assumed to be constant.

One is not given any information concerning $[\text{HS}^-]$. $[\text{HS}^-]$ is common to both k_1 and k_2 , so that if one solves one equation for $[\text{HS}^-]$ and substitutes it into the other equation, $[\text{HS}^-]$ is eliminated. Proceed as follows:

If one solves for $[\text{HS}^-]$ in k_1 , one obtains

$$[\text{HS}^-] = \frac{k_1[\text{H}_2\text{S}]}{[\text{H}_3\text{O}^+]} = \frac{6.3 \times 10^{-8} [\text{H}_2\text{S}]}{[\text{H}_3\text{O}^+]}$$

Substituting this into k_2 , one obtains

$$k_2 = 1 \times 10^{-14} = \frac{[\text{H}_3\text{O}^+][\text{S}^{2-}]}{[\text{HS}^-]} = \frac{[\text{H}_3\text{O}^+][\text{S}^{2-}]}{\frac{6.3 \times 10^{-8} [\text{H}_2\text{S}]}{[\text{H}_3\text{O}^+]}}$$

Rewriting in terms of $[\text{S}^{2-}]$,

$$[S^{-2}] = \frac{(1 \times 10^{-14})(6.3 \times 10^{-8})[H_2S]}{[H_3O^+]^2}$$

One is given that $[H_2S]$ and $[H_3O^+]$ equal 0.1 M and 0.3 M, respectively.

Therefore, one can substitute these values into this equation to solve for $[S^{-2}]$.

$$[S^{-2}] = \frac{(1 \times 10^{-14})(6.3 \times 10^{-8})(0.1)}{(0.3)^2} = 7 \times 10^{-22} \text{ M.}$$

• PROBLEM 320

At 1000°K , $K = 2.37 \times 10^{-3}$ for the reaction, $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$. If you inject one mole of each N_2 and H_2 in a one-liter box at 1000°K , what per cent of the H_2 will be converted to NH_3 at equilibrium?

Solution: To answer this question, determine the concentration of H_2 at equilibrium. Once this is known, subtract it from the initial concentration. The difference yields the amount that reacted to produce NH_3 . By multiplying the quotient of the difference divided by the original amount of H_2 by 100, you obtain the per cent.

To find $[H_2]$ at equilibrium, employ the equilibrium constant expression. It states that K , the equilibrium constant, is equal to the concentration ratio of the products to reactants, each raised to the power of its coefficient in the chemical reaction. For this reaction, then,

$$K = 2.37 \times 10^{-3} = \frac{[NH_3]^2}{[H_2]^3[N_2]}$$

To find $[H_2]$, proceed as follows. Let x = amount of N_2 that dissociates. The initial concentrations of all species are 1 M, since molarity = moles/liter, and 1 mole of each is placed in a one-liter box. If the N_2 's initial concentration is one and x moles/liter dissociates, then, at equilibrium, there is $(1 - x)$ moles/liter left of N_2 . In other words, at equilibrium, $[N_2] = 1 - x$. Now, from the chemical equation, it is seen that for every mole of N_2 that reacts, 3 moles of H_2 are necessary. Thus, when x moles/liter of N_2 dissociate, $3x$ moles/liter of H_2 are required. The initial concentration is 1, so that, at equilibrium

$$[H_2] = (1 - 3x) \text{ moles/liter.}$$

Notice, also, that for every mole/liter of nitrogen that dissociates, 2(mole/liter) of ammonia is obtained. Substituting these values into the equilibrium constant expression, one obtains

$$2.37 \times 10^{-3} = \frac{[NH_3]^2}{[N_2][H_2]^3} = \frac{(2x)^2}{(1-x)(1-3x)^3}$$

Solving for x , one obtains $x = .0217$ moles/liter of N_2 that dissociate. Thus, $[H_2] = 1 - 3x = .935$ moles/liter at equilibrium. The initial concentration was 1 mole/liter. The difference is the amount

that dissociated, i.e., the H_2 . The difference = $1 - .935 = .065$. Thus, the percent that dissociated equals

$$\frac{.065M}{1M} \times 100 = 6.5\%$$

• PROBLEM 321

For $PCl_5 \rightleftharpoons PCl_3 + Cl_2$, $K = 33.3$ at $760^\circ K$. Suppose 1.00g of PCl_5 is injected into a 500-ml evacuated flask at $760^\circ K$ and it is allowed to come into equilibrium. What percent of the PCl_5 will decompose? M.W. of $PCl_5 = 208.235$.

Solution: To find the percent of decomposition of PCl_5 , you need to know its initial concentration and final concentration (i.e., equilibrium concentration).

Initially, there is 1.00g of PCl_5 in the 500ml flask. Concentration = moles/liter. There are 1000ml in 1 liter, so that 500ml is .5 liters. The molecular weight of $PCl_5 = 208.235$ grams/mole. A mole is defined as

$$\frac{\text{grams (mass)}}{\text{molecular wt.}}$$

Thus, number of moles of

$$PCl_5 = \frac{1g}{208.235g/mole} = .0048 \text{ moles.}$$

Therefore, the initial concentration of

$$PCl_5 = \frac{.0048 \text{ moles}}{.5 \text{ liter}} = .0096M.$$

To find its concentration at equilibrium, use the equilibrium constant expression, which states that K , the equilibrium constant, is equal to the ratio of the concentrations of products to reactants, each raised to the power of its coefficient in the equation. Here, you have

$$K = \frac{[Cl_2][PCl_3]}{[PCl_5]}.$$

But, you are told $K = 33.3$. You can equate to obtain

$$33.3 = \frac{[Cl_2][PCl_3]}{[PCl_5]}.$$

Let x = concentration of $[Cl_2]$ at equilibrium. If this is the case, then $x = [PCl_3]$ also, since from the chemical equation it is shown that they are formed in equimolar amounts. If x moles/liter of each Cl_2 and PCl_3 form, and the only source is PCl_5 , then the $[PCl_5]$ at equilibrium is the initial concentration minus

$$x \frac{\text{moles}}{\text{liter}} = .0096 - x.$$

Substituting these values into the equilibrium constant expression, you obtain

$$33.3 = \frac{(x)(x)}{.0096 - x}.$$

Solving for x , using the quadratic equation, you obtain

$$x = 9.597 \times 10^{-3} = [PCl_3] = [Cl_2]$$

at equilibrium. $[PCl_5] = .0096 - .009597 = 3 \times 10^{-6}$. This means percent dissociation =

$$100 \times \frac{(\text{Initial} - \text{final}) \text{ concentration}}{\text{initial concentration}} =$$

$$100 \times \frac{.0096 - 3 \times 10^{-6}}{.0096} = 99.9 \% \text{ dissociated.}$$

• PROBLEM 322

At 273°K and 1 atm, 1 liter of N_2O_4 decomposes to NO_2 according to the equation



To what extent has decomposition proceeded when the original volume is 25 percent less than the existing volume?

Solution: Volume is proportional to the number of moles present when Pressure (P) and Temperature (T) are held constant. This can be seen from the equation of state,

$$PV = nRT \quad \text{or} \quad V = \frac{nRT}{P},$$

where R = universal gas constant, n = moles, and V = volume. When P and T are constant, RT/P is a constant, and V varies directly with n. Thus, in this problem you can discuss volumes in terms of fractions decomposed.


Let a be the volume fraction of the original N_2O_4 that decomposes. Since moles are proportional to volumes and 2 moles of NO_2 are produced for every mole of N_2O_4 , this fraction, a, results in the production of 2a of NO_2 and 1 - a of N_2O_4 . The total final volume fraction is 1 - a + 2a = (1 + a); that is, the final volume = (1 + a) times the original volume. The original volume is 1 liter of N_2O_4 . The existing volume is

$$(1 + a)(1 \text{ liter}) = (1 + a) \text{ liter.}$$

But 1 = .75 (1 + a). Solving, a = .333. Thus, you have 33% decomposition.

• PROBLEM 323

At 395 K, chlorine and carbon monoxide react to form phosgene, COCl_2 , according to the equation




The equilibrium partial pressures are $p_{\text{Cl}_2} = 0.128 \text{ atm}$, $p_{\text{CO}} = 0.116 \text{ atm}$, and $p_{\text{COCl}_2} = 0.334 \text{ atm}$. Determine the equilibrium constant K_p for the dissociation of phosgene

and the degree of dissociation at 395 K under a pressure of 1 atm.

Solution: After obtaining an expression for K_p for the dissociation of phosgene, the degree of dissociation under 1 atm of total pressure will be obtained by combining K_p with Dalton's law of partial pressures.

The dissociation of phosgene may be written as



By definition, K_p is the product of the partial pressures of the products divided by the product of the partial pressure of the reactants. Hence,

$$K_p = \frac{P_{\text{CO}} P_{\text{Cl}_2}}{P_{\text{COCl}_2}} = \frac{(0.116 \text{ atm})(0.128 \text{ atm})}{(.334 \text{ atm})} \\ = 0.0444 \text{ atm}.$$

K_p for the dissociation of phosgene is thus 0.0444 atm.

Let α denote the fraction of the original number of moles of phosgene that decomposed. Then, $1 - \alpha$ is the fraction of the original number of moles of phosgene remaining. From the stoichiometry of the dissociation reaction, one mole of CO and one mole of Cl_2 are formed for every mole of phosgene that decomposes. Thus, α moles of phosgene decomposes to α moles of CO and α moles of Cl_2 .

From Dalton's law of partial pressures,

$$P_{\text{COCl}_2} = X_{\text{COCl}_2} P_T$$

$$P_{\text{CO}} = X_{\text{CO}} P_T \quad \text{and} \quad P_{\text{Cl}_2} = X_{\text{Cl}_2} P_T$$

where X_{COCl_2} is the mole fraction of phosgene, X_{CO} is the mole fraction of CO, X_{Cl_2} is the mole fraction of Cl_2 , and P_T is the total pressure. Mole fraction may be defined as the number of moles of that particular substance divided by the total number of moles present. Since the total number of moles present after α moles of phosgene decomposes is

$$(1 - \alpha) (\text{from remaining phosgene}) + \alpha (\text{from CO}) + \alpha (\text{from Cl}_2) \\ = 1 + \alpha$$

$$\text{we have} \quad P_{\text{COCl}_2} = X_{\text{COCl}_2} P_T = \frac{1 - \alpha}{1 + \alpha} P_T$$

$$P_{\text{CO}} = X_{\text{CO}} P_T = \frac{\alpha}{1 + \alpha} P_T$$

and $P_{\text{Cl}_2} = x_{\text{Cl}_2} P_T = \frac{\alpha}{1 + \alpha} P_T$.

Substituting these into the expression for K_p , we obtain

$$K_p = \frac{P_{\text{CO}} P_{\text{Cl}_2}}{P_{\text{COCl}_2}} = \frac{\left(\frac{\alpha}{1 + \alpha} P_T \right) \left(\frac{\alpha}{1 + \alpha} P_T \right)}{\frac{1 - \alpha}{1 + \alpha} P_T} = \frac{\alpha^2}{(1 + \alpha)(1 - \alpha)} P_T$$

$$= \frac{\alpha^2}{(1 - \alpha^2)} P_T,$$

Now, the total pressure is $P_T = 1$ atm and K_p has been determined as $K_p = 0.0444$ atm. Hence,

$$K_p = \frac{\alpha^2}{1 - \alpha^2} P_T,$$

$$0.0444 \text{ atm} = \frac{\alpha^2}{1 - \alpha^2} \times 1 \text{ atm},$$

$$0.0444 - 0.0444 \alpha^2 = \alpha^2,$$

or, $\alpha = \left(\frac{0.0444}{1.0444} \right)^{1/2} = 0.206.$

The degree of dissociation of phosgene is equal to the fraction, α , of original moles that have dissociated. Hence, the degree of dissociation of phosgene at 395 K under a pressure of 1 atm is 0.206.

• PROBLEM 324

If 1.588 g of nitrogen tetroxide gives a total pressure of 1 atm when partially dissociated in a 500-cm³ glass vessel at 25°, what is the degree of dissociation, α ? What is the value of K_p ? The equation for this reaction is $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$.

Solution: The determination of the density of a partially dissociated gas provides one method for measuring the extent to which the gas dissociates. When a gas dissociates, more molecules are produced, and at constant temperature and pressure the volume increases. The density at constant pressure then decreases, and the difference between the density of the undissociated gas and that of the partially dissociated gas is directly related to the degree of dissociation.

If α represents the fraction of gas dissociated, then $1 - \alpha$ represents the fraction of gas undissociated. Assum-

ing one starts with 1 mole of gas the number of moles of gaseous products in the balanced chemical equation is $1 + \sum v_i$ (where $\sum v_i$ is the summation of all the coefficients; positive coefficients for the products and negative coefficients for the reactants). Therefore, the number of moles of gas present at equilibrium is

$$(1 - \alpha) + (1 + \sum v_i)\alpha = 1 + \alpha \sum v_i$$

The density of an ideal gas at constant pressure and temperature is inversely proportional to the number of moles for a given weight and the ratio of the density ρ_1 of the undissociated gas to the density ρ_2 of the partially dissociated gas is given by the expression

$$\frac{\rho_1}{\rho_2} = 1 + \alpha \sum v_i$$

$$\alpha = \frac{\rho_1 - \rho_2}{\rho_2 \sum v_i}$$


If there is no dissociation, then $\alpha = 0$ and $\rho_1 = \rho_2$, if dissociation is complete, then $\alpha = 1$, $\rho_2 \sum v_i = \rho_1 - \rho_2$ and $\rho_1 = (1 + \sum v_i)\rho_2$.

Molecular weights are proportional at constant temperature and pressure to the gas densities. Therefore, molecular weights can be substituted for the densities:

$$\alpha = \frac{M_1 - M_2}{M_2 \sum v_i}$$

where M_1 is the molecular weight of the undissociated gas, and M_2 is the average molecular weight of the gases when the gas is partially dissociated.

The dissociation of nitrogen tetroxide proceeds by the following equation:



The degree of dissociation is α ; $(1 - \alpha)$ is proportional to the number of moles of undissociated N_2O_4 ; 2α is proportional to the number of moles of NO_2 ; and $(1 - \alpha) + 2\alpha$ or $1 + \alpha$ is proportional to the total number of moles.

If the total pressure of N_2O_4 plus NO_2 is P , the partial pressures are:

$$P_{\text{N}_2\text{O}_4} = \frac{1 - \alpha}{1 + \alpha} P \quad \text{and} \quad P_{\text{NO}_2} = \frac{2\alpha}{1 + \alpha} P$$

$$\text{Then, } K_p = \frac{\left(\frac{2\alpha}{1+\alpha}P\right)^2}{\left(\frac{1-\alpha}{1+\alpha}P\right)} = \frac{4\alpha^2 P}{1-\alpha^2}$$

$$M_1 = 92.02 \text{ g/mole.}$$

Using the Combined gas Law,

$$PV = n RT$$

and setting n , the number of moles of substance, equal to the mass of substance, m , divided by its molecular weight, M_2 , one obtains

$$PV = \frac{m}{M_2} RT$$

Rearranging and solving for M_2 ,

$$M_2 = \frac{RT}{P} \times \frac{m}{V}$$

$$M_2 = \frac{RT}{P} \frac{m}{V} = \frac{(0.082 \text{ liter atm}^\circ\text{K}^{-1} \text{ mole}^{-1})(298^\circ\text{K})(1.588 \text{ g})}{(1 \text{ atm})(0.500 \text{ liter})}$$

$$= 77.68 \text{ g/mole}$$

$$\alpha = \frac{M_1 - M_2}{M_2 \sum v_i}$$

where $\sum v_i$ for the reaction equals [(coefficient of NO_2) - (coefficient of N_2O_4) = (2 - 1) = 1] and $M_1 = 92.02$ and $M_2 = 77.68$.

$$\alpha = \frac{92.02 - 77.68}{77.68} = 0.1846$$

$$K_p = \frac{4\alpha^2 P}{1-\alpha^2} = \frac{4(0.1846)^2(1 \text{ atm})}{1-(0.1846)^2} = 0.141.$$

THE SHIFTING OF EQUILIBRIUM-LE CHATELIER'S PRINCIPLE

• PROBLEM 325

A solute of formula AB is slightly dissociated into A^+ and B^- . In this system, there is a dynamic equilibrium such that $\text{A}^+ + \text{B}^- \rightleftharpoons \text{AB}$. Explain what happens if more acid is introduced into this system.

Solution: An acid is a species which, when added to a solvent (such as H_2O), dissociates into protons (H^+) and anions. In this particular case, the proton is represented as A^+ . When more acid is added to this general solvent system, more A^+ is introduced. The increased concentration of A^+ places a stress on the equilibrium and the result is a shift in this equilibrium. According to Le Châtelier's principle, an equilibrium system will readjust to reduce a stress if one is applied. Thus, the equilibrium $\text{A}^+ + \text{B}^- \rightleftharpoons \text{AB}$ will readjust to relieve the stress of the increased A^+ concentration. The stress is relieved by the reaction of A^+ with B^- to produce more AB . The concentration of B^- will decrease as compared to its concentration prior to the addition of the acid. Also, the concentration of the product AB will increase with the addition of the acid.

• PROBLEM 326

At 986°C , $K = 1.60$ for the reaction, $\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{H}_2\text{O}(\text{g}) + \text{CO}(\text{g})$.

If you inject one mole each of H_2 , CO_2 , H_2O , and CO simultaneously in a 20-liter box at time $t = 0$ and allow them to equilibrate at 986°C , what will be the final concentrations of all the species? What would happen to these concentrations if additional H_2 was injected and a new equilibrium was established?

Solution: Final concentrations of the species can be found by using the equilibrium constant expression. This expression equates K , the equilibrium constant, to the concentration ratio of products to reactants, each raised to the power of its coefficient in the chemical equation. Thus, for this reaction, you can say

$$K = \frac{[\text{H}_2\text{O}][\text{CO}]}{[\text{H}_2][\text{CO}_2]} = 1.60$$

Initially, there was 1 mole of each component in the 20 liter container. Since, concentration = moles/liter, all had an initial concentration of

$$\frac{1 \text{ mole}}{20 \text{ liter}} = .05\text{M}$$

At equilibrium, let x = the number of moles/liter of H_2 that have reacted. Thus, its concentration, at equilibrium, becomes $.05 - x$. If x moles/liter of H_2 react, the same number of moles/liter of CO_2 must react also, since they react in equimolar amounts; this is seen from the chemical reaction. Thus, $[\text{CO}_2] = .05 - x$, at equilibrium. These

x moles/liter have been converted to products. Thus, $[\text{H}_2\text{O}] = [\text{CO}] = .05 + x$, at equilibrium. The two products have the same concentration, since, again, the reaction shows they are formed in equimolar amounts. As such, you can substitute these values to obtain

$$\frac{(.05+x)(.05+x)}{(.05-x)(.05-x)} = 1.60$$

or

$$\frac{.05 + x}{.05 - x} = \sqrt{1.60}$$

Solving for x , you obtain $x = .00585$. Thus, the concentrations become $[H_2] = [CO_2] = .0442$ and $[H_2O] = [CO] = .0558$. If more H_2 is injected, the equilibrium is subjected to a stress, one component's concentration has been increased, and according to Le Chatelier's principle, the system will act to relieve the stress by shifting the equilibrium. To do this, more H_2 reacts with CO_2 , thus, decreasing their concentrations, to produce more H_2O and CO , thereby, increasing their concentrations.

● PROBLEM 327

You are given a box in which $PCl_5(g)$, $PCl_3(g)$, and $Cl_2(g)$ are in equilibrium with each other at $546^\circ K$. Assuming that the decomposition of PCl_5 to PCl_3 and Cl_2 is endothermic, what effect would there be on the concentration of PCl_5 in the box if each of the following changes were made? (a) Add Cl_2 to the box, (b) Reduce the volume of the box, and (c) Raise the temperature of the system.


Solution: You are told that the following equilibrium exists in the box $PCl_5 \rightleftharpoons PCl_3 + Cl_2$ (all gases) and asked to see what happens to $[PCl_5]$ when certain changes are made. This necessitates the use of Le Chatelier's principle, which states that if a stress is applied to a system at equilibrium, then the system readjusts to reduce the stress. With this in mind, proceed as follows:

(a) Here, you are adding Cl_2 to the box. This results in a stress, since one of the components in the equilibrium has its concentration increased. According to Le Chatelier's principle, the system will act to relieve this increased concentration of Cl_2 - the stress. It can do so, if the Cl_2 combines with PCl_3 to produce more PCl_5 . In this fashion, the stress is reduced, but the concentration of PCl_5 is increased.

(b) When the volume of the box is reduced, the concentration of the species is increased, i.e., the molecules are crowded closer together. Thus, a stress is applied. The stress can only be relieved (Le Chatelier's principle) if the molecules could be reduced in number. Notice, in our equilibrium expression you have 2 molecules, 1 each of PCl_3 and Cl_2 , producing 1 molecule of PCl_5 . In other words, the number of molecules is reduced if the equilibrium shifts to the left, so that more PCl_5 is produced. This is exactly what happens. As such, the $[PCl_5]$ increases.

(c) You are told that the decomposition of PCl_5 is endothermic (absorbing heat). In other words, it must absorb heat from the surroundings to proceed. If you increase the temperature, more heat is available, and the decomposition proceeds more readily, which means $[PCl_5]$ decreases. This fact can also be seen from the equilibrium constant of the reaction, K . This constant measures the ratio of products to reactants, each raised to the power of its coefficients in the chemical reaction. Now, when a reaction is endothermic, K is increased. For K to increase, the reactant's concentration must decrease. Again, therefore, you see that $[PCl_5]$ decreases.

A 1.00-liter reaction vessel containing the equilibrium mixture



was found to contain 0.400 mole of COCl_2 , 0.100 mole of CO , and 0.500 mole of Cl_2 . If 0.300 mole of CO is added at constant temperature, what will be the new concentrations of each component at equilibrium?

Solution: If a stress is placed on a system in equilibrium, whereby the equilibrium is altered, that change will take place which tends to relieve or neutralize the effect of the added stress. Thus, in this reaction when more CO is added after equilibrium has been established more COCl_2 will be formed to re-establish the equilibrium. One is given the concentrations of the components at equilibrium, thus the equilibrium constant (K_{eq}) can be calculated. The equilibrium constant for this reaction can be stated:

$$K_{eq} = \frac{[\text{COCl}_2]}{[\text{CO}][\text{Cl}_2]}, \text{ the ratio of products to}$$

reactants, where [] indicate concentration. One can solve for the K_{eq} by using the concentrations given for the first equilibrium,

$$K_{eq} = \frac{[\text{COCl}_2]}{[\text{CO}][\text{Cl}_2]} \quad \begin{array}{l} [\text{COCl}_2] = 0.400 \text{ mole/liter} \\ [\text{CO}] = 0.100 \text{ mole/liter} \\ [\text{Cl}_2] = 0.500 \text{ mole/liter} \end{array}$$

$$K_{eq} = \frac{(0.400)}{(0.100)(0.500)} = \frac{(0.400)}{(0.05)} = 8.0$$

One knows, from the definition of K_{eq} , that when more CO is added to the mixture that the concentrations of the components rearrange so that the equation for K_{eq} is equal to 8.

Using Le Chatelier's principle, stated in the first sentence of this explanation, one knows that when more CO is added to this mixture more COCl_2 is formed. Let new concentrations of $\text{COCl}_2 = [0.400 + x]$.

From the equation, one knows that for each mole of COCl_2 formed one mole of Cl_2 is used, thus the new concentration of $\text{Cl}_2 = [0.500 - x]$. One mole of CO is also used for each mole of COCl_2 formed, thus the new concentration of CO will be equal to x subtracted from the sum of the number of moles originally present and the number added. The new concentration of $\text{CO} = [0.100 + 0.300 - x]$. Because one knows that $K_{eq} = 8$, one can now solve for x .

$$K_{eq} = \frac{[\text{COCl}_2]}{[\text{CO}][\text{Cl}_2]} = 8 \text{ where } \begin{array}{l} [\text{COCl}_2] = (0.400 + x) \text{ moles} \\ [\text{CO}] = (0.100 + 0.300 - x) \text{ moles} \\ [\text{Cl}_2] = (0.500 - x) \text{ moles} \end{array}$$

Solving,

$$8 = \frac{(0.400 + x)}{(0.100 + 0.300 - x)(0.500 - x)}$$

$$\frac{(0.400 + x)}{(0.2 - 0.9x + x^2)} = 8$$

$$(0.400 + x) = 8(0.2 - 0.9x + x^2)$$

$$(0.400 + x) = (1.6 - 7.2x + 8x^2)$$

$$0 = 1.2 - 8.2x + 8x^2$$

One uses the quadratic equation to solve for x :

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, \text{ where}$$

$$0 = c + bx + ax^2$$

$$x = \frac{8.2 \pm \sqrt{(-8.2)^2 - 4(8 \times 1.2)}}{2 \times 8}$$

$$x = \frac{8.2 \pm 5.37}{16}$$

$$x = \frac{8.2 + 5.37}{16} = 0.85$$

or

$$x = \frac{8.2 - 5.37}{16} = .177$$

One cannot use $x = .85$ in this problem because when it is used, the concentrations of CO and Cl_2 will be negative values. Thus

$x = .177$. The new concentrations can now be found.


$$[\text{COCl}_2] = (0.400 + x) = .577 \text{ moles}$$

$$[\text{CO}] = (0.400 - x) = .223 \text{ moles}$$

$$[\text{Cl}_2] = (0.500 - x) = .323 \text{ moles}$$

• PROBLEM 329

At a certain temperature, an equilibrium mixture of



is analyzed and found to contain the following molar concentrations:

$$[\text{NO}_2] = 0.100, [\text{SO}_2] = 0.300, [\text{NO}] = 2.00, [\text{SO}_3] = 0.600$$

If 0.500 moles of SO_2 are introduced at constant temperature, what will be the new concentrations of reactants and products when equilibrium is re-established?

Solution: Le Chatelier's principle states: If a stress is placed on a system in equilibrium, whereby the equilibrium is altered, that change will take place which tends to relieve or neutralize the effect of the added stress. Thus, in this reaction, if more SO_2 is added more NO and SO_3 will be formed. If stress is placed on the left side of the equation, the reaction will be forced to the right (and vice versa). One can determine the equilibrium constant (K_{eq}) for this reaction by using the concentrations of the original mixture. The equilibrium constant is defined:

$$K_{eq} = \frac{[\text{NO}][\text{SO}_3]}{[\text{NO}_2][\text{SO}_2]}$$

where [] indicate concentrations. Solving for K_{eq} with

$$\begin{aligned} [\text{NO}] &= 2.00 \\ [\text{SO}_3] &= 0.600 \end{aligned}$$

$$\begin{aligned} [\text{NO}_2] &= 0.100 \\ [\text{SO}_2] &= 0.300 \end{aligned}$$

one obtains

$$K_{eq} = \frac{(2.00)(0.600)}{(0.100)(0.300)} = 40.0$$

One can solve for the new concentrations by using the K_{eq} . From Le Chatelier's principle, one knows that when SO_2 is added to this mixture, the amounts of NO and SO_3 will increase. Let x = the number of moles by which NO and SO_3 will increase. For each mole of SO_3 and NO formed, one mole of SO_2 and one mole of NO_2 will react, thus the new concentrations of these components will be equal to the original concentrations less x moles. The new concentrations can be stated.

$$\begin{aligned} [\text{NO}] &= 2.00 + x \\ [\text{SO}_3] &= 0.600 + x \\ [\text{NO}_2] &= 0.100 - x \\ [\text{SO}_2] &= 0.300 + \text{the amount added} - x \\ &= 0.300 + 0.500 - x = 0.800 - x \end{aligned}$$

Using the formula for the equilibrium constant, one can solve for x .

$$K_{eq} = \frac{[\text{NO}][\text{SO}_3]}{[\text{NO}_2][\text{SO}_2]} = 40.0$$

Substituting,

$$\begin{aligned} 40.0 &= \frac{(2.00 + x)(0.600 + x)}{(0.100 - x)(0.800 - x)} \\ 40.0 &= \frac{1.20 + 2.6x + x^2}{.080 - 0.90x + x^2} \end{aligned}$$

$$(0.80 - 0.90x + x^2)40 = 1.20 + 2.6x + x^2$$

$$3.20 - 36.0x + 40x^2 = 1.20 + 2.6x + x^2$$

$$2.0 - 38.6x + 39x^2 = 0$$

One can use the quadratic formula to solve for x .

$$\begin{aligned} ax^2 + bx + c &= 0 \\ x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ 39x^2 - 38.6x + 2.0 &= 0 \\ x &= \frac{38.6 \pm \sqrt{(38.6)^2 - 4 \times 2.0 \times 39}}{2 \times 39} \end{aligned}$$

$$x = \frac{38.6 \pm 34.44}{76}$$

$$x = \frac{38.6 + 34.44}{76} = 0.96$$

or

$$x = \frac{38.6 - 34.44}{76} = 0.055$$

One cannot use $x = 0.96$ because $[\text{NO}_2]$ and $[\text{SO}_2]$ will be negative.

Concentrations cannot have negative values, which means $x = .055$. One can now find the new concentrations

$$[\text{NO}] = 2.00 + x = 2.055 \text{ moles}$$

$$[\text{SO}_3] = 0.600 + x = 0.65 \text{ moles}$$

$$[\text{NO}_2] = 0.100 - x = 0.045 \text{ moles}$$

$$[\text{SO}_2] = 0.800 - x = 0.745 \text{ moles}$$

CHAPTER 10

ACID-BASE EQUILIBRIA

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 305 to 366 for step-by-step solutions to problems.

A simple definition of an acid in an aqueous solution is any compound which produces hydrated hydrogen ions, H_3O^+ (also called hydronium ions). Bases are compounds which produce hydroxyl, OH^- , ions. There are more sophisticated definitions such as Lewis acids and bases (compounds which can “accept” or “donate,” respectively, a share in an electron pair) and Brönsted acids and bases (compounds which act as proton donors and proton acceptors, respectively), but the simple definition works well for most aqueous phase acid-base reactions. While the hydrated hydrogen ion, H_3O^+ , is always the species that actually exists in aqueous solution, the writing of most reactions is simplified by showing it as the hydrogen ion, H^+ . Such a simplification is helpful in writing acid-base reactions because the hydrating H_2O molecules are left out.

Water ionizes to form hydrogen (or hydronium) and hydroxyl ions according to the equation



(Note that if the same reaction is written with a hydronium ion, the equation becomes $2\text{H}_2\text{O} \leftrightarrow \text{H}_3\text{O}^+ + \text{OH}^-$.)

Reaction 10-1, the autoionization of water, has an equilibrium constant of 10^{-14} . Since the activity of pure water is one by definition, the product of the H^+ and OH^- concentrations in an aqueous solution is always 10^{-14} .

$$(\text{H}^+)(\text{OH}^-) = 10^{-14} = K_w \quad 10-2$$

Similar reactions in non-aqueous solvents (e.g., $2\text{NH}_3 \leftrightarrow \text{NH}_4^+ + \text{NH}_2^-$) are called autoprotolysis. They are simply reversible ionization reactions resulting from the shift of a proton. Usually they reach equilibrium quickly.

A widely used measure of acidity, pH, is defined as the negative logarithm of the H^+ ion concentration.

$$\text{pH} = -\log_{10}(\text{H}^+) \quad 10-3$$

Hence, the H^+ ion concentration is equal to $10^{(-\text{pH})}$. When both the H^+ and OH^- ion concentrations are equal, Equation 10-2 dictates that they both equal 10^{-7} . The neutral pH, therefore, is 7; acidic solutions have a pH less than 7, basic solutions greater than 7. A similar quantity, pOH, is defined for the OH^- ion concentration, but it is much less common than pH. The sum of pH and pOH will always equal 14 in order to satisfy Equation 10-2.

The ionization constant, dissociation constant, and hydrolysis constant, which are illustrated in Problems 347–363, are all equilibrium constants. The problems can all be solved using the basic principles for solving for equilibrium compositions discussed in Chapter 9. Do not conclude from the different names assigned these equilibria that there are different principles involved. In each case the problem can be solved by the following steps:

- 1) Write the reaction.
- 2) Write the equation relating the equilibrium constant to compositions.
- 3) Write each composition in terms of a single variable from the stoichiometry.
- 4) Solve the equation for the single variable and calculate the equilibrium compositions.

Acid-base neutralization occurs in aqueous solution when the H^+ ions of the acid react with the OH^- ions of the base to form water. This reaction occurs so fast that it is usually considered instantaneous, and the H^+ and OH^- concentrations always obey the equilibrium relation of Equation 10-2. The other product of the neutralization of an acid and base — from the cation from the base and the anion from the acid — is called a salt. Normal table salt (NaCl) is the salt formed from the neutralization of hydrochloric acid (HCl) and sodium hydroxide (NaOH). It is, however, only one of numerous salts that form from the neutralization of acids and bases. The key to working the neutralization problems lies in remembering that one mole of H^+ ions reacts


with one mole of OH^- ions to form a neutral solution. The other ions will form a salt.

Diprotic or triprotic acids possess two or three hydrogens, respectively, that can ionize to form H^+ ions. In all such cases, the first H^+ ion is more readily formed than the second or third. A very common example is sulfuric acid which can form H^+ ions by both of the following reactions.



The equilibrium constant for the first H^+ ion, Equation 10-4, is very large and all the H_2SO_4 reacts. However, the equilibrium constant for the second is .012, so little H^+ is formed from this reaction. Let us calculate, for example, the SO_4^{2-} ion concentration in a 1M aqueous solution of sulfuric acid. Follow the procedure outlined above.

1) Write the equation containing SO_4^{2-} .



2) Write the equation relating the equilibrium constant to compositions.

$$0.012 = (\text{H}^+)(\text{SO}_4^{2-})/(\text{HSO}_4^-) \quad 10-7$$

3) Write the composition in terms of a single variable from the stoichiometry.

We can assume, since the equilibrium constant is small, that very little HSO_4^- is consumed and that, since Equation 10-4 goes essentially to completion, HSO_4^- and H^+ concentrations are equal and equal to 1M.

$$.012 = (1)(\text{SO}_4^{2-})/(1) \quad 10-8$$

4) Solve the equation for the single variable and calculate the equilibrium compositions.

$$(\text{SO}_4^{2-}) = .012\text{M} \quad 10-9$$

The assumption that very little HSO_4^- ion is consumed is reasonable. In this case, .012 moles or 1.2% of the HSO_4^- is consumed.

Aqueous buffered solutions, or aqueous solutions that maintain a very nearly constant pH or constant H^+ ion concentration, are formed in two common ways.

- 1) By dissolving, in water, a weak acid and a soluble ionic salt containing the same anion as the weak acid
- 2) By dissolving, in water, a weak base and a soluble ionic salt containing the same cation as the weak acid

When the anion of a weak acid is maintained at a constant, relatively large concentration by dissolving a salt, it will control, through equilibrium, the H^+ ion concentration and, hence, the pH. For example, acetic acid is a weak acid with a ionization equilibrium constant of 1.76×10^{-5} . We can write the reaction for the ionization of acetic acid, using the abbreviation Ac^- , for the acetate ion.



$$1.76 \times 10^{-5} = (H^+) (Ac^-)/(HAc) \quad 10-11$$


$$H^+ = 1.76 \times 10^{-5} (HAc)/(Ac^-) \quad 10-12$$

The concentration of HAc, unionized acetic acid, is constant because very little ionizes or reacts by Equation 10-10. If the concentration of acetate ion, Ac^- , is maintained at a large constant value (large by comparison to that produced by Equation 10-10) by adding a soluble ionic salt of acetic acid, such as sodium acetate, to the solution, then the ratio of HAc and Ac^- will be constant. The addition of small amounts of acids or bases to this solution will not result in significant changes in the Ac^- or HAc concentrations and, from Equation 10-12, the H^+ ion concentration or the pH will not change; the solution is said to be buffered against change in the pH.

Indicators are usually large organic molecules that change color by reaction with H^+ or OH^- ions. When the concentration of H^+ , for example, reaches an appropriate level, the equilibrium with the indicator molecule will shift to form a species of a different color. Many acid-base indicators exist; it is possible to select one that will change color at almost any desired pH. The reactions of H^+ with indicators reach equilibrium quickly and can be treated like any other equilibrium reaction.

Complex ions are simply ions that form by addition of a molecule (called a complexing agent) to existing ions. They can be treated like any other ion

and all of the principles of calculating ionic equilibrium apply. Ammonia, NH_3 , is a complexing agent with many metallic ions. A typical example is the one given in Problem 379.



Such ions are typically highly colored.

Electrolytes are compounds which form charged ions when dissolved. Most salts are electrolytes, as are most common inorganic acids and bases. Strong electrolytes ionize completely when dissolved; weak electrolytes, on the other hand, have small ionization equilibrium constants. Acetic acid, discussed previously, is a weak electrolyte with an ionization equilibrium constant of 1.76×10^{-5} . Treatment of the problems involving electrolytes can be solved by following the same principles discussed for other equilibrium reactions.

Step-by-Step Solutions to Problems in this Chapter, "Acid-Base Equilibria"

ACIDS AND BASES

• PROBLEM 330

Can I^+ (the iodine cation) be called a Lewis base? Explain your answer.

Solution: A Lewis base may be defined as an electron pair donor. Writing out its electronic structure is the best way to answer this question, because it will show the existence of any available electron pairs.

The electronic structure of I^+ may be written as

$[: I :]^+$. There are three available electron pairs. This might lead one to suspect that it is indeed a Lewis base. But note, I^+ does not have a complete octet of electrons, it does not obey the octet rule. According to this rule, atoms react to obtain an octet (8) of electrons. This confers stability.

Therefore, I^+ would certainly rather gain two more electrons than lose six. In reality, then, I^+ is an electron pair acceptor. Such substances are called Lewis acids.

• PROBLEM 331



Write the equations for the stepwise dissociation of pyrophosphoric acid, $H_4P_2O_7$. Identify all conjugate acid-base pairs.

Solution: Pyrophosphoric acid is an example of a polyprotic acid. Polyprotic acids furnish more than one proton

per molecule. From its molecular formula, $H_4P_2O_7$, one can see there exist four hydrogen atoms. This might lead one to suspect that it is tetraprotic, i.e. having 4 protons that can be donated per molecule. This is in fact the case, which means there exist four dissociation reactions. In general, the equation for a dissociation reaction is,



Polyprotic acids follow this pattern. Thus, one can write the following equations for the step-wise dissociation of $H_4P_2O_7$.



To identify all conjugate acid-base pairs, note the definition of the term. The base that results when an acid donates its proton is called the conjugate base. The acid that results when a base accepts a proton is called the conjugate acid. From these definitions, one sees that in all cases H_3O^+ is the conjugate acid of H_2O (the base in these reactions) and $H_3P_2O_7^-$, $H_2P_2O_7^{2-}$, $HP_2O_7^{3-}$ and $P_2O_7^{4-}$ are the conjugate bases of $H_4P_2O_7$, $H_3P_2O_7^-$, $H_2P_2O_7^{2-}$ and $HP_2O_7^{3-}$, respectively.




• PROBLEM 332

The dissociation sequence of the polyprotic acid H_3PO_4 shows three Bronsted-Lowry acids. Rank them in order of decreasing strengths.

Solution: Polyprotic acids are ones which furnish more than one proton per molecule. From its molecular formula, H_3PO_4 , it is observed that there are 3 available hydrogen atoms available for release. In general, the equation for a dissociation reaction is



where HA is the acid and water is acting as a weak base. With this in mind, one can write the 3 dissociation reactions as



From this, one can see that the three acids are

H_3PO_4 , H_2PO_4^- , and HPO_4^{2-} . The acids decrease in strength in the order of H_3PO_4 , H_2PO_4^- , HPO_4^{2-} .

Ranking can be explained by noting that equivalent H-O bonds are being broken to give off H^+ . The second and third protons that dissociate leave a progressively more negative ion. This means it is more difficult for the ion to dissociate in order to produce additional H^+ ions. This stems from the fact that the increased negativity results in increased attraction for the proton (H^+). In summary, as the negative charge of the acid increases, the weaker the acid becomes.

• PROBLEM 333

If you place HClO_4 , HNO_3 or HCl in water, you find that they are strong acids. However, they show distinct differences in acidities when dissolved in acetic acid. Such an occurrence is referred to as the leveling effect of the solvent, water: a) Explain the basis for this leveling effect by comparing acid reactions in the water solvent system to the acetic acid solvent system. b) Discuss the leveling effect in terms of basicities instead of acidities.

Solution: (a) To explain the leveling effect, you must consider the relative acidic or basic properties of the species involved. In water, the general reaction for the acid is $\text{HA} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{A}^-$. In acetic acid, however, the reaction is (assuming HA is a stronger acid than acetic acid (CH_3COOH), and the three acids given are) $\text{HA} + \text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COOH}_2^+ + \text{A}^-$. Let us consider the strengths of the acids in these two solvent systems. Water is less acidic, thus more basic and more strongly proton-attracting, than acetic acid. This means that when strong acids are dissolved the equilibrium will be shifted far to the right in water, but not as far in acetic acid. Thus, more products will be produced in the water solution than in the acetic acid. The acidities of the 3 given acids depend upon how much H^+ ion they produce. If the equilibrium is not shifted to the right, less H^+ ion is being produced. Thus, the acidities of the three given acids will be less in acetic acid, since the equilibrium is shifted less to the right, and thus, not much H^+ ion is generated.

(b) The leveling effect can also be thought of in terms of basicity. For the two types of bases, you have $\text{B} + \text{H}_2\text{O} \rightleftharpoons \text{BH}^+ + \text{OH}^-$ in water and $\text{B} + \text{CH}_3\text{COOH} \rightleftharpoons \text{BH}^+ + \text{CH}_3\text{COO}^-$ in acetic acid. Now OH^- is a stronger base than CH_3COO^- , and, therefore, the equilibrium is further to the left in water than in acetic acid. Thus, when a base is added to water it will ionize less than in acetic acid.

THE AUTOIONIZATION OF WATER

• PROBLEM 334

A 0.10 M solution of HCl is prepared. What species of ions are present at equilibrium, and what will be their equilibrium concentrations?

Solution: Two processes are occurring simultaneously: the reaction of HCl with H_2O (dissociation of HCl) and the autoionization of H_2O .


HCl reacts with H_2O according to the equation



For every mole of HCl that dissociates, one mole of Cl^- and one mole of H_3O^+ are produced. The initial concentration of HCl is 0.10 M. Thus, if we assume that HCl dissociates completely,

$$[\text{H}_3\text{O}^+] = 0.10 \text{ M} \quad \text{and} \quad [\text{Cl}^-] = 0.10 \text{ M}.$$

Water autoionizes according to the equation



The water constant for this process is

$$K_w = 10^{-14} \text{ moles}^2/\text{liter}^2 = [\text{H}_3\text{O}^+][\text{OH}^-].$$

$$\text{Hence, } [\text{OH}^-] = \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{[\text{H}_3\text{O}^+]}$$

Since $[\text{H}_3\text{O}^+]$ was determined to be $0.10 \text{ M} = 10^{-1} \text{ M} = 10^{-1} \text{ moles/liter}$,

$$\begin{aligned} [\text{OH}^-] &= \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{[\text{H}_3\text{O}^+]} \\ &= \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{10^{-1} \text{ moles/liter}} = 10^{-13} \text{ moles/liter} = 10^{-13} \text{ M}. \end{aligned}$$

Hence, at equilibrium, H_3O^+ , OH^- , and Cl^- are present in the concentrations

$$[\text{H}_3\text{O}^+] = 0.10 \text{ M}, [\text{OH}^-] = 10^{-13} \text{ M}, [\text{Cl}^-] = 0.10 \text{ M}.$$

• PROBLEM 335

A 0.10 M solution of NaOH is prepared. What species of ions are present at equilibrium, and what will be their equilibrium concentrations?

Solution: Two processes are occurring simultaneously, the dissociation of NaOH, and the autoionization of H_2O .

NaOH dissociates according to the equation



For every mole of NaOH that dissociates, one mole of Na^+ and one mole of OH^- are produced. The initial concentration of NaOH is 0.10 M. Thus, if we assume that NaOH dissociates completely $[Na^+] = 0.10$ M and $[OH^-] = 0.10$ M.

Water autoionizes according to the equation



The water constant for this process is

$$K_w = 10^{-14} \text{ moles}^2/\text{liter}^2 = [H_3O^+][OH^-].$$

$$\text{Hence, } [H_3O^+] = \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{[OH^-]}$$

Since $[OH^-]$ was determined to be 0.10 M = 10^{-1} M = 10^{-1} moles/liter, by substitution, we obtain

$$\begin{aligned} [H_3O^+] &= \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{[OH^-]} = \frac{10^{-14} \text{ moles}^2/\text{liter}^2}{10^{-1} \text{ moles/liter}} \\ &= 10^{-13} \text{ moles/liter} = 10^{-13} \text{ M.} \end{aligned}$$

Hence, at equilibrium, H_3O^+ , OH^- , and Na^+ are present in the concentrations

$$[H_3O^+] = 10^{-13} \text{ M}, [OH^-] = 0.10 \text{ M}, [Na^+] = 0.10 \text{ M.}$$

AUTOPROTOLYSIS

• PROBLEM 336


Find the equation for the autoprotolysis of water. Indicate which species is the acid, the base, the conjugate acid, and the conjugate base.

Solution: One can begin by defining autoprotolysis. It may be defined as the donation of a proton from a molecule of one specie to another molecule of the same specie to produce positive and negative ions. Thus, for water, the equation is $\text{H}_2\text{O} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{OH}^-$. An acid is defined as a specie that donates protons. A base is a substance that accepts protons. From the equation, one sees that either water (H_2O) molecule can be the base or acid. A conjugate base is a specie obtained by abstracting a proton (H^+). If one abstracts a proton from water, one obtains OH^- . Thus, OH^- is the conjugate base. The conjugate acid is defined as the base plus a proton. It was stated that either H_2O molecule could be the base. If one adds a proton to one of them, one obtains H_3O^+ . Thus, H_3O^+ is the conjugate acid.

• PROBLEM 337

Indicate the equilibrium equation and constant expression for the autoprotolysis of liquid ammonia. If $K_{\text{NH}_3} = 10^{-22}$, how many molecules of ammonia are ionized in 1 mole of ammonia? Assume a density of 0.771g/ml for ammonia.

Solution: Autoprotolysis is that phenomenon whereby an ammonia molecule can donate a proton to another NH_3 molecule to form positive and negative charged species. The equation of the autoprotolysis can be written $\text{NH}_3 + \text{NH}_3 \rightleftharpoons \text{NH}_4^+ + \text{NH}_2^-$ or



To find the constant expression, consider the equilibrium constant expression for the reaction:

$$K = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2}$$

Note, though, that the concentration of NH_3 in pure ammonia is always constant. By analogy with the autoprotolysis of water (where the K_w expression is written $[\text{OH}^-][\text{H}_3\text{O}^+]$, - without $[\text{H}_2\text{O}]^2$ in the denominator), the constant expression for the autoprotolysis of NH_3 is

$$(11) \quad K_{\text{NH}_3} = [\text{NH}_4^+][\text{NH}_2^-]$$

To find the number of molecules of ammonia ionized in 1 mole of ammonia, use the equation

$$K = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2}$$

Let x be the number of moles of ammonia ionized. Then the NH_3 remaining nonionized is $1-x$ moles. Since each 2 ammonia molecules must ionize to produce one NH_4^+ and one NH_2^- , the number of NH_4^+ =

number of $\text{NH}_2^- = \frac{x}{2}$. Let V be the volume of one mole of ammonia. The concentration of NH_4^+ and NH_2^- can be rewritten as $\frac{x/2}{V}$, and the concentration of nonionized NH_3 is $\frac{1-x}{V}$. The equation for K can be rewritten as

$$(iii) \quad K = \frac{(\frac{x/2}{V})(\frac{x/2}{V})}{(\frac{1-x}{V})^2} = \frac{x^2/4}{(1-x)^2} \cdot \frac{1/V^2}{1/V^2} = \frac{x^2}{4(1-x)^2}$$

x , the number of moles of ionized ammonia, can be calculated if K is known. To solve for K , consider a more general case of the equation

$$K = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2}$$

The numerator $[\text{NH}_4^+][\text{NH}_2^-]$ is the constant expression for the autoprotolysis of NH_3 and must always equal K_{NH_3} . K_{NH_3} is given as 10^{-22} .

To find $[\text{NH}_3]^2$, use the fact that the density of ammonia is 0.771g/ml. The mole weight of ammonia is 17.03g. Thus, 0.771g is $\frac{0.771\text{g}}{17.03\text{g/mole}} = 0.0453$ moles of ammonia; and the density of ammonia is 0.0453 moles/ml. = 45.3 moles/liter. Thus, $[\text{NH}_3] = 45.3\text{M}$. Substitute these results in

$$(iv) \quad K = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2} = \frac{10^{-22}}{45.3^2} = 4.9 \times 10^{-26}$$

Substitute this value of K into (iii)

$$(v) \quad 4.9 \times 10^{-26} = \frac{x^2}{4(1-x)^2}$$

To simplify the problem, note that the dissociation of ammonia is very small and thus $x \ll 1$. Thus, approximate $1-x^2$ as 1. Then, (v) becomes

$$(vi) \quad \frac{x^2}{4} = 4.9 \times 10^{-26}$$

Solve to obtain $x = 4.42 \times 10^{-13}$. This is the number of moles of NH_3 that ionized. To find the number of molecules, remember that 1 mole = 6.02×10^{23} molecules. Thus,

$$\begin{aligned} \text{No. of molecules ionized} &= (4.42 \times 10^{-13} \text{ moles}) (6.02 \times 10^{23} \text{ molecules/mole}) \\ &= 2.66 \times 10^{11} \text{ molecules.} \end{aligned}$$

2.66×10^{11} molecules of ammonia are ionized.

pH

• PROBLEM 338

A) Determine the pH of a solution with a hydrogen ion concentration of 3.5×10^{-4} .

B) If a solution has a pH of 4.25, what is the hydrogen ion concentration?

Solution: To determine the acidity or basicity of an aqueous solution, the hydrogen ion concentration must be measured. The pH of a solution expresses this concentration. pH is defined as the negative logarithm of the hydrogen ion concentration. In other words, $\text{pH} = (-\log [\text{H}^+])$, where the brackets around H^+ signify concentration. As such, to solve the problem, you must substitute into the equation. For part "A", you have $\text{pH} = -\log [3.5 \times 10^{-4}]$
now $-\log [3.5 \times 10^{-4}] = -\log 3.5 - \log 10^{-4}$

$$= -.54 - (-4)$$

$$= -.54 + 4$$

$$= 3.46$$

It follows, then, that the $\text{pH} = 3.46$ for a hydrogen ion concentration of 3.5×10^{-4} . Part "B" is similar, but here you are given the pH and asked to find the ion concentration. Therefore, you have $4.25 = -\log [\text{H}^+]$ or $-4.25 = \log [\text{H}^+]$. Now, logarithm numbers give only positive mantissas. As such, -4.25 must be in the form of $-5 + .75$. If you take the antilogarithm of each, $.75$ is 5.6 and -5 is 10^{-5} , you obtain a hydrogen ion concentration of 5.6×10^{-5} mole/liter

• PROBLEM 339

Determine the pH of each of the following solutions:
(a) 0.20 M HCl , (b) 0.10 M NaOH .

Solution: A pH scale has been devised to express the H_3O^+ concentration in solution. By definition,

$$\text{pH} = -\log [\text{H}_3\text{O}^+] \quad \text{or} \quad [\text{H}_3\text{O}^+] = 10^{-\text{pH}}$$

It has been shown that water dissociates to H_3O^+ and OH^-

ions to a small degree.



The equilibrium constant is defined as K_w for this reaction and is expressed as $[\text{H}_3\text{O}^+][\text{OH}^-]$. The H_2O does not appear, since it is presumed to be a constant. From the dissociation equation, it can be seen that the concentration of H_3O^+ equals OH^- . By experimentation, K_w has been shown to equal 1.0×10^{-14} . This means that in water, therefore, H_3O^+ and OH^- each have a concentration of 1.0×10^{-7} M. With this information in mind, one can now solve the problem.

(a) The concentration of HCl is 0.20 M. Since HCl is a strong electrolyte, dissociation is complete. Therefore, the concentration of H_3O^+ is also 0.20 M = 2.0×10^{-1} M. By definition, then

$$\text{pH} = -\log (2.0 \times 10^{-1}) = 1 - 0.3 = 0.7.$$

(b) The $[\text{OH}^-]$ equals the concentration of NaOH , since it is also a strong electrolyte. One wants the pH, therefore, employ the expression for K_w .

$$[\text{H}_3\text{O}^+] = \frac{K_w}{[\text{OH}^-]} = \frac{1 \times 10^{-14}}{0.10} = 1.0 \times 10^{-13} \text{ M.}$$

$$\text{Therefore, } \text{pH} = -\log (1.0 \times 10^{-13}) = 13.$$

• PROBLEM 340

A certain solution has pH 3.89 at 0°C . Find pOH and $[\text{OH}^-]$.

Solution: pH is a measure of the $[\text{H}^+]$ and pOH is a measure of $[\text{OH}^-]$. Their product gives K_w , the ionization constant of water:

$$[\text{H}^+][\text{OH}^-] = K_w.$$

pH and pOH are related by the equation,

$$\text{pOH} + \text{pH} = \text{p}K_w.$$

At 0°C , $\text{p}K_w = 14.94$. Therefore:

$$\text{pOH} = \text{p}K_w - \text{pH} = 14.94 - 3.89 = 11.05.$$

To find $[\text{OH}^-]$ use the equation,

$$\text{pOH} = -\log [\text{OH}^-]$$

$$11.05 = -\log [\text{OH}^-]$$

$$[\text{OH}^-] = 10^{-11.05} = 10^{-12+.95} = (10^{-12})(10^{.95}).$$

Find the antilog of $10^{.95}$. It is 8.9 which gives

$$[\text{OH}^-] = 8.9 \times 10^{-12}.$$


• PROBLEM 341

Assuming complete ionization, calculate (a) the pH of 0.0001 N HCl, (b) the pOH of 0.0001 N KOH.

Solution: (a) pH is defined as the negative log of the hydrogen ion concentration.

$$\text{pH} = -\log [\text{H}^+]$$

The normality of an acid is defined as the number of equivalents of H^+ per liter of solution. The ionization of HCl can be written



This means that there is one H^+ for every HCl, and that the concentration of H^+ equals the concentration of completely ionized HCl.

$$[\text{H}^+] = [\text{HCl}]$$

We are told that $[\text{HCl}] = 0.0001 \text{ N} = 1 \times 10^{-4} \text{ N}$. Therefore, $[\text{H}^+] = 1 \times 10^{-4} \text{ N}$. We can now solve for pH. Note: In this problem, normality = molarity (concentration), since equivalent weight = M.W.

$$\text{pH} = -\log [\text{H}^+]$$


$$\text{pH} = -\log (1 \times 10^{-4}) = 4.$$

The pH of this solution is 4.

(b) The pOH is defined as the negative log of the OH^- ion concentration.

$$\text{pOH} = -\log [\text{OH}^-].$$

The ionization of KOH can be stated



Therefore, one OH^- is formed for every KOH, and when KOH is completely ionized, their concentrations are equal.

$$[\text{KOH}] = [\text{OH}^-]$$

We are told that $[\text{KOH}] = 0.0001 \text{ N}$, thus $[\text{OH}^-] = 0.0001 \text{ N}$ (again, normality = molarity.)

Solving for pOH :

$$\begin{aligned}\text{pOH} &= -\log [\text{OH}^-] \\ &= -\log (0.0001) = -\log (1 \times 10^{-4}) = 4.\end{aligned}$$

The pOH of this solution is 4.

• PROBLEM 342

Find the pH of a solution in which $[\text{H}^+] = 6.38 \times 10^{-6}$ mole/liter.

Solution: pH indicates the acidity or basicity of a solution. It runs on a scale of 1 to 14. 1 is most acidic, 7 is neutral, and 14 is most basic. pH is a measure of $[\text{H}^+]$. The higher the $[\text{H}^+]$, the lower the pH , and the stronger the acid. To calculate pH use the equation:

$$\begin{aligned}\text{pH} &= -\log [\text{H}^+] = -\log [6.38 \times 10^{-6}] \\ &= -[(-6) + (\log 6.38)] = 6 + (-.805) \\ \text{pH} &= 5.195.\end{aligned}$$

• PROBLEM 343

What is the pH of a neutral solution at 50°C ?
 $\text{pK}_w = 13.26$ at 50°C .

Solution: A neutral solution is defined as $[\text{H}^+] = [\text{OH}^-]$; an acid solution has $[\text{H}^+] > [\text{OH}^-]$, and a basic solution has $[\text{H}^+] < [\text{OH}^-]$. For a solution at 25° , $\text{pK}_w = 14$.

pK_w indicates the amount of dissociation of water. To find the neutral pH , one lets $\text{pH} = \text{pOH} = x$. Since

$$\text{pH} + \text{pOH} = 2x = 14 = \text{pK}_w,$$

$$x = 7.$$

However, the solution in question is at 50° . At 50° $\text{pK}_w = 13.26$. Therefore, to find the neutral pH ,

$$\text{pH} + \text{pOH} = 2x = \text{pK}_w = 13.26$$

$$x = 6.63 = \text{neutral pH}.$$

Before the advent of pH meters, a urologist collected 1.3 liters of urine from a hospitalized patient over the course of a day. In order to calculate what the pH was a laboratory technician determined the number of equivalents of acid present in the sample. A given sample of urine contains 1.3×10^{-6} equivalents of dissociated acid. What is the pH of this sample?

Solution: To solve this problem, we have to note that 1 equivalent of H^+ is the same as 1 mole of H^+ . We then determine $[H^+]$ and from this the pH.

1.3×10^{-6} equivalent of H^+ is the same as 1.3×10^{-6} mole of H^+ . The concentration of H^+ in the sample is then

$$\begin{aligned}[H^+] &= \text{moles of } H^+ / \text{volume} \\ &= 1.3 \times 10^{-6} \text{ mole} / 1.3 \text{ liter} = 10^{-6} \text{ M.}\end{aligned}$$

The pH is defined as $pH = -\log [H^+]$, hence

$$pH = -\log [H^+] = -\log (10^{-6}) = -(-6) = 6,$$


which is the pH of normal urine.

A chemist wants to make up a 50 ml solution of pH 1.25 HCl. The HCl is labeled 11.6 M. How should she go about the preparation?

Solution: To answer this question, find how many moles of HCl exist in 50 ml of $pH = 1.25$. Once this is determined, one can calculate the volume of concentrated HCl needed. The total volume of water to be added will be the difference between the total volume of the solution and this amount of concentrated HCl. Thus, proceed as follows:

One can find the number of moles of HCl present from the pH. $pH = -\log [H^+]$. As given, $pH = 1.25$.

Thus, $1.25 = -\log [H^+]$. Solving, $[H^+] = 5.65 \times 10^{-2} \text{ M}$. The concentration of H^+ must be equal to the concentration of Cl^- and HCl, since



The solution is to have a volume of 50 ml or 0.05 liters. (Note: 1000 ml = 1 liter.) M = molarity = moles/liters.

Thus, the number of moles is equal to the volume in liters \times molarity or

$$(5.65 \times 10^{-2} \text{ M})(0.05 \text{ l}) = 2.82 \times 10^{-3} \text{ moles HCl.}$$

It is given that the molarity of the solution is 11.6 M. One needs 2.82×10^{-3} moles. Recalling that molarity = moles/liter, one finds that

$$\frac{2.82 \times 10^{-3} \text{ moles}}{11.6 \text{ M}} = 2.44 \times 10^{-4} \text{ l} = 0.24 \text{ ml}$$

of concentrated HCl that is required. If the total volume of the solution is to be 50 ml, then, add $(50 - .24 =)$ 49.76 ml of H_2O to 0.24 ml of the concentrated HCl.

• PROBLEM 346

Both HCl and NaOH are strong electrolytes. What is the pH of the solution formed by adding 40 ml of 0.10 M NaOH to 10 ml of 0.45 M HCl?

Solution: We will solve this problem by considering the number of moles of H_3O^+ and of OH^- formed by the complete dissociation of HCl and NaOH, respectively.


If we assume that HCl and NaOH dissociate completely, then the concentration of H_3O^+ in the HCl solution is equal to the initial concentration of HCl, or $[\text{H}_3\text{O}^+] = 0.45 \text{ M}$, and the concentration of OH^- in the NaOH solution is equal to the initial concentration of NaOH, or $[\text{OH}^-] = 0.10 \text{ M}$. Since moles = concentration \times volume, the number of moles of H_3O^+ in 10 ml of the acid solution is

$$\begin{aligned} \text{moles } \text{H}_3\text{O}^+ &= [\text{H}_3\text{O}^+] \times \text{volume} = 0.45 \text{ M} \times 10 \text{ ml} \\ &= 0.45 \text{ M} \times 0.01 \text{ liter} \\ &= .0045 \text{ mole} = 4.5 \times 10^{-3} \text{ mole.} \end{aligned}$$

Similarly, the number of moles of OH^- in 40 ml of the basic solution is

$$\begin{aligned} \text{moles } \text{OH}^- &= [\text{OH}^-] \times \text{volume} = 0.10 \text{ M} \times 40 \text{ ml} \\ &= 0.10 \text{ M} \times 0.04 \text{ liter} \\ &= 0.004 \text{ mole} = 4 \times 10^{-3} \text{ mole} \end{aligned}$$

H_3O^+ and OH^- neutralize each other according to the reaction



Ignoring the dissociation of water, we can assume that

this reaction is complete. Hence, the 4×10^{-3} moles of OH^- will be neutralized by 4×10^{-3} moles of H_3O^+ , leaving $4.5 \times 10^{-3} - 4.0 \times 10^{-3} = 0.5 \times 10^{-3} = 5 \times 10^{-4}$ mole of H_3O^+ remaining. Thus, when the two solutions are mixed, no OH^- remains and 5×10^{-4} mole of H_3O^+ remains. Since the final volume of the solution is $40 \text{ ml} + 10 \text{ ml} = 50 \text{ ml} = 0.05 \text{ liter} = 5 \times 10^{-2} \text{ liter}$, the concentration of H_3O^+ is

$$[\text{H}_3\text{O}^+] = 5 \times 10^{-4} \text{ mole} / 5 \times 10^{-2} \text{ liter} = 10^{-2} \text{ mole/liter.}$$

The pH is defined as $\text{pH} = -\log [\text{H}_3\text{O}^+]$, hence

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log (10^{-2}) = -(-2) = 2.$$

THE IONIZATION CONSTANT

• PROBLEM 347

The ionization constant for acetic acid is 1.8×10^{-5} .

- Calculate the concentration of H^+ ions in a 0.10 molar solution of acetic acid.
- Calculate the concentration of H^+ ions in a 0.10 molar solution of acetic acid in which the concentration of acetate ions has been increased to 1.0 molar by addition of sodium acetate.

Solution: The ionization constant (K_a) is defined as the concentration of H^+ ions times the concentration of the conjugate base ions of a given acid divided by the concentration of unionized acid. For an acid, HA,

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]},$$

where K_a is the ionization constant, $[\text{H}^+]$ is the concentration of H^+ ions, $[\text{A}^-]$ is the concentration of the conjugate base ions and $[\text{HA}]$ is the concentration of unionized acid. The K_a for acetic acid is stated as

$$K_a = \frac{[\text{H}^+][\text{acetate ion}]}{[\text{acetic acid}]} = 1.8 \times 10^{-5}.$$

The chemical formula for acetic acid is $\text{HC}_2\text{H}_3\text{O}_2$. When it is ionized, one H^+ is formed and one $\text{C}_2\text{H}_3\text{O}_2^-$ (acetate) is formed, thus the concentration of H^+ equals the concentration of $\text{C}_2\text{H}_3\text{O}_2^-$.

$$[\text{H}^+] = [\text{C}_2\text{H}_3\text{O}_2^-].$$

The concentration of unionized acid is decreased when ionization occurs. The new concentration is equal to the concentration of H^+ subtracted from the concentration of unionized acid.

$$[\text{HC}_2\text{H}_3\text{O}_2] = 0.10 - [\text{H}^+].$$

Since $[\text{H}^+]$ is small relative to 0.10, one may assume that 0.10 -

$[H^+]$ is approximately equal to 0.10.

Using this assumption, and the fact that $[H^+] = [C_2H_3O^-]$, K_a can be rewritten as

$$K_a = \frac{[H^+][H^+]}{0.10} = 1.8 \times 10^{-5}.$$

Solving for the concentration of H^+ :

$$[H^+]^2 = (1.0 \times 10^{-1})(1.8 \times 10^{-5}) = 1.8 \times 10^{-6}$$

$$[H^+] = \sqrt{1.8 \times 10^{-6}} = 1.3 \times 10^{-3}.$$

The concentration of H^+ is thus $1.3 \times 10^{-3} M$.

b) When the acetate concentration is increased, the concentration of H^+ is lowered to maintain the same K_a . The K_a for acetic acid is stated as

$$K_a = \frac{[H^+][C_2H_3O^-]}{[HC_2H_3O]} = 1.8 \times 10^{-5}$$

As previously shown for acetic acid equilibria in a solution of 0.10 molar acid, the concentration of acid after ionization is

$$[HC_2H_3O] = 0.10 - [H^+].$$

Because $[H^+]$ is very small compared to 0.10, $0.10 - [H^+] \approx 0.10$ and $[HC_2H_3O] = 0.10$.

In this problem, we are told that the concentration of acetate is held constant at 1.0 molar by addition of sodium acetate. Because one now knows the concentrations of the acetate and the acid, the concentration of H^+ can be found.

$$\frac{[H^+][C_2H_3O^-]}{[HC_2H_3O]} = 1.8 \times 10^{-5}$$

$$\frac{[H^+][1.0]}{[0.10]} = 1.8 \times 10^{-5}$$

$$[H^+] = 1.8 \times 10^{-6}.$$

• PROBLEM 348

Find the hydronium ion concentration of .1M HOAC (acetic acid) solution. Assume $k_a = 1.75 \times 10^{-5}$ for acetic acid.

Solution: You want to represent the equilibrium constant expression for the reaction, which necessitates a balanced equation. After writing the expression, you want to express the concentrations in terms of the same variables and solve for it. Begin by writing the balanced equation for the reaction of acetic acid in water. The acid will donate a proton (H^+) to the only available base, H_2O . Thus, $HOAC + H_2O \rightleftharpoons H_3O^+ + OAC^-$. $[H_3O^+]$, the hydronium concentration, is the quantity you are looking for. The equilibrium constant expression measures the ratio of the concentrations of the products to the reactants, each raised to the power of their

respective coefficients in the chemical equation. Thus, the constant, $K_a = \frac{[\text{OAC}^-][\text{H}_3\text{O}^+]}{[\text{HOAC}]}$. Note: H_2O is omitted,

since it is considered a constant. $K_a = 1.75 \times 10^{-5}$. Equating, $\frac{[\text{OAC}^-][\text{H}_3\text{O}^+]}{[\text{HOAC}]} = 1.75 \times 10^{-5}$. Let $x = \text{concentration of } \text{H}_3\text{O}^+$.

According to the reaction, $[\text{H}_3\text{O}^+] = [\text{OAC}^-]$, thus, $x = \text{concentration of } [\text{OAC}^-]$, also. If the initial concentration of HOAC is .1 and x moles of $[\text{H}_3\text{O}^+]$ are formed, then you have $(.1-x)$ moles/liter of HOAC left. Substituting these variables into the equilibrium constant expression, you have

$$\frac{x^2}{.1-x} = 1.75 \times 10^{-5}.$$

Solving, $x = [\text{H}_3\text{O}^+] = 0.0013\text{M}$

• PROBLEM 349

The ionization constant for NH_4OH is 1.8×10^{-5} .

- (a) Calculate the concentration of OH^- ions in a 1.0 molar solution of NH_4OH .

Solution: The ionization constant (K_b) is defined as the concentration of OH^- ions times the concentration of the conjugate acid ions of a given base divided by the concentration of unionized base. For a base, BA,

$$K_b = \frac{[\text{B}^-][\text{A}^+]}{[\text{BA}]},$$

where K_b is the ionization constant, $[\text{B}^-]$ is the concentration of ionized base ions, $[\text{A}^+]$ is the concentration of the conjugate acid, and $[\text{BA}]$ is the concentration of unionized base. The K_b for NH_4OH is stated as

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} = 1.8 \times 10^{-5}$$

When NH_4OH is ionized, one NH_4^+ ion is formed and one OH^- ion is formed,



Thus, the concentrations of each ion are equal.

$$[\text{NH}_4^+] = [\text{OH}^-]$$

The concentration of unionized base is decreased when ionization occurs. The new concentration is equal to the concentration of OH^- subtracted from the concentration of NH_4OH .

$$[\text{NH}_4\text{OH}] = 1.0 - [\text{OH}^-]$$

Since $[\text{OH}^-]$ is small relative to 1.0, one may assume that $1.0 - [\text{OH}^-]$ is approximately equal to 1.0

$$[\text{NH}_4\text{OH}] = 1.0 - [\text{OH}^-] \approx 1.0$$

Using this assumption, and the fact that $[\text{OH}^-] = [\text{NH}_4^+]$, K_b can be rewritten as

$$K_b = \frac{[\text{OH}^-][\text{OH}^-]}{1.0} = 1.8 \times 10^{-5}$$

Solving for $[\text{OH}^-]$:

$$\frac{[\text{OH}^-][\text{OH}^-]}{1.0} = 1.8 \times 10^{-5}$$

$$[\text{OH}^-]^2 = 1.8 \times 10^{-5}$$

$$[\text{OH}^-] = \sqrt{1.8 \times 10^{-5}} = 4.2 \times 10^{-3}$$

• PROBLEM 350

Find the degree of ionization of 0.05 M NH_3 in a solution of pH 11.0. $K_b = 1.76 \times 10^{-5}$.


Solution: The degree of ionization is the fraction of the total acid or base present (α) that ionizes. Namely,

$$\alpha = \frac{x}{c},$$

where x is the number of moles of acid or base that dissociate and c is the original number of moles present.

To solve this problem, let it first be stated that NH_3 is a base and that $[\text{OH}^-]$ can be calculated from the pH value.

The reaction equation is



After obtaining $[\text{OH}^-]$, $[\text{NH}_4^+]$ will be the only unknown in the equilibrium constant equation, which is

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.76 \times 10^{-5}.$$

To find $[\text{OH}^-]$, note that $\text{pOH} = -\log [\text{OH}^-]$ and that $\text{pH} + \text{pOH} = 14$. Substituting the given value of pH,

$$\text{pOH} = 14 - \text{pH} = 14 - 11 = 3.$$

Thus, $[\text{OH}^-] = -\log 3 = 1 \times 10^{-3}$.

Next, remember that for each x moles of NH_3 that dissociates, x moles of NH_4^+ will form. The problem becomes clearer if one observes what is happening from the table below.



Before Reaction: 0.05 0 0 0

After Reaction: 0.05- x x 1×10^{-3}

Upon substituting these values into the K_b equation,

one obtains:

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.76 \times 10^{-5} = \frac{x(10^{-3})}{0.05 - x}.$$

Solving for $[\text{NH}_4^+]$,


$$x = [\text{NH}_4^+] = 8.65 \times 10^{-4}.$$

$[\text{NH}_4^+]$ also represents the unknown value in the degree of ionization equation, since its concentration must be the amount of NH_3 that was ionized. Thus,

$$\alpha = \frac{8.65 \times 10^{-4}}{0.05} = 1.73 \times 10^{-2}.$$

• PROBLEM 351

Given K_i for acetic acid is 1.8×10^{-5} , calculate the percentage of ionization of 0.5 M acetic acid. The dissociation reaction is



Solution: K_i is the ionization constant and indicates to what degree acids and bases will dissociate in solution. Acetic acid is a weak acid because K_i is so small. The larger the value of K_i , the greater the % dissociation and the stronger the acid. A useful range of values for K_i is given below.

<u>Strength</u>	<u>Range</u>
Very strong	greater than 1×10^3
strong	1×10^3 to 1×10^{-2}
weak	1×10^{-2} to 1×10^{-7}
very weak	less than 1×10^{-7}

K_i is calculated from the ratio of products to reactants, each raised to the power of its coefficient in the reaction equation;

$$K_i = \frac{[\text{products}]^{\text{reaction moles (or coefficient)}}}{[\text{reactants}]^{\text{reaction moles (or coefficient)}}}$$

From the chemical reaction,

$$K_i = \frac{[\text{H}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]} = 1.8 \times 10^{-5}$$

To solve this problem, one uses the following method. Before the reaction begins, no dissociation occurs, so that only acetic acid is present, and in its full concentration of 0.5 M. After dissociation occurs, b number of moles/liter has dissociated to the products in equal amounts. Or b moles/liter of each H^+ and $C_2H_3O_2^-$ are produced since the coefficients of the reaction indicate that they are formed in equimolar amounts. If b moles/liter dissociate, then there are $0.5 - b$ moles/liter of acetic acid left.

This can be summarized as follows:

Before:	0.5 moles.	0	0
After:	$0.5 - b$	b	b

This gives a K_i of,

$$K_i = \frac{[b][b]}{[0.5 - b]} = 1.8 \times 10^{-5}.$$

Since K_i is so small, very few moles of acetic acid dissociate, which causes b to be insignificantly small. Therefore, $0.5 - b$ is approximately equal to 0.5, giving

$$K_i = \frac{[b][b]}{[0.5]} = 1.8 \times 10^{-5}.$$

Solving for b, one obtains:

$$b^2 = 9.0 \times 10^{-6}$$

$$b = 3.0 \times 10^{-3} \text{ mole of acetic acid ionized.}$$

Using the following equation to find % ionization:

$$\begin{aligned} \% \text{ ionization} &= \frac{\text{number of moles ionized}}{\text{original number of moles of acetic acid}} \times 100 \\ &= \frac{3.0 \times 10^{-3}}{0.5} \times 100 = 0.60 \%. \end{aligned}$$

• PROBLEM 352

A solution of 10.0 g of HF in 500 g H_2O freezes at $-1.98^\circ C$. Calculate the degree of ionization of HF. (M.W. HF = 20.0 The Freezing point depression of H_2O is 1.86° .)


Solution: When an acid is dissolved in solution, it dissociates into ions. For example, when one mole of HF is dissolved in H_2O , one mole of H^+ and one mole of F^- are present after complete dissociation. The freezing

point of a solution is dependent on the number of particles in the solution, which means that the freezing point of a solution will be lowered more by a compound which ionizes than by the same amount of a compound which does not ionize. The degree of ionization of a compound is a measure of what percent of the compound is ionized when it is placed in a particular solution.

In this problem, one is told that when 10 g of HF is added to 500 g of H_2O the original freezing point of the water (0°) is lowered to $-1.98^\circ C$. The freezing point depression is related to the concentration of particles in the solution by the statement

$$\text{freezing point depression} = \text{molality} \times \text{freezing point constant}$$

The freezing point constant of water is 1.86° . This means that one mole of a substance (except those substances which ionize) dissolved in 1000 g of water lowers the freezing point 1.86° . Since HF ionizes, we cannot use its molality in this equation. You can use the effective molality, which is the sum of the molalities of H^+ , F^- and HF. The molality of H^+ and F^- will be equal to the degree of ionization of the HF. The concentrations of H^+ and F^- will be equal because, when HF ionizes, one H^+ and one F^- will be formed. This reaction is given by the equation



To solve for the degree of ionization of HF, one must: (a) find the molality of HF as if it were not ionized, (b) define a variable for the molalities of H^+ and F^- ; here, x will be used, (c) find the molality of HF after ionization is taken into account, (d) find the effective molality of the species, (e) use the effective molality in the freezing point depression equation to solve for x , the molality of H^+ and F^- .

(a) The molality of HF before ionization.

The molality is defined as the number of moles of solute in 1000 g of solvent. The number of moles of HF can be found by dividing the number of grams present by the molecular weight.

$$\text{number of moles} = \frac{\text{number of grams}}{\text{molecular weight}}$$

$$\text{moles of HF} = \frac{10.0 \text{ g}}{20.0 \text{ g/mole}} = 0.5 \text{ moles}$$

The molality can now be found by dividing the number of moles of HF by the number of kg of water present.

$$\text{molality} = \frac{\text{no. of moles of HF}}{\text{no. of kg of } H_2O}$$

$$\text{molality} = \frac{0.5 \text{ moles}}{0.500 \text{ kg}} = 1.0 \text{ moles/kg}$$

(b) $x = \text{molality of } H^+ = \text{molality of } F^-$

(c) After ionization of x molal of H^+ and F^- , the molality of HF will be $1.0 - x$.

(e) The effective molality of the species will be equal to the sum of the molalities of all of the species present.

The effective molality = molality of H^+ + molality of F^- + molality of HF

$$\text{effective molality} = x + x + (1.0 - x) = 1.0 + x.$$

To solve for x , the concentration of H^+ and F^- , the effective molality of the species will be used in the freezing point depression equation.

$$\text{freezing point depression} = \text{eff. molality} \times \text{freezing pt constant}$$

The freezing point depression in this case is 1.98° and the freezing point constant of water is 1.86° . Solving the equation

$$1.98^\circ = (1.0 + x) \times 1.86^\circ$$

$$1.0 + x = \frac{1.98^\circ}{1.86^\circ}$$

$$x = .06$$

The percent of ionization is the molality of H^+ and F^- divided by the molality of the unionized HF multiplied by 100.

$$\text{degree of ionization} = \frac{\text{molality of ion}}{\text{molality of unionized species}} \times 100$$

$$\text{degree of ionization} = \frac{0.06}{1.00} \times 100 = 6\%$$

Hence, the HF in this solution is 6 % ionized.

THE DISSOCIATION CONSTANT

• PROBLEM 353

If 1 mole of HCl and 1 mole of $NaC_2H_3O_2$ are mixed in enough water to make one liter of solution, what will be the concentrations of the species in the final equilibrium? $K_{diss} = 1.8 \times 10^{-5}$ for $NaC_2H_3O_2$.

Solution: To answer this question, you must consider what is happening at equilibrium. This necessitates defining $K_{\text{dissociation}}$, which is an equilibrium constant


HCl and $\text{NaC}_2\text{H}_3\text{O}_2$ are strong electrolytes, which means that, in solution, they are completely dissociated. You have, therefore, H^+ , Cl^- , Na^+ , and $\text{C}_2\text{H}_3\text{O}_2^-$ ions present in the solution. The Na^+ and Cl^- do not associate, and need not be considered. Thus, you must only consider the formation of $\text{HC}_2\text{H}_3\text{O}_2$ from H^+ and $\text{C}_2\text{H}_3\text{O}_2^-$. The equation for this reaction can be written



This reaction can proceed in both directions, an equilibrium exists, as the double arrow indicates. The equilibrium constant (K_{eq}) for this reaction is equal to

$$\frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{H}^+][\text{C}_2\text{H}_3\text{O}_2^-]}$$

$K_{\text{dissociation}}$ measures the equilibrium quantitatively. The dissociation reaction for $\text{HC}_2\text{H}_3\text{O}_2$ can be written



The dissociation constant,

$$K_{\text{diss}} = \frac{[\text{H}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]} = 1.8 \times 10^{-5}$$

By examination, you can see that K_{eq} for the association reaction is equal to $1/K_{\text{diss}}$. Thus,

$$K_{\text{eq}} = \frac{1}{K_{\text{diss}}} = \frac{1}{1.8 \times 10^{-5}} = \frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{H}^+][\text{C}_2\text{H}_3\text{O}_2^-]}$$

To rewrite into a more convenient form for solving, take the reciprocal of each side.

$$1.8 \times 10^{-5} = \frac{[\text{H}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]}$$

The final concentrations of the species, the unknowns, will be those at the equilibrium. Let y be the concentration of $\text{HC}_2\text{H}_3\text{O}_2$ at equilibrium. The concentrations of both H^+ and $\text{C}_2\text{H}_3\text{O}_2^-$ can be represented by $1 - y$. Initially, you started with 1 mole/liter of each, therefore, each y mole/liter that associates to form $\text{HC}_2\text{H}_3\text{O}_2$ must be subtracted from the initial concentration. You can now substitute these variables into the expression for K_{diss} to obtain

$$\frac{(1 - y)(1 - y)}{y} = 1.8 \times 10^{-5}$$

Solving for y , using the quadratic formula, you obtain $y = .996$. Therefore, the concentrations of the species are

$$[H^+] = 1 - y = .004 \text{ M}$$


$$[C_2H_3O_2^-] = .004 \text{ M}$$

$$[HC_2H_3O_2] = .996 \text{ M.}$$

• PROBLEM 354

There exists a 0.5 M HF solution for which $K_{\text{diss}} = 6.71 \times 10^{-4}$. Determine how much dilution is necessary to double the percent dissociation of HF.

Solution: Percent dissociation means the ratio of $[H^+]$ to original $[HF]$ concentration times 100. To find the amount of dilution necessary to double the percent dissociation of an acid, first, establish what the percent dissociation is before dilution. This can be determined from K_{diss} , the equilibrium dissociation constants, which measures the ratio of products to reactants (i.e. their concentrations), each raised to the power of their coefficients in the balanced chemical equation. The general reaction for the dissociation of an acid, e.g., HA, is



for the dissociation reaction. This means that

$$K_{\text{diss}} = 6.71 \times 10^{-4} = \frac{[H_3O^+][F^-]}{[HF]}.$$

It is given that the initial concentration of HF is 0.5. Thus, to find percent dissociation, one needs to know $[H_3O^+]$. To find this, perform the following operations:

Let $x = [H_3O^+]$. Since H_3O^+ and F^- are formed in equimolar amounts, as can be seen in the chemical reaction, $[H_3O^+] = [F^-] = x$. If the initial $[HF] = 0.5$, and x moles/liter dissociate to give $[H_3O^+]$ and $[F^-]$, then, at equilibrium, $[HF] = 0.5 - x$. Substituting these values:

$$\frac{x \cdot x}{0.5 - x} = 6.71 \times 10^{-4}.$$

Solving for x , using the quadratic formula, $x = 0.018 = [H_3O^+]$. Thus, percentage dissociation becomes

$$\frac{0.018}{0.5} \times 100 = 3.6 \% \text{ HF dissociated.}$$

Then, to determine the final answer, 7.2 % HF dissociation is needed when one dilutes. Recall that $[H^+] = [F^-]$. At 7.2 % dissociation,

$$\frac{[F^-]}{[HF]} = \frac{0.072}{1 - .072} = \frac{0.072}{0.928}.$$

From the K_{diss} expression, one has

$$K_{\text{diss}} = 6.71 \times 10^{-4} = \frac{[H^+][F^-]}{[HF]} \quad \text{or}$$

$$\frac{6.71 \times 10^{-4}}{[H^+]} = \frac{[F^-]}{[HF]}.$$

$$\frac{[F^-]}{[HF]} = \frac{0.072}{0.928}, \text{ when percent dissociation is}$$

doubled to 7.2 %. After substitution,

$$\frac{6.71 \times 10^{-4}}{[H^+]} = \frac{0.072}{0.928}.$$

Solving for $[H^+]$, $[H^+] = 8.65 \times 10^{-3} = [F^-]$. Substituting these actual molar concentrations into the equilibrium constant expression, one obtains

$$\frac{6.71 \times 10^{-4}}{8.65 \times 10^{-3}} = \frac{8.65 \times 10^{-3}}{[HF]}$$

Solving: $[HF] = 0.112$ at equilibrium

$$[H^+] = 8.65 \times 10^{-3}.$$

Before dissociation $[HF]$ = amount at equilibrium plus amount dissociated = $8.65 \times 10^{-3} + 0.112 = 0.121$ M. Thus, when the percent dissociation is doubled, the initial amount equals 0.121 M for $[HF]$. The $[HF]$, when the percent dissociation was unchanged, was 0.5 M (given). Therefore, dilute by factor $0.500/0.121 = 4.13$. Remember that concentration or molarity is a parameter of volume, actually $M = \text{moles/liter}$. One is going from 0.5 M to 0.121 M, which means volume must be increased by a certain factor. When one dilutes a solution, one is adding volume to it and, dividing the solution. This account for using division to obtain the factor of dilution, once initial concentrations are known.

A chemist wants the percent dissociation of $\text{HC}_2\text{H}_3\text{O}_2$ to be 1 %. If the K_{diss} of $\text{HC}_2\text{H}_3\text{O}_2$ is 1.8×10^{-5} , what concentration of $\text{HC}_2\text{H}_3\text{O}_2$ is required?

Solution: To answer this question, consider the dissociation of an acid in H_2O . Then write an equilibrium constant expression which relates the equilibrium constant for dissociation, K_{diss} , to the ratio of the concentrations of products to reactants, each raised to the power of its coefficients in the reaction. In general, for an acid, there exists the following reaction for dissociation:



where HA is the acid. For this reaction,



The equilibrium constant expression is

$$K_{\text{diss}} = 1.8 \times 10^{-5} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]}$$

One wants to find $[\text{HC}_2\text{H}_3\text{O}_2]$ with the knowledge that the percent dissociation is 1 % (note: exclude water from the expression, as it is assumed to be constant). If the percent dissociation is 1 %, then

$$[\text{H}_3\text{O}^+]/[\text{HC}_2\text{H}_3\text{O}_2] = 1/99.$$

The H^+ can only come from $\text{HC}_2\text{H}_3\text{O}_2$. If 1 % of H^+ exists, then 100% (initial percentage) - 1% = 99% of $\text{HC}_2\text{H}_3\text{O}_2$ must be left at equilibrium. Thus,

$$1.8 \times 10^{-5} = \frac{1}{99} [\text{C}_2\text{H}_3\text{O}_2^-]. \text{ Solving for } [\text{C}_2\text{H}_3\text{O}_2^-];$$

$$[\text{C}_2\text{H}_3\text{O}_2^-] = 1.78 \times 10^{-3} \text{ M.}$$

This concentration of $\text{C}_2\text{H}_3\text{O}_2^-$ must equal the concentration of H_3O^+ , since they are formed in equimolar amounts. One can substitute into the equation

$$1.8 \times 10^{-5} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]}$$


$$\text{to obtain } \frac{(1.78 \times 10^{-3})(1.78 \times 10^{-3})}{[\text{HC}_2\text{H}_3\text{O}_2]} = 1.8 \times 10^{-5}.$$

Solving for $[\text{HC}_2\text{H}_3\text{O}_2]$, $1.76 \times 10^{-1} \text{ M.}$


Given a solution of 1.00M $\text{HC}_2\text{H}_3\text{O}_2$, what is the concentration of all solute species? What is the percentage of acid that dissociated? Assume $K_{\text{diss}} = 1.8 \times 10^{-5}$.

Solution: Determine the equilibrium equation, write an equilibrium constant expression, and substitute the concentrations of the species into the expression.

By definition, an acid is a substance which donates protons (H^+). Since water is the only other species in the solution, it must act as a base; it receives protons.



There is a constant called $K_{\text{dissociation}}$, which measures the extent of the $\text{HC}_2\text{H}_3\text{O}_2$ donation of protons. For the general reaction,



For the reaction in this problem, you have

$$\frac{[\text{C}_2\text{H}_3\text{O}_2^-][\text{H}_3\text{O}^+]}{[\text{HC}_2\text{H}_3\text{O}_2]} = K_{\text{diss}} = 1.8 \times 10^{-5}$$

You are asked to find these concentrations. Let x = the moles per liter of $\text{HC}_2\text{H}_3\text{O}_2$ that dissociated. From the chemical equation, you see that x must also be the concentration of H_3O^+ and $\text{C}_2\text{H}_3\text{O}_2^-$, since for each mole of $\text{HC}_2\text{H}_3\text{O}_2$ that dissociates, one mole of H_3O^+ and one mole of $\text{C}_2\text{H}_3\text{O}_2^-$ is produced. With this in mind, you can represent $[\text{HC}_2\text{H}_3\text{O}_2]$ as $1-x$, since you started with 1M of $\text{HC}_2\text{H}_3\text{O}_2$ and x moles per liter dissociate. You have left $1-x$ moles per liter. Now substituting, you obtain

$$\frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]} = 1.8 \times 10^{-5} = \frac{x \cdot x}{1-x}$$

Solving for x , you obtain $x = .0042$. Therefore, the concentrations of the species are

$$[\text{HC}_2\text{H}_3\text{O}_2] = 1 - .0042 = .9958\text{M}$$


$$[\text{H}_3\text{O}^+] = x = .0042\text{M}$$

$$[\text{C}_2\text{H}_3\text{O}_2^-] = x = .0042\text{M}$$

The percentage of acid that dissociated is, thus,

$$\frac{.0042}{1} \times 100 = .42\%$$

The density of liquid NH_3 = 0.68 g/ml. Liquid NH_3 dissociates according to the reaction



for which $K = 1.0 \times 10^{-33}$ at -33.4°C . Determine the concentrations of NH_4^+ and NH_2^- at equilibrium.

Solution: To solve this problem, write the equilibrium constant expression for the dissociation of liquid NH_3 . This equates the equilibrium constant, K , to the ratio of the concentrations of products to reactants, each raised to the power of its coefficient in the chemical reaction.

$$K = 1.0 \times 10^{-33} = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2}$$

One is asked to find $[\text{NH}_4^+]$ and $[\text{NH}_2^-]$. To do this, one needs the initial concentration of the liquid ammonia, which can be found from the density. (density = mass/volume). It is given that density = 0.68 g/ml. The molecular weight of ammonia (NH_3) is 17 g/mole. For the correct units, convert 0.68 g/ml to 680 g/l. Thus concentration (moles/liter) = $(680 \text{ g/l}) / (17 \text{ g/mole}) = 40 \text{ molar } \text{NH}_3$. Let $x = [\text{NH}_4^+]$.

From the chemical reaction, it can be seen that NH_4^+ and NH_2^- are formed in equimolar amounts, which means $[\text{NH}_4^+] = [\text{NH}_2^-]$, so that $[\text{NH}_2^-] = x$. If the original $[\text{NH}_3] = 40 \text{ M}$, and x moles/liter form $[\text{NH}_4^+]$ and $[\text{NH}_2^-]$ each, then, at equilibrium, $[\text{NH}_3] = 40 - x$.

Substituting these values into the equilibrium constant expression, one obtains

$$1.0 \times 10^{-33} = \frac{x \cdot x}{(40 - x)^2} \quad \text{Solving for } x, \text{ using the}$$

quadratic formula:

$$x = 1.26 \times 10^{-15} = [\text{NH}_4^+] = [\text{NH}_2^-] \text{ at equilibrium.}$$

• PROBLEM 358

Given the equilibrium: $\text{ClCH}_2\text{COOH} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{ClCH}_2\text{COO}^-$ exists at 25°C . $K_a = 1.35 \times 10^{-3}$. (a) Determine the H_3O^+ concentration for a 0.1 M solution of monochloroacetic acid (ClCH_2COOH) in water (H_2O). (b) Can one make the assumption that the dissociated acid is negligible with respect to the undissociated acid? (c) Calculate to what degree this solution is more acidic than 1.0 M acetic acid ($K_a = 1.8 \times 10^{-5}$ at 25°C).

Solution: The concentration of H_3O^+ can be calculated through the equilibrium constant expression. This is a measure of the ratio between the concentrations of products to reactants, each raised to the power of their respective

coefficients in the chemical equation.

(a) For this reaction, the equilibrium constant,

$$K_a = \frac{[\text{ClCH}_2\text{COO}^-][\text{H}_3\text{O}^+]}{[\text{ClCH}_2\text{COOH}]} . \quad \text{Because } K_a = 1.35 \times 10^{-3},$$

$$1.35 \times 10^{-3} = \frac{[\text{ClCH}_2\text{COO}^-][\text{H}_3\text{O}^+]}{[\text{ClCH}_2\text{COOH}]} .$$

Letting $x = [\text{H}_3\text{O}^+]$ at equilibrium, then $\text{ClCH}_2\text{COO}^- = x$ since the reaction shows that the products are formed in equimolar amounts. The original concentration of the $\text{ClCH}_2\text{COOH} = 0.1 \text{ M}$.

If x moles of H_3O^+ and x moles of $\text{ClCH}_2\text{COO}^-$ form, then, at equilibrium, there are $0.1 - x$ moles of $\text{ClCH}_2\text{COO}^-$ left. Substituting these variables into the equilibrium constant expression:

$$\frac{x \cdot x}{(0.1 - x)} = 1.35 \times 10^{-3} .$$

Solving for x , $x = 0.0110 \text{ M}$.

(b) If the dissociated acid is negligible with respect to the amount left undissociated, then $0.1 - x$ (from previous part) is equal to 0.1 . This means that

$$\frac{x \cdot x}{0.1} = 1.35 \times 10^{-3} .$$

Solving for x ; $x = 0.0116 \text{ M} = [\text{H}_3\text{O}^+]$. This means that when you make this assumption, $[\text{H}_3\text{O}^+] = 0.0116 \text{ M}$ instead of 0.0110 M , the percent error is over 5 %. The error is too high and, therefore, this assumption cannot be made.


To answer (c), compare the $[\text{H}_3\text{O}^+]$ in acetic acid solution of 1.0 M with that of the $[\text{H}_3\text{O}^+]$ calculated in part (a). $[\text{H}_3\text{O}^+]$ for acetic acid is calculated by using the same procedure as in part (a), except that $K_a = 1.8 \times 10^{-5}$ and the dissociation equation is



After the calculations, $[\text{H}_3\text{O}^+] = 0.00133 \text{ M}$. For acetic acid $[\text{H}_3\text{O}^+] = 0.00133 \text{ M}$ and for ClCH_2COOH , $[\text{H}_3\text{O}^+] = 0.0110 \text{ M}$. Thus, the ClCH_2COOH has 8.27 times the H_3O^+ concentration of CH_3COOH (acetic acid), and is therefore more acidic. This answer is found by dividing $[\text{H}_3\text{O}^+]$ dissociated from ClCH_2COOH by $[\text{H}_3\text{O}^+]$ dissociated from CH_3COOH .

$$\frac{0.0110 \text{ M}}{0.00133 \text{ M}} = 8.27 .$$

Sulfuryl chloride decomposes according to the equation,



Given that at 300°K and 1 atm pressure, the degree of dissociation is .121, find the equilibrium constant, K_p . What will be the degree of dissociation when pressure = 10 atm?

Solution: To find the equilibrium constant, K_p , one uses the expression for K_p . This is the product of the pressures of the products divided by pressure of the reactant. These pressures are brought to the powers of their coefficients. From this, employ Dalton's law concerning partial pressures to help solve for K_p .

At $p = 10$ atm, assume K_p is pressure-independent because of ideal gas behavior. This, then, allows the determination of the degree of dissociation from the K_p expression.

Thus, one commences by the definition of K_p .

$$K_p = \frac{P_{\text{SO}_2} P_{\text{Cl}_2}}{P_{\text{SO}_2\text{Cl}_2}}$$

To find K_p , evaluate these partial pressures. To do this, use Raoult's law, which states partial pressure of gas = mole fraction of gas in mixture times total pressure, i.e., $p = X_{pT}$. The p_T is given as 1 atm. Thus, the key is to determine or represent the mole fraction X . Let "a" = the degree of dissociation of SO_2Cl_2 . From the chemical equation, it is seen that SO_2 and Cl_2 are formed in equimolar amounts. Thus, at equilibrium there are "a" moles of both SO_2 and Cl_2 . Starting with 1 mole of SO_2Cl_2 and "a" moles dissociated, then, at equilibrium, there remain $1 - a$ moles. In the mixture, therefore, total number of moles = $(1-a) + a + a = (1+a)$. Thus, $P_{\text{SO}_2\text{Cl}_2} = X_{\text{SO}_2\text{Cl}_2} P_T = \left(\frac{1-a}{1+a}\right) P_T$,

$P_{\text{SO}_2} = P_{\text{Cl}_2} = \frac{a}{1+a} P_T$. Therefore,

$$K_p = \frac{P_{\text{SO}_2} P_{\text{Cl}_2}}{P_{\text{SO}_2\text{Cl}_2}} = \frac{\left(\frac{a}{1+a}\right) P_T \left(\frac{a}{1+a}\right) P_T}{\left(\frac{1-a}{1+a}\right) P_T}$$

This can be simplified to

$$K_p = \frac{a^2}{1-a^2} P_T .$$

Given the degree of dissociation, a, is equal to .121 and $P_T = 1$ atm; K_p therefore, equal

$$\frac{(.121)^2}{1-(.121)^2} (1) = 1.49 \times 10^{-2} .$$

At 10 atm, then,

$$K_p = \frac{a^2}{1-a^2} (10) = 1.49 \times 10^{-2}.$$


Solving for a , $a = .0386$.

THE HYDROLYSIS CONSTANT

• PROBLEM 360

If the hydrolysis constant of Al^{3+} is 1.4×10^{-5} , what is the concentration of H_3O^+ in 0.1 M AlCl_3 ?

Solution: Hydrolysis refers to the action of the salts of weak acids and bases with water to form acidic or basic solutions. Consequently, to answer this question, write out the reaction, which illustrates this hydrolysis, and write out an equilibrium constant expression. From this, the concentration of H_3O^+ can be defined. The net hydrolysis reaction is



Water is excluded in this expression since it is considered as a constant. Let x = the moles/liter of $[\text{H}_3\text{O}^+]$. Since H_3O^+ and AlOH^{2+} are formed in equal mole amounts, the concentration of $[\text{AlOH}^{2+}]$ can also be represented by x . If one starts with 0.1 M of Al^{3+} , and x moles/liter of it forms H_3O^+ (and AlOH^{2+}), one is left with $0.1 - x$ at equilibrium. Substituting these representations into the K_{hyd} expression,

$$\frac{x \cdot x}{0.1 - x} = 1.4 \times 10^{-5}.$$

If one solves for x , the answer is $x = 1.2 \times 10^{-3}$ M, which equals $[\text{H}_3\text{O}^+]$.

• PROBLEM 361

Calculate the hydrolysis constants of the ammonium and cyanide ions, assuming $K_w = 1 \times 10^{-14}$ and $K_a = 4.93 \times 10^{-10}$

for HCN and $K_b = 1.77 \times 10^{-5}$ for NH_3 . For each, determine the percent hydrolysis in a .1M solution.

Solution: To find the hydrolysis constant, you must know what it defines. Hydrolysis is the process whereby an acid or base is regenerated from its salt by the action of water. The hydrolysis constant measures the extent of this process. Quantitatively, it is defined as being equal to $\frac{K_w}{K_a \text{ or } K_b}$, where K_w = the equilibrium constant

for the autodissociation of water, K_a = dissociation of acid, and K_b = dissociation of base. You are given K_w , K_a , and K_b . Thus, the hydrolysis constants can be easily found by substitution. Let K_h = hydrolysis constant.

$$\text{For cyanide ion: } K_h = \frac{K_w}{K_a} = \frac{1 \times 10^{-14}}{4.93 \times 10^{-10}} = 2.02 \times 10^{-5}$$

$$\text{For ammonium ion: } K_h = \frac{K_w}{K_b} = \frac{1 \times 10^{-14}}{1.77 \times 10^{-5}} = 5.64 \times 10^{-10}$$

To find the percent hydrolysis in a .1M solution, write the hydrolysis reaction and express the hydrolysis constant just calculated in those terms. After this, represent the concentrations of the hydrolysis products in terms of variables and solve. For cyanide ion: The hydrolysis reaction is $\text{CN}^- + \text{H}_2\text{O} \rightleftharpoons \text{HCN} + \text{OH}^-$. Therefore,

$$K_h = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]}. \text{ But you calculated that } K_h = 2.02 \times 10^{-5}.$$

$$\text{Equating, } 2.02 \times 10^{-5} = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]}.$$

You start with a .1M solution of CN^- . Let $x = [\text{HCN}]$ formed. Thus, $x = [\text{OH}^-]$ also, since they are formed in equimolar amounts. If x moles/liter of substance are formed from CN^- , then, at equilibrium you have .1-x moles/liter left.

$$\text{Substituting these values, } 2.02 \times 10^{-5} = \frac{x \cdot x}{.1 - x}$$

Solving, $x = 1.4 \times 10^{-3}\text{M}$. The percent is just 100 times $\frac{x}{.10\text{M}}$ since the initial concentration is .10M, so

that you have 1.4% hydrolysis in a .1M solution.

$$\text{For ammonium ion: The hydrolysis reaction is } \text{NH}_4^+ + \text{H}_2\text{O} \rightleftharpoons \text{NH}_3 + \text{H}_3\text{O}^+. \text{ } K_h \text{ for this reaction} = \frac{[\text{NH}_3][\text{H}_3\text{O}^+]}{[\text{NH}_4^+]}$$

The calculated $K_h = 5.6 \times 10^{-10}$.

$$\text{Equating, } 5.6 \times 10^{-10} = \frac{[\text{NH}_3][\text{H}_3\text{O}^+]}{[\text{NH}_4^+]}$$

From this point, you follow the same reasoning as was used with the cyanide.

Solving:

$$\text{Let } x = [\text{NH}_3]$$

$$5.6 \times 10^{-10} = \frac{(x)(x)}{(.1-x)}$$

$$x^2 = (5.6 \times 10^{-11}) - (5.6 \times 10^{-10})x$$

$$x^2 + (5.6 \times 10^{-10})x - 5.6 \times 10^{-11} = 0$$

Using the quadratic formula one can solve for x , where $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-5.6 \times 10^{-10} \pm \sqrt{(5.6 \times 10^{-10})^2 - 4(1)(-5.6 \times 10^{-11})}}{2(1)}$$

$$x = \frac{-5.6 \times 10^{-10} \pm \sqrt{2.24 \times 10^{-10}}}{2}$$

$$x = \frac{-5.6 \times 10^{-10} \pm 1.50 \times 10^{-5}}{2}$$

$$x = \frac{-1.50 \times 10^{-5}}{2} \text{ or } x = \frac{1.50 \times 10^{-5}}{2}$$

x cannot be negative, because concentration cannot be negative. Thus, $x = 7.5 \times 10^{-6}$

Solving for the percent:

$$\frac{7.5 \times 10^{-6}}{.1} \times 100\% = 7.5\%$$


Thus, you find that the percent hydrolysis is $7.5 \times 10^{-3}\%$.

• PROBLEM 362

Find an expression for the hydrolysis constant of the bicarbonate ion HCO_3^- , $K_h(\text{HCO}_3^-)$, using only the first dissociation constant for H_2CO_3 , $K_{a1}(\text{H}_2\text{CO}_3)$, and the water constant, K_w .

Solution: The solution to this problem is a direct application of the definitions of K_h , K_{a1} , and K_w .

The removal of one proton may be represented by



Hence, the first dissociation constant is

$$K_{a1}(\text{H}_2\text{CO}_3) = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$$

The water constant is given by

$$K_w = [\text{H}^+][\text{OH}^-].$$

Hydrolysis of HCO_3^- proceeds according to the reaction



We are trying to determine an expression for the hydrolysis constant $K_h(\text{HCO}_3^-)$ in terms of $K_{a1}(\text{H}_2\text{CO}_3)$ and K_w . By definition,

$$K_h(\text{HCO}_3^-) = \frac{[\text{H}_2\text{CO}_3][\text{OH}^-]}{[\text{HCO}_3^-]}.$$

$$\begin{aligned} \text{Hence, } K_h(\text{HCO}_3^-) &= \frac{[\text{H}_2\text{CO}_3][\text{OH}^-]}{[\text{HCO}_3^-]} \\ &= \frac{[\text{H}_2\text{CO}_3][\text{OH}^-]}{[\text{HCO}_3^-]} \times \frac{[\text{H}^+]}{[\text{H}^+]} \\ &= \frac{[\text{H}_2\text{CO}_3]}{[\text{H}^+][\text{HCO}_3^-]} \times [\text{H}^+][\text{OH}^-] \\ &= \frac{1}{K_{a1}(\text{H}_2\text{CO}_3)} \times K_w, \end{aligned}$$

$$\text{or, } K_h(\text{HCO}_3^-) = \frac{K_w}{K_{a1}(\text{H}_2\text{CO}_3)}.$$


• PROBLEM 363

What is the pH of a 1.0 M solution of the strong electrolyte sodium acetate? The dissociation constant of acetic acid is $K_a = 1.8 \times 10^{-5}$ mole/liter.

Solution: The first step is the determination of the hydrolysis constant for sodium acetate. From this we obtain the concentration of hydroxyl contributed by the hydrolysis of sodium acetate. The concentration of hydronium ion, and consequently the pH, is determined by using the water constant.

The dissociation of acetic acid (HAc) into hydronium

ions and acetate ions (Ac^-) may be represented by the equation




The dissociation constant for this reaction is

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{Ac}^-]}{[\text{HAc}]}$$

This constant and the water constant, $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$ mole²/liter² will be used to determine the hydrolysis constant for acetate.

Hydrolysis of acetate proceeds according to the following equation:



The hydrolysis constant is $K_h = [\text{HAc}][\text{OH}^-]/[\text{Ac}^-]$. This may be rewritten in terms of K_a and K_w as follows:

$$\begin{aligned} K_h &= \frac{[\text{HAc}][\text{OH}^-]}{[\text{Ac}^-]} = \frac{[\text{HAc}][\text{OH}^-]}{[\text{Ac}^-]} \times \frac{[\text{H}_3\text{O}^+]}{[\text{H}_3\text{O}^+]} \\ &= \frac{[\text{HAc}]}{[\text{Ac}^-][\text{H}_3\text{O}^+]} \times [\text{H}_3\text{O}^+][\text{OH}^-] = \frac{1}{K_a} \times K_w = \frac{K_w}{K_a}. \text{ Hence,} \\ K_h &= \frac{[\text{HAc}][\text{OH}^-]}{[\text{Ac}^-]} = \frac{K_w}{K_a} \\ &= \frac{10^{-14} \text{ mole}^2/\text{liter}^2}{1.8 \times 10^{-5} \text{ mole/liter}} = 5.6 \times 10^{-10} \text{ mole/liter.} \end{aligned}$$

Let the equilibrium concentration of HAc formed by the hydrolysis of acetate be x . Since one mole of OH^- is formed per mole of HAc formed, the equilibrium concentration of OH^- is also x . Furthermore, if we assume that sodium acetate dissociates completely, then the initial concentration of Ac^- is equal to the concentration of sodium acetate (1.0 M) and the equilibrium concentration of acetate is $1.0 - x$. Note that we have neglected the contribution to $[\text{OH}^-]$ from the hydrolysis of water.

Substituting these concentrations into the expression for K_h , we obtain

$$\begin{aligned} K_h &= 5.6 \times 10^{-10} \text{ mole/liter} = \frac{[\text{HAc}][\text{OH}^-]}{[\text{Ac}^-]} \\ &= \frac{x \cdot x}{1.0 - x} = \frac{x^2}{1.0 - x} \end{aligned}$$

To avoid use of the quadratic formula, we will assume that x is much smaller than 1.0 so that $1.0 - x \approx 1.0$. (This assumption will be justified later on in the solution). Hence, we obtain

$$5.6 \times 10^{-10} \text{ mole/liter} = \frac{x^2}{1.0 - x} \approx \frac{x^2}{1.0}$$

or $x = (1.0 \times 5.6 \times 10^{-10})^{1/2} = 2.4 \times 10^{-5} \text{ mole/liter}$.

Since $[\text{OH}^-] = x$, $[\text{OH}^-] = 2.4 \times 10^{-5} \text{ mole/liter}$. Hence, x is much smaller than 1.0, justifying our earlier assumption.

We will find $[\text{H}_3\text{O}^+]$ by use of the water constant, $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$, or,

$$\begin{aligned} [\text{H}_3\text{O}^+] &= \frac{K_w}{[\text{OH}^-]} = \frac{10^{-14} \text{ mole}^2/\text{liter}^2}{2.4 \times 10^{-5} \text{ mole/liter}} \\ &= 4.2 \times 10^{-9} \text{ mole/liter.} \end{aligned}$$

The pH is then

$$\begin{aligned} \text{pH} &= -\log [\text{H}_3\text{O}^+] = -\log (4.2 \times 10^{-9}) \\ &= -(-9.4) = 9.4. \end{aligned}$$

NEUTRALIZATION

• PROBLEM 364

Assuming complete neutralization, calculate the number of milliliters of 0.025 M H_3PO_4 required to neutralize 25 ml of 0.030 M $\text{Ca}(\text{OH})_2$.

Solution: This problem can be solved by two methods: mole method or equivalent method.

Mole Method

This method requires one to write out the balanced equation that illustrates the neutralization reaction. The balanced equation is $3\text{Ca}(\text{OH})_2 + 2\text{H}_3\text{PO}_4 \rightarrow \text{Ca}_3(\text{PO}_4)_2 + 6\text{H}_2\text{O}$. From this equation, one can see that 2 moles of H_3PO_4 react for every 3 moles of $\text{Ca}(\text{OH})_2$. This means that one must first calculate how many moles of $\text{Ca}(\text{OH})_2$ are involved. The molarity of the $\text{Ca}(\text{OH})_2$ is 0.030. (Molarity = no. of moles/liters.)

As given, the $\text{Ca}(\text{OH})_2$ solution is 25 ml or 0.025 liters. Therefore, the number of moles of $\text{Ca}(\text{OH})_2$ is

$(0.030)(0.025) = 0.00075$ moles. As the balanced equation indicates, the number of moles of H_3PO_4 is $2/3$ the moles of $Ca(OH)_2$ or $2/3 (0.00075) = 0.00050$ moles of H_3PO_4 .

From the definition of molarity for H_3PO_4 , one has

$$0.025 \text{ M} = \frac{0.00050 \text{ moles}}{\text{liters}} .$$

The molarity, 0.025, is given. Solving for liters, one obtains 0.020 liters or 20 ml. The key to solving this problem with the mole method, is to write a balanced equation, which will indicate the relative amounts of moles required for complete neutralization .

Equivalent Method

This method requires that one consider normality and the definition of an equivalent. An equivalent is defined as the molecular weight or mass of an acid or base that furnished one mole of protons (H^+) or hydroxyl (OH^-) ions. For example, the number of equivalents contained in a mole of H_2SO_4 is $98/2$ or 49. Since each mole of H_2SO_4 produces two protons, divide the molecular weight by 2.

The number of equivalents of an acid must equal that of the base in a neutralization reaction. Normality is defined as equivalents of solute per liter. In this problem, it is given that there are 25 ml of 0.03 $Ca(OH)_2$.

To solve the problem, determine how many equivalents are present. The number of moles of $Ca(OH)_2$ is $(0.025)(0.03)$ or 0.00075 from the definition of molarity. The molecular weight of $Ca(OH)_2$ is 74.08. Therefore, there are $(0.00075)(74.08)$ or 0.06 grams of $Ca(OH)_2$.

The number of equivalents per gram is $74.08/2$ since two OH^- can be produced. The number of equivalents is

$$\frac{(0.00075)(74.08)g}{\left(\frac{74.08}{2} \text{ g/equiv}\right)} = 0.0015 \text{ equiv of } Ca(OH)_2 .$$

This indicates that 0.0015 equivalents of H_3PO_4 are required. The molarity of H_3PO_4 is 0.025 M, which means its normality is 0.075 N, because there are 3 ionizable protons per mole. Recalling the definition of normality, there are

$$0.075 \text{ N} = 0.0015 \text{ equiv./liters}$$

The reason one knows that there is 0.0015 equiv in the H_3PO_4 present is because one knows that for the neutralization to occur the number of equivalents of acid must equal the number of equivalents of base. In this problem, one has already calculated that there are

0.0015 equiv of base present. Thus,

$$\text{volume} = 0.20 \text{ l or } 20 \text{ ml.}$$

• PROBLEM 365

A 50 ml solution of sulfuric acid was found to contain 0.490 g of H_2SO_4 . This solution was titrated against a sodium hydroxide solution of unknown concentration. 12.5 ml of the acid solution was required to neutralize 20.0 ml of the base. What is the concentration of the sodium hydroxide solution.

Solution: At the neutralization point, the number of equivalents of acid in the 12.5 ml volume is equal to the number of equivalents of base in the 20.0 ml volume. Since the normality is defined as the number of equivalents per liter of solution, the number of equivalents is equal to the normality times the volume, at the neutralization point we have

$$N_a V_a = N_b V_b$$

where N_a = normality of acid, V_a = volume of acid, N_b = normality of base, V_b = volume of base.

The normality of the 50.0 ml (0.050 l) sulfuric acid solution is

$$\begin{aligned} N_a &= \frac{\text{number of equivalents in } 0.050 \text{ l}}{0.05 \text{ l}} \\ &= \frac{\text{mass of acid/gram equivalent weight}}{0.05 \text{ l}} \end{aligned}$$

The gram equivalent weight of sulfuric acid is 49.0 g/equivalent, because there are 2 equiv per molecule. The MW of H_2SO_4 is 98 g/mole.

$$\begin{aligned} N_a &= \frac{\text{mass of acid/gram equivalent weight}}{0.05 \text{ l}} \\ &= \frac{0.490 \text{ g}/49.0 \text{ g/equivalent}}{0.05 \text{ l}} = 0.200 \text{ equivalent/l} \\ &= 0.200 \text{ N} \end{aligned}$$

The normality of the base is then found as follows:

$$N_a V_a = N_b V_b$$

$$N_b = \frac{N_a V_a}{V_b} = \frac{0.200 \text{ N} \times 12.5 \text{ ml}}{20.0 \text{ ml}} = 0.125 \text{ N}.$$

Therefore, the sodium hydroxide solution is 0.125 N, which because there is 1 ionizable OH^- in NaOH, is equal to 0.125 M.

• PROBLEM 366

Determine the molarity of an H_2SO_4 solution, 15.0 ml of which is completely neutralized by 26.5 ml of 0.100 M NaOH.

Solution: Acids and bases react with each other to produce salts and water. Such reactions are called neutralizations. To find the molarity of the neutralized acid solution, first write out the balanced reaction.



When neutralization is complete, there is no longer any acid or base left. It is given that 26.5 ml of 0.100 M NaOH is used. From this, one wants to compute how many moles of NaOH were present.

It is known from the neutralization reaction that the number of moles of acid must be $\frac{1}{2}$ the number of moles of base for a complete neutralization.

From the number of moles of base present, the number of moles of acid needed are determined. Since molarity = moles/liter and one is given the volume, the molarity can be calculated. Proceed as follows:

There exists 26.5 ml of 0.100 M NaOH. There are 1,000 ml in one liter. This means that 26.5 ml is equal to 0.0265 liters. Recalling the definition of molarity,

$$0.1 = \frac{\text{no. of moles of NaOH}}{0.0265 \text{ l}} \quad \text{Solving}$$

$$\text{moles} = 0.00265 \text{ of NaOH}.$$

This, then, must be twice the number of moles of acid. For the acid, therefore, molarity = $(0.00265) \frac{1}{2}$ /liters. One is told that the volume of the H_2SO_4 solution is 15 ml, or 0.015 liters (1000 ml = 1 liter). Molarity of the acid becomes

$$\frac{0.001325}{0.015} = 0.088 \text{ M}.$$

One of the two most common ingredients in medication designed for the relief of excess stomach acidity is aluminum hydroxide ($\text{Al}(\text{OH})_3$, formula weight = 78 g/mole). If a patient suffering from a duodenal ulcer displays a hydrochloric acid (HCl , formula weight = 36.5 g/mole), concentration of $80 \times 10^{-3} \text{ M}$ in his gastric juice and he produces 3 liters of gastric juice per day, how much medication containing 2.6 g $\text{Al}(\text{OH})_3$ per 100 ml of solution must he consume per day to neutralize the acid?

Solution: $\text{Al}(\text{OH})_3$ neutralizes HCl according to the following reaction:



When all the HCl has been neutralized, the number of equivalents of $\text{Al}(\text{OH})_3$ is equal to the number of equivalents of HCl . But since the number of equivalents is equal to the product of the normality, N , and the volume, V , this condition may be written as

$$N_{\text{Al}(\text{OH})_3} V_{\text{Al}(\text{OH})_3} = N_{\text{HCl}} V_{\text{HCl}}.$$

We must solve for $V_{\text{Al}(\text{OH})_3}$,

$$V = \frac{N_{\text{HCl}} V_{\text{HCl}}}{N_{\text{Al}(\text{OH})_3}}$$

The molarity of the $\text{Al}(\text{OH})_3$ solution is equal to the number of moles of $\text{Al}(\text{OH})_3$ divided by the volume in liters. To determine the number of moles corresponding to 2.6 g $\text{Al}(\text{OH})_3$, we divide 2.6 g by the formula weight of $\text{Al}(\text{OH})_3$. We then divide this by 100 ml = 100 ml \times 1 liter/1000 ml = 0.100 liter to obtain the molarity. Hence,

$$\begin{aligned} \text{concentration of } \text{Al}(\text{OH})_3 &= \frac{2.6 \text{ g}}{100 \text{ ml}} \\ &= \frac{2.6 \text{ g}/78 \text{ g/mole}}{0.100 \text{ liter}} = 0.33 \text{ M}. \end{aligned}$$

Since HCl has one ionizable proton, its normality is equal to its molarity, or $N_{\text{HCl}} = 80 \times 10^{-3} \text{ M} = 80 \times 10^{-3} \text{ N}$. $\text{Al}(\text{OH})_3$ contains three hydroxyl groups, hence, its normality is equal to three times its molarity, or $N_{\text{Al}(\text{OH})_3} = 3 \times 0.33 \text{ M} = 1.0 \text{ N}$. The required volume of medication is then

$$\begin{aligned} V_{\text{Al}(\text{OH})_3} &= \frac{N_{\text{HCl}} V_{\text{HCl}}}{N_{\text{Al}(\text{OH})_3}} = \frac{80 \times 10^{-3} \text{ N} \times 3 \text{ liters}}{1.0 \text{ N}} = 0.240 \text{ liter} \\ &= 0.240 \text{ liter} \times 1000 \text{ ml/liter} = 240 \text{ ml}. \end{aligned}$$

A chemist performs a neutralization reaction. She finds that 1 g of $C_6H_{10}O_4$, an acid, requires 0.768 g of KOH for complete neutralization. Determine the number of neutralizable protons in this molecule.

Solution: To solve this problem, first determine how many equivalents of the acid or base are involved. Once this is known, the number of neutralizable protons will also be known. One equivalent is defined as the mass of the substance, if it is an acid, needed to furnish one mole of H_3O^+ or, if it is a base, needed to furnish one mole of OH^- .

The number of grams/equiv. for KOH is 56.1, since only 1 mole of OH^- can be furnished. The number of equivalents is

$$\frac{0.768 \text{ g}}{56.1 \text{ g/equiv}} = 0.0137 \text{ equiv of KOH.}$$

The number of equivalents of base must equal that of the acid. The number of equivalents of acid = 0.0137. The number of grams/equiv is

$$\frac{1.00 \text{ g of } C_6H_{10}O_4}{0.0137 \text{ equiv.}} = 73.0 \text{ g/equiv.}$$

One mole of $C_6H_{10}O_4$ weighs 146.1 g, since a mole = weight in grams/molecular weight. Therefore, the number of moles/equiv is

$$\frac{73.0 \text{ g/equiv}}{146.1 \text{ g/mole}} = 0.5 \text{ moles/equiv.}$$

It follows that each mole of acid furnishes two H_3O^+ . As such, there are two neutralizable protons in $C_6H_{10}O_4$.

A common method for commercially peeling potatoes is to soak them for 1 - 5 minutes in a 10 - 20% solution of NaOH (molecular weight = 40.0 g/mole) at 60 - 88°C, and to spray off the peel once the potatoes are removed from solution. As an economy measure, a manufacturer titrates the NaOH solution with standardized H_2SO_4 (molecular weight = 98.0 g/mole) at the end of each day to determine whether the solution is still capable of peeling potatoes. If, at the end of one day, he finds that it takes 64.0 ml of a 0.200 M solution of H_2SO_4 to titrate a 10.0 ml sample of NaOH solution to neutrality, what concentration of NaOH did he find?

Solution: At the neutralization point, the number of equivalents of NaOH is equal to the number of equivalents of H_2SO_4 . But since the number of equivalents is equal to the product of the normality, N , and the volume, V , this condition may be stated as

$$N_{\text{NaOH}} V_{\text{NaOH}} = N_{\text{H}_2\text{SO}_4} V_{\text{H}_2\text{SO}_4}$$

To solve this problem, we must solve for N_{NaOH} ,

$$N_{\text{NaOH}} = \frac{N_{\text{H}_2\text{SO}_4} V_{\text{H}_2\text{SO}_4}}{V_{\text{NaOH}}}$$

Since H_2SO_4 has two ionizable protons, its normality is equal to twice its molarity, or $N_{\text{H}_2\text{SO}_4} = 2 \times 0.200 \text{ M} = 0.400 \text{ N}$. NaOH has only one hydroxyl group, so that its normality is equal to its molarity. Then

$$N_{\text{NaOH}} = \frac{N_{\text{H}_2\text{SO}_4} V_{\text{H}_2\text{SO}_4}}{V_{\text{NaOH}}} = \frac{0.400 \text{ N} \times 64.0 \text{ ml}}{10.0 \text{ ml}} = 2.56 \text{ N}.$$

The concentration of NaOH solution is therefore 2.56 N, or 2.56 M.

• PROBLEM 370

A potato peeling solution was found to be 2.56 M in NaOH (formula weight = 40.0 g/mole) at the end of the day. To operate, the solution must be at least 10 % NaOH by weight (100 g NaOH per 1000 g of solution). What weight percent corresponds to 2.56 M? The density of a 2.56 M solution of NaOH is about 1.10 g/ml.

Solution: To solve this problem, the concentration 2.56 M must be converted to a weight-weight basis. $2.56 \text{ M} = 2.56 \text{ moles NaOH/l liter solution} = 2.56 \text{ moles NaOH/1000 ml solution}$. To obtain the mass corresponding to 2.56 moles of NaOH, we multiply by the formula weight of NaOH, or, $2.56 \text{ moles} \times 40.0 \text{ g/moles} = 102.4 \text{ g NaOH}$.

To obtain the mass of NaOH contained in 1000 ml of solution, we multiply by the density of the solution, $1000 \text{ ml} \times 1.10 \text{ g/ml} = 1100 \text{ g}$. Hence,

$$2.56 \text{ M} = \frac{2.56 \text{ moles NaOH}}{1000 \text{ ml solution}} = \frac{102.4 \text{ g NaOH}}{1100 \text{ g solution}};$$

$$\frac{102.4 \text{ g NaOH}}{1100 \text{ g solution}} \times 100 \% = 9.3 \% \text{ by weight}.$$

Since this is less than 10 % by weight, the solution is no longer capable of peeling potatoes.

A lab technician prepared a calcium hydroxide solution by dissolving 1.48 g of Ca(OH)_2 in water. How many milliliters of 0.125 N HCl solution would be required to neutralize this calcium hydroxide solution?

Solution: For a neutralization reaction to occur, the number of equivalents of acid must equal the number of equivalents of base. Therefore, one can solve this problem by computing how many equivalents are present in 1.48 g of Ca(OH)_2 ; the equivalents of acid (HCl) must equal this number.

An equivalent is defined as the weight of a substance in grams that releases one mole of protons (H^+) or hydroxyl ions. When one mole of Ca(OH)_2 dissolves in water, it produces 2 moles of hydroxyl ions (OH^-) according to the equation $\text{Ca(OH)}_2 \rightarrow \text{Ca}^{2+} + 2\text{OH}^-$. The weight of 1 mole of Ca(OH)_2 is equal to its molecular weight. Therefore, Ca(OH)_2 has a weight of 74 grams.

Thus, 74 grams of Ca(OH)_2 produces 2 moles of OH^- ions. Recalling the definition of equivalency, $74/2 = 37$ grams is the equivalent weight of Ca(OH)_2 . The number of equivalents of Ca(OH)_2 is:

$$\text{Ca(OH)}_2 = \frac{\text{weight in grams}}{\text{equivalent weight}} = \frac{1.48 \text{ g}}{37 \text{ g}} = 0.04 \text{ equiv.}$$

The number of equivalents of HCl must also be .04 for neutralization to occur. The normality of HCl is given as 0.125. Normality is defined as the number of equivalents divided by liters of solution, i.e., $N = \text{equivalents/volume}$.

This means that $0.125 \text{ N} = 0.04 \text{ equivalents/volume}$ for HCl. The volume in liters required for neutralization can be obtained by solving this equation.


$$\text{liters} = \frac{0.04}{0.125} = 0.32 \text{ l}$$

To convert to milliliters multiply by the conversion factor 1000 ml/l. One has


$$0.32 \text{ l} \times 1000 \text{ ml/l} = 320 \text{ ml.}$$

50 ml of rhubarb juice is titrated against 0.25 N NaOH. 20 ml of NaOH solution is required for neutralization. Assuming the acidity of the juice is due to oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$) determine (a) the weight of oxalic acid per liter of juice, (b) normality of the juice.

Solution: (a) For the neutralization to occur, there must be the same number of OH^- ions as there are H^+ ions present.



The normality of a base is defined as the number of equivalents of base in one liter of solution. An equivalent is the weight of an acid or base that produces one mole of H^+ or OH^- ions, respectively. When NaOH ionizes, there is 1 OH^- ion formed by each NaOH that ionizes.




Therefore, in a 0.25 N NaOH solution there are 0.25 equivalents of OH^- , which, in this case is also the number of moles. 20 ml is 0.02 liters. The number of moles of NaOH present will equal the number of liters of solution times the molarity of the solution. As previously indicated, normality equals molarity. Thus,

no. of moles = no. of liters \times normality

$$= 0.02 \text{ liters} \times \frac{.25 \text{ moles}}{\text{liter}} = 0.005 \text{ moles.}$$

Thus, there must be 0.005 moles of H^+ ions from the oxalic acid to neutralize the NaOH. When oxalic acid ionizes, there are 2 H^+ ions formed for each molecule ionized.



Therefore, each mole of oxalic acid ionizes 2 moles of NaOH. Thus, for neutralization to occur, only one half as much oxalic acid is needed as NaOH. There are 0.005 moles of NaOH, present, therefore, 0.0025 moles of oxalic acid is needed to neutralize it. One now knows that there are 0.0025 moles of oxalic acid in the 50 ml of rhubarb juice. One can find the number of moles in 1 liter by multiplying 0.0025 moles/50 ml by the conversion factor 1000 ml/1 liter.

$$\begin{aligned} \text{no. of moles/liter of oxalic acid} &= \frac{0.0025 \text{ moles}}{50 \text{ ml}} \times \frac{1000 \text{ ml}}{1 \text{ liter}} \\ &= 0.05 \text{ moles/liter.} \end{aligned}$$

Since there are 0.05 moles of $\text{H}_2\text{C}_2\text{O}_4$ in 1 liter of juice, one can find the weight of $\text{H}_2\text{C}_2\text{O}_4$ in this quantity of juice by multiplying 0.05 moles by the MW of $\text{H}_2\text{C}_2\text{O}_4$. (MW = 90).

$$\begin{aligned} \text{weight of } \text{H}_2\text{C}_2\text{O}_4 \text{ in 1 liter} &= 0.05 \text{ moles} \times 90 \text{ g/moles} \\ &= 4.5 \text{ g} \end{aligned}$$

(b) For titrations, the following relation is found:

$$N_{\text{acid}} V_{\text{acid}} = N_{\text{base}} V_{\text{base}}$$

where N_{acid} is the normality of the acid, V_{acid} is the volume of the acid, N_{base} is the normality of the base, and V_{base} is the volume of the base.

Here, one is given N_{base} , V_{base} , and V_{acid} . One is asked to find N_{acid} .

$$N_b = .25 \text{ N}$$

$$N_a V_a = N_b V_b$$

$$V_b = 20 \text{ ml}$$

$$N_a \times 50 \text{ ml} = .25 \text{ N} \times 20 \text{ ml}$$



$$N_a = ?$$

$$V_a = 50 \text{ ml}$$

$$N_a = \frac{0.25 \text{ N} \times 20 \text{ ml}}{50 \text{ ml}} = 0.10 \text{ N.}$$

• PROBLEM 373

The $^+\text{NH}_3\text{CH}_2\text{COOH}$ ion (glycine $^+$) is a cation of the amino acid glycine ($\text{NH}_2\text{CH}_2\text{COOH}$). Glycine $^+$ undergoes the following two successive dissociations:



What is the isoelectric point of this molecule (the pH at which the number of positive charges on the molecule equals the number of negative charges).

Solution: We will approach this problem by considering the equilibrium constants k_1 and k_2 . For the first reaction, k_1 is equal to

$$k_1 = \frac{[\text{H}^+][^+\text{NH}_3\text{CH}_2\text{COO}^-]}{[^+\text{NH}_3\text{CH}_2\text{COOH}]}$$

and for the second reaction, k_2 is equal to

$$k_2 = \frac{[\text{H}^+][\text{NH}_2\text{CH}_2\text{COO}^-]}{[^+\text{NH}_3\text{CH}_2\text{COO}^-]}$$

If we take the product of these two constants, the concentration of isoelectric species $^+\text{NH}_3\text{CH}_2\text{COO}^-$ cancels out, giving

$$\begin{aligned}
 k_1 k_2 &= \frac{[H^+][^+NH_3CH_2COO^-]}{[^+NH_3CH_2COOH]} \times \frac{[H^+][NH_2CH_2COO^-]}{[^+NH_3CH_2COO^-]} \\
 &= \frac{[H^+]^2 [NH_2CH_2COO^-]}{[^+NH_3CH_2COOH]}
 \end{aligned}$$

At the isoelectric point, the concentrations of all charged species are equal, that is

$$[NH_2CH_2COO^-] = [^+NH_3CH_2COOH].$$

Hence, $\frac{[NH_2CH_2COO^-]}{[^+NH_3CH_2COOH]} = 1$ and

$$k_1 k_2 = \frac{[H^+]^2 [NH_2CH_2COO^-]}{[^+NH_3CH_2COOH]} = [H^+]^2.$$

Taking the logarithm of both sides,

$$[H^+]^2 = k_1 k_2$$

$$2 \log [H^+] = \log k_1 k_2 = \log k_1 + \log k_2.$$

Multiplying this equation by negative one and using the definitions $pH = -\log [H^+]$, $pk_1 = -\log k_1$, and $pk_2 = -\log k_2$, we obtain

$$-2 \log [H^+] = -\log k_1 + (-\log k_2)$$

$$2 pH = pk_1 + pk_2, \quad \text{or,} \quad pH = \frac{1}{2}(pk_1 + pk_2).$$

Thus, the isoelectric point is

$$pH = \frac{1}{2}(pk_1 + pk_2) = \frac{1}{2}(2.35 + 9.78) = 6.07.$$


BUFFERS

• PROBLEM 374


Explain the buffering action of a liter of 0.10 M acetic acid containing 0.1 mole of sodium acetate. In the explanation, use ionic equations.

Solution: By buffering action, one means the ability of a substance to maintain relatively constant conditions of pH in the face of changes that might otherwise affect the acidity or basicity in solution. For example, a weak acid, such as acetic acid, will dissociate according to the

following equation:



The sodium acetate particle will be completely ionized in solution:



These are the two processes that occur in this buffer. Suppose one increases the acetate concentration. By doing this the equilibrium is shifted to the left, i.e., to acetic acid.

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{OAc}^-]}{[\text{HOAc}]} \quad \text{or} \quad [\text{H}_3\text{O}^+] = \frac{[\text{HOAc}]}{[\text{OAc}^-]} K_a.$$

The key to the buffering action is this $[\text{HOAc}]$ to $[\text{OAc}^-]$ ratio. If $[\text{HOAc}]$ changes, $[\text{OAc}^-]$ will change accordingly so that the same value of the ratio is obtained. Thus, the ratio is a constant. This means, therefore, that $[\text{H}_3\text{O}^+]$ is maintained as a constant. $\text{pH} = -\log [\text{H}^+]$, it is also constant.

• PROBLEM 375

Design a buffer system that will function in the low, near neutral, and high pH levels, using combinations of the three sodium phosphates (Na_3PO_4 , Na_2HPO_4 , NaH_2PO_4) and phosphoric acid (H_3PO_4). The equilibrium constants for this polyprotic acid are $k_1 = 7.5 \times 10^{-3}$, $k_2 = 6.2 \times 10^{-8}$, and $k_3 = 4.8 \times 10^{-13}$. With this information, calculate the pH at both extremes of the buffer, assume 10 : 1 and 1 : 10 ratios, and at the mid-range of the buffer, assume a 1 : 1 ratio. Assume, also, that the acid-salt ratio in making up the buffer is equal to HA/A^- in solution.

Solution: Solutions that contain appreciable amounts of a weak acid, such as phosphoric acid, and its salt or salts, such as sodium phosphates, are called buffers. Their utility is in maintaining a relatively constant pH. This problem requires one to find various pH values for this buffer system. To do this, note that the pH of a buffer is given by

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]} = \text{p}K_a + \log \frac{[\text{salt}]}{[\text{acid}]}.$$

One knows that to find the pH, one must determine $[\text{H}_3\text{O}^+]$, since $\text{pH} = -\log [\text{H}_3\text{O}^+]$. To do this consider the ionization of the polyprotic acid, phosphoric acid, which can dissociate to produce 3 protons. k_1 , the first ionization constant, corresponds to $\text{H}_3\text{PO}_4 + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{H}_2\text{PO}_4^-$ so that

$$k_1 = \frac{[\text{H}_3\text{O}^+][\text{H}_2\text{PO}_4^-]}{[\text{H}_3\text{PO}_4]}.$$

k_2 , the second ionization constant, corresponds to $\text{H}_2\text{PO}_4^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HPO}_4^{2-}$ so that

$$k_2 = \frac{[\text{H}_3\text{O}^+][\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]}.$$

k_3 , the third ionization constant, corresponds to $\text{HPO}_4^{2-} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{PO}_4^{3-}$ and

$$k_3 = \frac{[\text{H}_3\text{O}^+][\text{PO}_4^{3-}]}{[\text{HPO}_4^{2-}]}.$$

In general, the k_a of any acid is given by the expression

$$k_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]} \quad \text{or} \quad [\text{H}_3\text{O}^+] = k_a \frac{[\text{HA}]}{[\text{A}^-]}.$$

It is from this equation that the $[\text{H}_3\text{O}^+]$ must be calculated, and thus yield the pH value wanted. The three systems:

Buffer system with k_1

Consider the ratios of 10 : 1, 1 : 1, and 1 : 10 for the acid : salt ratios, given that the ratio equals $[\text{HA}]/[\text{A}^-]$ for the low, neutral and high pH values, respectively. This means that $[\text{H}_3\text{O}^+] = 10 k_a$, k_a and $0.1 k_a$ for low, neutral, and high, respectively. Here,

$k_a = k_1$ which is $= 7.5 \times 10^{-3}$. $\text{pk}_1 = 2.12$. Since $[\text{H}_3\text{O}^+] = 10 k_a$, k_a , and $0.1 k_a$; $\text{pH} = -\log [\text{H}_3\text{O}^+] = (\text{pk}_a - 1)$, pk_a and $(\text{pk}_a + 1)$ for low, middle, and high pH values, respectively. Thus, with $\text{pk}_1 = 2.12$, $\text{pH} = 1.12$ (low), 2.12 (middle) and 3.12 (high).

Buffer system with k_2

One still has $\text{pH} = (\text{pk}_a - 1)$, pk_a , and $\text{pk}_a + 1$. But now $k_a = k_2 = 6.2 \times 10^{-8}$ $\text{pk}_2 = 7.21$. Thus, $\text{pH} = 6.21$ (low), 7.21 (middle), and 8.21 (high).

Buffer system with k_3

One still has $\text{pH} = (\text{pk}_a - 1)$, pk_a , and $\text{pk}_a + 1$. Now, $k_a = k_3 = 4.8 \times 10^{-13}$ $\text{pk}_3 = 12.32$. Thus, $\text{pH} = 11.32$ (low), 12.32 (middle), and 13.32 (high).

The following bases, and their conjugate acids (as the chlorides), are available in the lab: ammonia, NH_3 ; pyridine, $\text{C}_5\text{H}_5\text{N}$; ethylamine, $\text{CH}_3\text{CH}_2\text{NH}_2$. A buffer solution of pH 9 is to be prepared, and the total concentration of buffering reagents is to be 0.5 mole/liter. (a) Choose the best acid-base pair. (b) Give the recipe for preparing one liter of the solution. (c) Calculate the pH of the solution after 0.02 mole of NaOH has been added per liter.

Solution: This problem deals with the preparation of a buffer solution and then to show the effect the addition of a base has on the pH.

If this is a truly buffered system, then, the pH should be only slightly altered when a small quantity of base is added.

Since we are using bases it will be more convenient to work initially with pOH. pH is converted to pOH by the equation:

$$14 = \text{pH} + \text{pOH} \quad \text{or} \quad \text{pOH} = 14 - \text{pH}.$$

Substituting the given value of pH, $\text{pOH} = 14 - 9 = 5$.

In part (a) one is asked to choose the strongest base. At a given pOH, the strongest base has pOH about equal to pK_b (the pH at which dissociation occurs). The maximum buffering capacity and minimum sensitivity to pH change occurs when the concentrations of the acid and base are initially equal. From a table of pK_b values, it is found that $\text{pK}_{\text{NH}_3} = 4.75$, $\text{pK}_{\text{C}_5\text{H}_5\text{N}} = 8.81$, and $\text{pK}_{\text{CH}_3\text{CH}_2\text{NH}_2} = 3.3$. Therefore, NH_3 has a pK_b closest to 5, which means it is the best base.

To prepare a one liter solution, as requested in part (b), find the initial concentrations of the components to be added and their ratio. NH_3 is the base and NH_4Cl dissociates to the acid NH_4^+ . The reaction equation is



$$K_b = 1.76 \times 10^{-5}.$$

To find the ratio, set up the equilibrium constant equation:

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.76 \times 10^{-5}.$$

Rewriting to obtain the ratio, one has:

$$\frac{[\text{NH}_4^+]}{[\text{NH}_3]} = \frac{K_b}{[\text{OH}^-]} = \frac{1.76 \times 10^{-5}}{1.0 \times 10^{-5}} = 1.76.$$

($[\text{OH}^-] = 1 \times 10^{-5}$, since $\text{pOH} = 5$ and $\text{pOH} = -\log [\text{OH}^-]$.)

Therefore, NH_4Cl and NH_3 should be added in the molar ratio of 1.76 : 1. Since it is given that the sum of the concentrations of the buffering reagents equals 0.5 mole/liter, the following equation can be written which says, in effect, that the sum of the parts is equal to the whole

$$[\text{NH}_3] + [\text{NH}_4^+] = 0.5.$$

Let $[\text{NH}_3] = x$ and since the ratio is 1.76 : 1, let $[\text{NH}_4^+] = 1.76 x$.

This results in

$$x + 1.76 x = 0.5$$


$$x = [\text{NH}_3] = 0.18 \text{ mole/liter}$$

$$1.76 x = [\text{NH}_4^+] = 0.32 \text{ mole/liter.}$$

Therefore, to prepare a 1 liter solution, mix 0.18 moles NH_3 and 0.32 moles NH_4Cl and then add H_2O until a total volume of 1 liter is obtained.

Part (c) tells one to add 0.02 M NaOH . Since it is a strong base, complete dissociation occurs giving 0.02 mole Na^+ and 0.02 mole OH^- . The base OH^- then reacts fully with the acid NH_4^+ and converts 0.02 mole NH_4^+ to NH_3 .

This is more clearly seen below.



Before: 0.32 mole 0.18 mole

After: 0.32-0.02=0.30 mole 0.18+0.02=0.20 mole

Substituting these new values into the equilibrium constant equation, one arrives at $[\text{OH}^-]$ and then pH. One then has

$$K_b = \frac{[\text{NH}_3][\text{OH}^-]}{[\text{NH}_4^+]}. \text{ Substituting,}$$

$$[\text{OH}^-] = \frac{K_b [\text{NH}_4^+]}{[\text{NH}_3]} = \frac{1.76 \times 10^{-5} (0.20)}{(0.30)}$$

$$= 1.17 \times 10^{-5} \text{ mole/liter.}$$


Thus, $\text{pOH} = -\log [\text{OH}^-] = -\log [1.17 \times 10^{-5}] = 4.93$.

Thus, $\text{pH} = 14 - \text{pOH} = 14 - 4.93 = 9.07$.

INDICATORS

• PROBLEM 377

Using methyl orange as an indicator, a solution of unknown pH was matched to a Bjerrum wedge at a point where the acid wedge was 40 percent of the total thickness of the combined wedges. This point is indicated by the arrow in the diagram. Determine the pH of the solution.



Bjerrum wedge

Solution: From the Bjerrum wedge, the ratio of the concentration of methyl orange indicator in basic solution ($[\text{Ind}^-]$) to the concentration in acid solution ($[\text{HInd}]$) will be determined. This ratio will then be used to find the pH of the unknown solution.

A Bjerrum wedge is a glass box divided into two wedge-shaped compartments by a glass plate placed diagonally in the box (see diagram, above). In one compartment is placed an acidic solution of the chosen indicator (HInd) and into the other a basic solution of the indicator (Ind^-). The concentration of indicator is the same in both wedges. A view from the top will produce a continuum of color varying from that of pure HInd on one side to that of pure Ind^- on the other. A solution of indicator of the same concentration as the Bjerrum wedge is prepared using a solution of unknown pH. The resulting solution is then

placed in a glass box having a thickness equal to that of the Bjerrum wedge. The boxes are placed side-by-side and the color of the unknown solution is matched to a color on the Bjerrum wedge. At the point where the two colors match, let the acidic wedge be x percent of the total thickness of the box. Then the basic wedge is $100 - x$ percent of the total thickness. The ratio $[\text{Ind}^-]/[\text{HInd}]$ is then

$$\frac{[\text{Ind}^-]}{[\text{HInd}]} = \frac{100 - x}{x} .$$

In this problem, the acidic wedge is 40 percent the total thickness, hence

$$\frac{[\text{Ind}^-]}{[\text{HInd}]} = \frac{100 - 40}{40} = \frac{60}{40} .$$

To make use of this ratio, we need an expression relating the pH to $[\text{Ind}^-]/[\text{HInd}]$. Consider the dissociation of acidic indicator:



The equilibrium constant, K_a , for this reaction is

$$K_a = [\text{H}_3\text{O}^+] \times \frac{[\text{Ind}^-]}{[\text{HInd}]} .$$

Taking the logarithm of both sides and multiplying by negative one gives

$$- \log K_a = - \log [\text{H}_3\text{O}^+] - \log \frac{[\text{Ind}^-]}{[\text{HInd}]} .$$

$$- \log K_a = \text{p}K_a \quad \text{and} \quad - \log [\text{H}_3\text{O}^+] = \text{pH} .$$

$$\text{Hence,} \quad \text{p}K_a = \text{pH} - \log \frac{[\text{Ind}^-]}{[\text{HInd}]} ,$$

$$\text{or,} \quad \text{pH} = \text{p}K_a + \log \frac{[\text{Ind}^-]}{[\text{HInd}]} .$$

For methyl orange, $\text{p}K_a = 3.7$. Using the value of $[\text{Ind}^-]/[\text{HInd}]$ obtained above, we get


$$\text{pH} = 3.7 + \log \left(\frac{60}{40} \right) = 3.9$$

as the pH of our unknown solution.

For the indicator phenolphthalein (In^-), HIn is colorless and In^- is red; K_{diss} is 10^{-9} . If one has a colorless solution of 50 ml phenolphthalein, how much 0.10 M NaOH solution would one have to add to make it red: $K_w = 10^{-14}$.

Solution: To answer this question, write out the dissociation reaction of phenolphthalein in solution. After this, write out the equilibrium constant expression. This allows one to calculate the $[\text{H}^+]$. This informs one of the concentration of OH^- required and allows the calculation of the amount of NaOH to be added.

With this in mind, proceed as follows: An indicator, which changes color, undergoes a dissociation reaction like an acid. For phenolphthalein the reaction is



It is given that In^- makes the solution red and HIn makes it colorless. The equilibrium constant expression, which measures the ratio of concentrations of products to reactants, each raised to the power of its coefficient in the equation, is

$$K_{\text{diss}} = 10^{-9} = \frac{[\text{H}^+][\text{In}^-]}{[\text{HIn}]}$$

The object is to make a red solution, from an originally colorless one. In general, a specific color of an indicator will show itself when the concentration of the corresponding species is 10 times as great as the concentration of the other species. Since one wants the solution to be red, and In^- = red color, $[\text{In}^-]/[\text{HIn}] = 10/1$. Substituting this into the equilibrium constant expression,

$$10^{-9} = [\text{H}^+] \frac{10}{1}. \text{ Solving for } [\text{H}^+], [\text{H}^+] = 10^{-10}.$$

From the K_w constant of water, it is known that $[\text{H}^+][\text{OH}^-] = 10^{-14}$, where K_w is the autodissociation constant of H_2O . If $[\text{H}^+] = 10^{-10}$, one can calculate $[\text{OH}^-]$ and obtain $[\text{OH}^-] = 10^{-4}$ M. Thus, to have a red solution, one wants $[\text{OH}^-]$ to be 10^{-4} M. The NaOH solution being added is 0.1 M. (M = molarity = moles/liter.) The volume of the solution is 50 ml. or 0.05 l. (1000 ml = 1 l.) Thus, one needs 10^{-4} M or $0.050 \text{ l} \times 10^{-4} \text{ M} = 5.0 \times 10^{-6}$ moles. Since volume in liters = number of moles/molarity, and molarity = 0.1, liters of NaOH to be added =


$$= \frac{5.0 \times 10^{-6}}{0.1} = 5.0 \times 10^{-5} \text{ liters.}$$

COMPLEX IONS

• PROBLEM 379

For the complex ion equilibrium for the dissociation of $\text{Cu}(\text{NH}_3)_4^{2+}$, calculate (1) the ratio of Cu^{2+} ion to $\text{Cu}(\text{NH}_3)_4^{2+}$ in a solution 1 M in NH_3 ; (2) the equilibrium concentration of NH_3 needed to attain 50 % conversion of Cu^{2+} to $\text{Cu}(\text{NH}_3)_4^{2+}$. K_d for $\text{Cu}(\text{NH}_3)_4^{2+} = 2 \times 10^{-13}$.

Solution: (1) The equilibrium constant for the reaction



is defined as

$$K = \frac{[\text{B}]^n [\text{C}]^m}{[\text{A}]}$$

where K is the equilibrium constant, $[\text{B}]$ is the concentration of B, n is the number of moles of B formed, $[\text{C}]$ is the concentration of C, m is the number of moles of C produced and $[\text{A}]$ is the concentration of A. The reaction in this problem is



This indicates that the equilibrium constant can be stated

$$K_d = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 2 \times 10^{-13}$$

Here, one is trying to find the ratio of Cu^{2+} ion to $\text{Cu}(\text{NH}_3)_4^{2+}$ in the solution or stated in another way

$\frac{[\text{Cu}^{2+}]}{[\text{Cu}(\text{NH}_3)_4^{2+}]}$. To obtain this ratio, one can use the equation for K_d . In the problem, one is told that the concentration of NH_3 is 1 M. Therefore,

$$\frac{[\text{Cu}^{2+}]}{[\text{Cu}(\text{NH}_3)_4^{2+}]} \text{ can be found.}$$

$$K_d = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 2 \times 10^{-13}$$

$$\frac{[\text{Cu}^{2+}][1.0]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 2 \times 10^{-13}$$

$$\frac{[\text{Cu}^{2+}]}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = \frac{2 \times 10^{-13}}{(1.0)^4} = 2 \times 10^{-13}$$

(b) To find the equilibrium concentration of NH_3 needed to convert 50 % of the Cu^{2+} ion present to $\text{Cu}(\text{NH}_3)_4^{2+}$ ion, the equation for the equilibrium constant will again be used.

$$K_d = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 2 \times 10^{-13}$$

When 50 % of Cu^{2+} is converted to $\text{Cu}(\text{NH}_3)_4^{2+}$, the concentration of Cu^{2+} will equal that of $\text{Cu}(\text{NH}_3)_4^{2+}$. This is true because there is one Cu^{2+} ion used in the formation of each $\text{Cu}(\text{NH}_3)_4^{2+}$ ion. If there were 50 Cu^{2+} ions in the solution and 25 of them were converted to $\text{Cu}(\text{NH}_3)_4^{2+}$, there would be 25 Cu^{2+} ions left and 25 $\text{Cu}(\text{NH}_3)_4^{2+}$ ions formed. This means that for a 50% conversion

$$\frac{[\text{Cu}^{2+}]}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 1$$

One can now solve for the $[\text{NH}_3]$ by using the equation for K_d .

$$K_d = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 2 \times 10^{-13}$$

$$(1)[\text{NH}_3]^4 = 2 \times 10^{-13}$$

$$[\text{NH}_3] = \sqrt[4]{2 \times 10^{-13}}$$

$$[\text{NH}_3] = 6.7 \times 10^{-3} \text{ M}$$

A concentration of $6.7 \times 10^{-3} \text{ M}$ of NH_3 will cause a 50 % conversion of Cu^{2+} to $\text{Cu}(\text{NH}_3)_4^{2+}$.

• PROBLEM 380

You have the four complex ions:

- | | |
|--|---------------------------|
| (1) $\text{Cu}(\text{NH}_3)_4^{2+} \rightleftharpoons \text{Cu}^{2+} + 4\text{NH}_3$ | $k = 1.0 \times 10^{-12}$ |
| (2) $\text{Co}(\text{NH}_3)_6^{2+} \rightleftharpoons \text{Co}^{2+} + 6\text{NH}_3$ | $k = 4.0 \times 10^{-5}$ |
| (3) $\text{Co}(\text{NH}_3)_6^{3+} \rightleftharpoons \text{Co}^{3+} + 6\text{NH}_3$ | $k = 6.3 \times 10^{-36}$ |
| (4) $\text{Ag}(\text{NH}_3)_2^+ \rightleftharpoons \text{Ag}^+ + 2\text{NH}_3$ | $k = 6.0 \times 10^{-8}$ |

Arrange them in order of increasing NH_3 concentration that would be in equilibrium with complex ion, if you start with 1 M of each.

Solution: To solve this problem determine the $[\text{NH}_3]$ for each complex ion equilibria. This necessitates writing

the equilibrium constant expression, which measures the ratio of concentrations of products to reactants, each raised to the power of their coefficients in the chemical reaction. Proceed as follows:

$$K = 1.0 \times 10^{-12} = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]}$$

Start with 1 M of $\text{Cu}(\text{NH}_3)_4^{2+}$. Let $x = [\text{NH}_3]$ at equilibrium. From the chemical reaction one can see that 1 mole of Cu^{2+} is formed as 4 moles of NH_3 are formed. Thus, $[\text{Cu}^{2+}] = \frac{1}{4}[\text{NH}_3] = 0.25x$. If x moles/liter of NH_3 form, and the only source is $\text{Cu}(\text{NH}_3)_4^{2+}$, then, at equilibrium, $[\text{Cu}(\text{NH}_3)_4^{2+}] = 1 - 0.25x$. It is initial concentration minus $\frac{1}{4}$ of $[\text{NH}_3]$, since they have a mole ratio of 1 : 4, as is seen in the chemical reaction. Substituting into the equilibrium equation:

$$\frac{(0.25x)(x)^4}{(1 - 0.25x)} = 1.0 \times 10^{-12}.$$

Solving for x , $x = [\text{NH}_3] = 5.3 \times 10^{-3}$.

To obtain this answer the denominator $(1 - 0.025x)$ is approximated as 1, since the value for x is relatively small compared to 1.

To find $[\text{NH}_3]$ at equilibrium in (2), (3), and (4), employ the same type of logic and reasoning. The only differences are the values of K and the mole ratios between the substances.

$$\text{For } \text{Co}^{2+} : 1/6 x^7 = 4.0 \times 10^{-5}$$

$$[\text{NH}_3] = x = 0.3$$

$$\text{For } \text{Co}^{3+} : 1/6 x^7 = 6.3 \times 10^{-36}$$

$$[\text{NH}_3] = x = 9.4 \times 10^{-6}$$

$$\text{For } \text{Ag}^+ : \frac{1}{2} x^3 = 6.0 \times 10^{-8}$$

$$[\text{NH}_3] = x = 5.0 \times 10^{-3}$$

In increasing order, rank them as


$$\begin{array}{c} \text{Co}^{3+}, \text{Ag}^+, \text{Cu}^{2+}, \text{Co}^{2+} \\ \hline \text{increasing } [\text{NH}_3] \end{array}$$

• PROBLEM 381


0.1 moles of $\text{Cu}(\text{NO}_3)_2$ are added to 1.0 moles of NH_3 and then diluted to 1000ml. Find the concentration of Cu^{2+} formed.

$$K_{\text{eq}} = 8.5 \times 10^{-13}.$$

Solution: The first thing to realize is that the complex $\text{Cu}(\text{NO}_3)_2$ dissociates to $\text{Cu}^{2+} + 2\text{NO}_3^-$ in H_2O



when NH_3 is added to the solution the following reaction occurs.



$\text{Cu}(\text{NH}_3)_4^{2+}$ is called a complex ion. $\text{Cu}(\text{NH}_3)_4^{2+}$ will be in equilibrium with $\text{Cu}^{2+} + 4\text{NH}_3$. You are given the K_{eq} for this reaction which is defined as

$$K = \frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]}$$

In general, if you have the equilibrium: $\text{H}(\text{A})_X \rightleftharpoons \text{H}^+ + \text{XA}^-$, then

$$K = \frac{[\text{H}^+][\text{A}^-]^X}{[\text{H}(\text{A})_X]}$$

That is, the coefficients of the balanced equation serve as the exponents in the K expression. The problem stated $K = 8.5 \times 10^{-13}$. Therefore,

$$\frac{[\text{Cu}^{2+}][\text{NH}_3]^4}{[\text{Cu}(\text{NH}_3)_4^{2+}]} = 8.5 \times 10^{-13}$$

You are solving for $[\text{Cu}^{2+}]$. Let $x = [\text{Cu}^{2+}]$, then $[\text{Cu}(\text{NH}_3)_4^{2+}] = .1 - x$, since it must be the original amount of $\text{Cu}(\text{NO}_3)_2$ present minus the Cu^{2+} formed. There is originally 1.0 moles NH_3 , at equilibrium 4 moles of NH_3 will have reacted with each mole of Cu^{2+} to form $\text{Cu}(\text{NH}_3)_4^{2+}$, thus, $[\text{NH}_3]$ may be represented as $1.0 - 4x$ $[\text{Cu}(\text{NH}_3)_4^{2+}] = 1.0 - 4(.10 - x)$. Since most of the Cu^{2+} present will be in the form of $\text{Cu}(\text{NH}_3)_4^{2+}$ and that there was only .1M of it in the first place, one can make the assumption that $x \ll .1$, therefore, $.10 - x = .1$ and $1.0 - 4(.10 - x) = 1.0 - 4(.1) = 0.6 = [\text{NH}_3]$. Using this same assumption, $[\text{Cu}(\text{NH}_3)_4^{2+}] = .1 - x = .1$.

You now substitute these values into the previous expression. That is,

$$\frac{x(0.6)^4}{.1} = 8.5 \times 10^{-13}, \quad x = 6.5 \times 10^{-13}$$

Therefore, $[\text{Cu}^{2+}] = 6.5 \times 10^{-13} \text{M}$.

An equilibrium solution of the complex ion $\text{Ag}(\text{NH}_3)_2^+$ contains 0.30 M of NH_4^+ and 0.15 M of the actual complex ion. To obtain a concentration of Ag^+ equal to 1.0×10^{-8} M, what must the pH of this solution be? K_{diss} of $\text{Ag}(\text{NH}_3)_2^+ = 6.0 \times 10^{-8}$, K_b of $\text{NH}_3 = 1.8 \times 10^{-5}$ and $K_w = 1 \times 10^{-14}$.


Solution: To find the pH required, calculate the $[\text{H}^+]$ ($\text{pH} = -\log [\text{H}^+]$). To find $[\text{H}^+]$, the $[\text{OH}^-]$ is needed. One is given K_w , which is the constant for the autodis-

sociation of H_2O . K_w is defined as $[\text{H}^+][\text{OH}^-]$ for $\text{H}_2\text{O} \rightleftharpoons \text{H}^+ +$

OH^- . If $[\text{OH}^-]$ was known, one could solve for $[\text{H}^+]$. To find $[\text{OH}^-]$, consider the K_b value given. When bases, such as NH_3 ,

are placed in water, these bases dissociate into positive and negative ions. The K_b , the equilibrium constant, measures the ratio of the concentration of products to re-


actants, each raised to the power of its coefficient in the chemical reaction. When NH_3 is placed in water, following dissociation exists:



$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]}$$

Note: H_2O is omitted; it is assumed to be a constant. Therefore, if $[\text{NH}_4^+]/[\text{NH}_3]$ was known, one could find $[\text{OH}^-]$. To find $[\text{NH}_4^+]/[\text{NH}_3]$, calculate $[\text{NH}_3]$ since $[\text{NH}_4^+]$ is given. To calculate $[\text{NH}_3]$, use the equilibrium constant expression for the complex ion dissociation. This constant, K_{diss} , is defined in the same way as K_b (i.e., ratio of products to reactants). The problem provides all the information needed to find $[\text{NH}_3]$. Proceed as follows:

The complex ion dissociation is similar in nature to other equilibrium dissociations.



$$K_{\text{diss}} = \frac{[\text{Ag}^+][\text{NH}_3]^2}{[\text{Ag}(\text{NH}_3)_2^+]}$$

As given, $[\text{Ag}(\text{NH}_3)_2^+] = 0.15 \text{ M}$, one wants $[\text{Ag}^+] = 1.0 \times 10^{-8} \text{ M}$ at equilibrium. Thus, substituting these values, one finds that

$$K_{\text{diss}} = 6.0 \times 10^{-8} = \frac{(1.0 \times 10^{-8})[\text{NH}_3]^2}{0.15}$$

Solving for $[\text{NH}_3]$, one obtains

$$[\text{NH}_3] = \sqrt{\frac{(6.0 \times 10^{-8})(0.15)}{1.0 \times 10^{-8}}} = 9.5 \times 10^{-2}$$

One is told $K_b = 1.8 \times 10^{-5}$. By definition,

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]}$$

$[\text{NH}_3]$ has been found and $[\text{NH}_4^+]$ is given. Thus, by substitution one can solve for $[\text{OH}^-]$.

$$1.8 \times 10^{-5} = \frac{(0.30)[\text{OH}^-]}{9.5 \times 10^{-2}} \quad \text{or}$$

$$[\text{OH}^-] = \frac{(1.8 \times 10^{-5})(9.5 \times 10^{-2})}{0.30} = 5.7 \times 10^{-6}$$

Recalling that $K_w = 1.0 \times 10^{-14} = [\text{OH}^-][\text{H}^+]$, one has

$$1.0 \times 10^{-14} = [\text{H}^+][5.7 \times 10^{-6}].$$

$$\text{Solving, } [\text{H}^+] = \frac{1.0 \times 10^{-14}}{5.7 \times 10^{-6}} = 1.75 \times 10^{-9}$$

$$\text{Since } \text{pH} = -\log [\text{H}^+],$$

$$\text{pH} = -\log [1.75 \times 10^{-9}] = 8.76.$$

Thus, bring the pH up to 8.76 to obtain an $[\text{Ag}^+] = 1.0 \times 10^{-6} \text{ M}$.

ELECTROLYTES


• PROBLEM 383

Describe, by both formal and ionic equations, what takes place when 0.10 mole of sodium sulfite is added to 1 liter of 1 M hydrochloric acid.


Solution: The formal equation for the reaction is



The ionic equation gives more detailed information than this. The soluble sodium sulfite salt will react with excess hydrochloric acid according to the ionic reaction



Sulfurous acid, H_2SO_3 , exists in equilibrium with both sulfur dioxide and hydronium ions:



SO_2 gas will escape from solution, thus pulling the reaction towards completion, to give the products $2(\text{Na}^+, \text{Cl}^-)$, SO_2 , and $2\text{H}_2\text{O}$.

• PROBLEM 384


A saturated solution of the strong electrolyte $\text{Ca}(\text{OH})_2$ is prepared by adding sufficient water to 5.0×10^{-4} mole of $\text{Ca}(\text{OH})_2$ to form 100 ml of solution. What is the pH of this solution?

Solution: We need to find $[H_3O^+]$ in order to find the pH. $[H_3O^+]$ is determined by substituting into the expression for the water constant, $K_w = [H_3O^+][OH^-]$

($K_w = 10^{-14}$ mole²/liter²), or

$$[H_3O^+] = \frac{K_w}{[OH^-]}$$

The hydroxyl concentration, $[OH^-]$, is contributed by the dissociation of $Ca(OH)_2$, which is given by the equation



$Ca(OH)_2$ is a strong electrolyte, hence we assume that it dissociates completely. Since two hydroxyl ions are formed for every $Ca(OH)_2$ that dissociates, the concentration of OH^- is equal to twice that of $Ca(OH)_2$, or $[OH^-] = 2 \times (0.0005 \text{ mole}/100 \text{ ml}) = 2 \times (0.0005 \text{ mole}/0.10 \text{ liter}) = 2 \times 0.005 \text{ mole/liter} = 0.010 \text{ mole/liter} = 10^{-2} \text{ mole/liter}$. The hydronium ion concentration is then

$$[H_3O^+] = \frac{K_w}{[OH^-]} = \frac{10^{-14} \text{ mole}^2/\text{liter}^2}{10^{-2} \text{ mole/liter}} = 10^{-12} \text{ mole/liter}.$$

The pH is defined as $pH = -\log [H_3O^+]$, or

$$pH = -\log [H_3O^+] = -\log (10^{-12}) = -(-12) = 12.$$


• PROBLEM 385

Find the pH of a 0.25 M solution of Na_2CO_3 , a strong electrolyte.


Solution: The crux of this problem is to realize that the solution contains the diprotic base CO_3^{2-} . Since Na_2CO_3 is a strong electrolyte, it dissociates completely. Diprotic means that 2 hydrogen ions will dissociate. Thus, there are 2 acid equilibrium constants. H_2CO_3 is the fully protonated acid. For diprotic acids, the second base ionization constant, K_{b2} , is related to the first acid ionization constant, K_{a1} , and K_{b1} is related to K_{a2} .

The first H^+ ion neutralized behaves as a stronger acid than the second H^+ ion, because in each successive stage, ionization is less complete.

The ionization and equilibrium constant equations are:




$$K_{b1} = \frac{[\text{HCO}_3^{-}][\text{OH}^{-}]}{[\text{CO}_3^{-2}]} \quad \text{and}$$



$$K_{b2} = \frac{[\text{H}_2\text{CO}_3][\text{OH}^{-}]}{[\text{HCO}_3^{-}]}$$

Since it is given that the $[\text{CO}_3^{-2}] = 0.25 \text{ M}$, attention shall be limited to the first ionization.

Each x moles of CO_3^{-2} that reacts with H_2O forms x moles each of HCO_3^{-} and OH^{-} .



Before: 0.25 0 0

After: 0.25-x x x

If the concentration of CO_3^{-2} is initially 0.25 M, and x moles of it react, then, at equilibrium, there are $0.25 - x$.

Before substituting these values into the K_{b1} equation first find K_{b1} . This is accomplished by referring to a table of K_a ionization constants. K_{a2} has a value of 5.62×10^{-11} . One can then use this information in the following equation:

$$K_{a2} K_{b1} = K_w$$

which relates the acid and base ionization constants to the constant for the autodissociation of water, K_w . Therefore,

$$K_{b1} = \frac{K_w}{K_{a2}} = \frac{1.0 \times 10^{-14}}{5.62 \times 10^{-11}} = 1.8 \times 10^{-4}$$

One can find $[\text{OH}^{-}]$ by writing the equilibrium constant expression and substituting the values for the concentrations.

$$K_{b1} = \frac{[\text{HCO}_3^{-}][\text{OH}^{-}]}{[\text{CO}_3^{-2}]}$$

$$\frac{(x)(x)}{0.25} = 1.8 \times 10^{-4}$$

$$x = [\text{OH}^{-}] = 6.7 \times 10^{-3}.$$

The x from $(0.25 - x)$ was eliminated since so few moles react, as CO_3^{-2} is a weak base.

In order to calculate pH, first obtain pOH using

$$\text{pOH} = -\log [\text{OH}^-]$$

$$= -\log [6.7 \times 10^{-3}] = 3 - \log 6.7 = 2.17.$$

$$\text{pH} = \text{p}K_w - \text{pOH} = 14 - 2.17 = 11.83.$$

It makes sense that $K_a K_b = K_w$ since K_a is a measure of $[\text{H}^+]$ and K_b is a measure of $[\text{OH}^-]$, and one knows that

$$[\text{H}^+][\text{OH}^-] = K_w = 10^{-14}.$$

• PROBLEM 386

Explain the following phenomena:

- a) Liquid HCl is a nonelectrolyte, but aqueous HCl is a strong electrolyte
 b) Liquid HCN is a nonelectrolyte as is aqueous HCN.

Solution: An electrolyte is a substance which exists as ions. Both liquid HCN and HCl are not electrolytes because they consist of neutral atoms. No ions exist. The case with an aqueous solution requires a little more investigation. In water, HCl is ionized via the reaction $\text{HCl} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{Cl}^-$. Thus, you have hydronium ions and chloride ions in solution. Thus, aqueous HCl is an electrolyte. HCN also dissociate when placed in water. But, to what extent? H_3O^+ and CN^- ions are only to a very slight extent produced, since HCN's dissociation constant is so small. Thus, while HCN may be called an electrolyte, it is an extremely weak one.

CHAPTER 11

SOLUBILITY AND THE ION PRODUCT CONSTANT

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 367 to 384 for step-by-step solutions to problems.

A solubility product or ion product constant is simply an equilibrium constant for the solubility reaction. The problems in this section can be solved by following the same principles discussed in Chapter 9 and illustrated for acid-base equilibrium in Chapter 10. If ions of several salts are present, the equilibrium solubility of each possible reaction must simultaneously be satisfied. Two examples will illustrate the application of these principles to solubility equilibrium.

First, the solution of a relatively insoluble salt such as silver chloride:




$$K_{sp} = (\text{Ag}^+) (\text{Cl}^-) / (\text{AgCl}) = 1.1 \times 10^{-10} \quad 11-2$$

The activity of solid AgCl is one if the solid is present. Also, from the stoichiometric relationship in Equation 11-1, it is apparent that the concentrations of Ag^+ and Cl^- are equal. Let this concentration = x .

$$1.1 \times 10^{-10} = (x)(x)/1 = x^2 \quad 11-3$$

$$x = 1.05 \times 10^{-5} = (\text{Ag}^+) = (\text{Cl}^-) \quad 11-4$$

Second, let us consider the concentration of silver and chloride ions in a solution initially containing 0.1M silver nitrate, a strong electrolyte, and 0.2M sodium chloride, also a strong electrolyte. Several possible ionic equilibrium reactions might be considered. The equilibrium of each must be satisfied.



The equilibrium constant K_{sp} , for strong electrolytes, is a large number and so the concentration of AgNO_3 and NaCl can be considered to be zero. The 0.2M Cl^- from Equation 11-6 will react with essentially all of the 0.1M Ag^+ from Equation 11-7 leaving a Cl^- concentration of approximately 0.1M . To calculate the (Ag^+) concentration in this case, Equation 11-2 can be employed. Note that (Cl^-) is very nearly 0.1 and the AgCl activity is one.

$$1.1 \times 10^{-10} = (\text{Ag}^+)(\text{Cl}^-)/(\text{AgCl}) = (\text{Ag}^+)(0.1)/(1) \quad 11-8$$

$$(\text{Ag}^+) = 1.1 \times 10^{-9} \quad 11-9$$

Step-by-Step Solutions to Problems in this Chapter, "Solubility and the Ion Product Constant"


• PROBLEM 387

Describe, using equations, what takes place when the following solutions are prepared:


- A. A solution containing equal amounts of 0.10M lead nitrate and 0.10M potassium chromate;
- B. A solution containing 1 mole of sodium chloride and 1 mole of potassium bromide.

Solution: In such problems, two processes can occur: mixing of ions in solution and precipitation.

- A. Both lead nitrate ($\text{Pb}(\text{NO}_3)_2$) and potassium chromate (K_2CrO_4) are highly soluble, so that four ions are present in solution:




There are four possible combinations of these ions: $\text{Pb}(\text{NO}_3)_2$, PbCrO_4 , KNO_3 , and $\text{K}_2\text{Cr}_2\text{O}_7$. Of these, only lead chromate, PbCrO_4 , is insoluble, so that the reaction



is driven to the right, and yields a precipitate, leaving K^+ and NO_3^- ions in solution (along with trace amounts of Pb^{2+} and CrO_4^{2-} , which exist in equilibrium with solid lead chromate).

- B. Both sodium chloride (NaCl) and potassium bromide (KBr) are highly soluble, giving rise to Na^+ , K^+ , Cl^- , and Br^- ions in solution. The four possible combinations of these ions are NaCl , NaBr , KCl , and KBr , all of which are highly soluble. Therefore no precipitate is formed, but only mixing of the ions according to the equation



• PROBLEM 388

Given that K_{sp} for $\text{Mg}(\text{OH})_2$ is 1.2×10^{-11} , calculate the solubility of this compound in grams per 100ml of solution. The re-

action equation is



Solution: K_{sp} is the solubility product constant; it measures the equilibrium established between the ions in the saturated solution and the excess solid phase. Knowing the K_{sp} , we can calculate the solubility of the compound. The K_{sp} equation for general compound A_xB_y is

$$K_{sp} = [\text{A}]^x[\text{B}]^y.$$

From the equation, it can be seen that if x moles per liter of Mg(OH)_2 dissolves, x moles of Mg^{+2} and $2x$ moles of OH^- form per liter,

$$K_{sp} = [\text{Mg}^{+2}][\text{OH}^-]^2$$

$$1.2 \times 10^{-11} = x(2x)^2 = 4x^3.$$

Solving,

$$x = 1.4 \times 10^{-4} \text{ mole/liter of } \text{Mg(OH)}_2 \text{ dissolved.}$$

One is asked, however, for grams per 100ml of solution. 1.4×10^{-4} mole/liter can be converted to grams/100ml by the following method:

100ml = .1 liters since there exists 1000ml = 1 liter. If one has

1.4×10^{-4} moles in 1 liter, then in .1 liters, there are $(.1)(1.4 \times 10^{-4})$

$= 1.4 \times 10^{-5}$ moles. The molecular weight of $\text{Mg(OH)}_2 = 58.312\text{g/mole}$.


Therefore, 1.4×10^{-5} moles/100ml translates into

$$(1.4 \times 10^{-5} \frac{\text{moles}}{100\text{ml}})(58.312\text{g/mole}) = 8.16 \times 10^{-4} \text{ grams/100ml.}$$

• PROBLEM 389

The solubility product constant of magnesium hydroxide is 8.9×10^{-12} , calculate its solubility in (a) water and (b) .05M NaOH.

Solution: Whenever an ionic solid is placed in water, an equilibrium is established between its ions and the excess solid phase. The solubility constant, K_{sp} , measures this equilibrium. For the general reaction $\text{A}_x\text{B}_y \rightleftharpoons x\text{A}^+ + y\text{B}^-$, the K_{sp} is defined as being equal to $[\text{A}^+]^x[\text{B}^-]^y$. The concentration of solid is always constant, no matter how much is in contact with the ions. This means the solid phase will not appear in the equilibrium constant expression. For this problem, part (a), you have



The

$$K_{sp} = [\text{Mg}^{2+}][\text{OH}^-]^2 = 8.9 \times 10^{-12}.$$

You are asked to find these concentrations. From the chemical equation, you find 2 moles of OH^- will be generated per mole of Mg^{2+} . Thus, at equilibrium, if the concentration of $\text{Mg}^{2+} = x$ (mol/liter), then the concentration of $\text{OH}^- = 2x$ (mol/liter). (Note: The dissociation of water contributes some OH^- , but this amount is very small, and can be ignored.) Thus, you have

$$[\text{Mg}^{2+}][\text{OH}^-]^2 = x(2x)^2 = 8.9 \times 10^{-12} = K_{sp}$$

Solving for x , you obtain $x = 1.3 \times 10^{-4}$ mol/liter. Thus, 1.3×10^{-4} mole of $\text{Mg}(\text{OH})_2$ dissolves per liter of water, producing a solution of 1.3×10^{-4} M Mg^{2+} and 2.6×10^{-4} M OH^- .

To find the solubility in .05M NaOH, part (b), perform the same process, except you must realize that the NaOH supplies OH^- in addition to the OH^- that dissolves from the salt. This means, as such,

$$[\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp} = (x)(2x + .050)^2 = 8.9 \times 10^{-12}$$


If you solve for x , you obtain

$$3.6 \times 10^{-9} \text{ mol/liter.}$$

• PROBLEM 390

A chemist dissolves BaSO_4 in pure water at 25°C . If its $K_{sp} = 1 \times 10^{-10}$, what is the solubility of the barium sulfate in the water?

Solution: The solubility of a compound is defined as the limiting concentration of the compound in a solution before precipitation occurs. To find the solubility of the barium sulfate, you need to know the concentration of its ions in solution. BaSO_4 will dissociate into ions because it is a salt. There will be an equilibrium between these ions and the BaSO_4 . The equilibrium can be measured in terms of a constant, K_{sp} , called the solubility constant. The K_{sp} is expressed in terms of the concentrations of the ions. As such, to answer the question, you want to represent this K_{sp} . For this reaction, the equation is



$$K_{sp} = [\text{Ba}^{2+}][\text{SO}_4^{2-}] = 1 \times 10^{-10}$$


Let $x = [\text{Ba}^{2+}]$. Thus, $x = [\text{SO}_4^{2-}]$, also, since both ions will be formed in equimolar amounts. Therefore, $x \cdot x = 1 \times 10^{-10}$. Solving,

$$x = 1 \times 10^{-5} \text{ M} = [\text{Ba}^{2+}] = [\text{SO}_4^{2-}]$$

• PROBLEM 391

Determine the approximate solubility of AgCl in 0.10M NaCl solution. K_{sp} for $\text{AgCl} = 1.1 \times 10^{-10}$.

Solution: The approximate solubility of AgCl in this solution is equal to the concentration of Ag^+ in the solution. This is because the concentration of Ag^+ is equal to the amount of AgCl ionized. The ionization of AgCl can be stated




This means that one Ag^+ ion is formed each time one molecule of AgCl is ionized. The concentration of Ag^+ can be obtained by using the

solubility product constant (K_{sp}). This constant is equal to the product of the concentration of Ag^+ ($[Ag^+]$) times the concentration of Cl^- ($[Cl^-]$).

$$K_{sp} = [Ag^+][Cl^-] = 1.1 \times 10^{-10}$$

The concentration of the Cl^- ions is also affected by the Cl^- ions from the NaCl solution. Because NaCl is a salt, it is 100% ionized in solution



There is one Cl^- ion formed for every NaCl ionized. The original concentration of NaCl is 0.10 molar, and after ionization, the solution is 0.10 molar in Cl^- ions. The Na^+ ions can be disregarded because here we are looking at the solubility of AgCl, which does not contain any Na^+ ions. One can assume that the total concentration of Cl^- ions in solution will be approximately 0.10, since the amount contributed by the AgCl ionization is small, and can be neglected.

The $[Ag^+]$ can now be found.

$$[Ag^+][Cl^-] = 1.1 \times 10^{-10}$$


$$[Ag^+](0.10) = 1.1 \times 10^{-10}$$

$$[Ag^+] = 1.1 \times 10^{-9} M$$

• PROBLEM 392

Assuming the K_{sp} for radium sulfate is 4×10^{-11} , what is its solubility in (a) pure water, and (b) .1M Na_2SO_4 ?

Solution: Whenever an ionic solid is placed in water, an equilibrium is established between its ions and the excess solid phase. A solubility constant, K_{sp} , measures this equilibrium. The concentration of solid is always constant, no matter how much it is in contact with the ions. This means the solid phase will not appear in the equilibrium constant expression. The dissociation is



The K_{sp} is, thus, $[Ra^{2+}][SO_4^{2-}] = 4 \times 10^{-11}$. From the chemical reaction, it becomes evident that the concentration of Ra^{2+} must equal that of SO_4^{2-} , since they are generated in equimolar amounts.

Therefore, let x = the solubility of each. Substituting into $[Ra^{2+}][SO_4^{2-}] = 4 \times 10^{-11}$, you have $x \cdot x = 4 \times 10^{-11}$. Solving for x , you obtain $x = 6 \times 10^{-6}$ mol/liter. Thus, the solubility of $RaSO_4$ is 6×10^{-6} mol/liter, the solution is $6 \times 10^{-6} M Ra^{2+}$ and $6 \times 10^{-6} M SO_4^{2-}$ in water.

b) The equation for this reaction in water is $RaSO_4 \rightleftharpoons Ra^{2+} + SO_4^{2-}$. Na_2SO_4 is a salt and therefore will ionize completely in water.

The concentration of the Na_2SO_4 solution is .1M, which is equivalent to a solution of .2M of Na^+ and .1M SO_4^{2-} .

If one adds more SO_4^{2-} to the system, the equilibrium is forced

to the left. Therefore, there will be even less Ra^{2+} present than in pure water. Thus,

$[\text{Ra}^{2+}] \ll 6.0 \times 10^{-6} \text{ M}$
when additional SO_4^{2-} is present. Let y = concentration of Ra^{2+} .

The solubility of RaSO_4 is equal to the product of $[\text{Ra}^{2+}]$ and $[\text{SO}_4^{2-}]$ in this solution. Solving: $[\text{Ra}^{2+}]$ is equal to y . Because RaSO_4 dissociates into Ra^{2+} and SO_4^{2-} , the RaSO_4 in this contributes y to $[\text{SO}_4^{2-}]$. There is also SO_4^{2-} already present in this solution, its concentration is .1M, therefore $[\text{SO}_4^{2-}] = (y + .1)$.

Writing the equation for the solubility
solubility of $\text{RaSO}_4 = [\text{Ra}^{2+}][\text{SO}_4^{2-}]$

$$= y(y + .1).$$

It has been shown that y is much smaller than .1 and the following approximation can be made. $y + .10 \approx .1$. Solving for y :

$$y(y + .1) = 4 \times 10^{-11}$$


$$.1y = 4 \times 10^{-11}$$

$$y = 4 \times 10^{-10}$$

• PROBLEM 393

A saturated solution of CaF_2 contains .00168g of CaF_2 per 100g of water. Determine the K_{sp} .

Solution: When a salt is added to water, it dissociates into ions. Once a saturated solution is obtained, the addition of more salt results in precipitation of the solid. In such a case, an equilibrium is established between the solid phase and the ions in solution. This equilibrium is expressed as the solubility product constant or K_{sp} . To determine the K_{sp} , you must set up an equation that expresses this equilibrium. You must also determine the molarity of the solution. Molarity refers to the number of moles per liter of solution. The equilibrium reaction is



The products have their oxidation states expressed because they are ions in solution. The K_{sp} for the reaction is defined as


$$K_{sp} = [\text{Ca}^{2+}][\text{F}^{-}]^2.$$

Because this is a heterogeneous equilibrium, there is no need to include $\text{CaF}_2(\text{s})$ in the expression for K_{sp} . You include only those substances with variable molar concentrations. The fluoride ion concentration is squared because 2 moles of fluoride ion are generated in the reaction. In general, then, the exponent of the concentration will equal the number of moles of that ion that is generated. Now that the K_{sp} is defined, determine the molarity of the solution. Since molarity is defined as moles of solute per liter of solution, you have

$$M = \text{molarity} = \frac{.00168\text{g}}{78\text{g/mole } (\text{CaF}_2)}$$

$$\frac{100 \text{ ml}}{1000\text{ml/l}}$$

Solving, you obtain $M = .214 \times 10^{-2} \text{M}$. You will notice that the 100 grams of water do not appear in this calculation. The density of water is 1g/ml . Since Density = Mass/Volume, $100\text{g H}_2\text{O}$ has a volume of 100ml . Now that you know the molarity, go back to



If $\text{CaF}_2(\text{s})$ has $M = .214 \times 10^{-2}$, then by looking at the coefficients, you find in solution

$$2.14 \times 10^{-3} \text{M Ca}^{2+} \text{ and } 4.28 \times 10^{-3} \text{M F}^{-}.$$

These results are obtained because for every mole/liter of CaF_2 in solution, 1 mole/liter of Ca^{2+} and 2 moles/liter of F^{-} is obtained.

Therefore, recalling

$$K_{sp} = [\text{Ca}^{2+}][\text{F}^{-}]^2,$$

you have

$$\begin{aligned} K_{sp} &= (.214 \times 10^{-2})(.428 \times 10^{-2})^2 \\ &= 3.92 \times 10^{-8}. \end{aligned}$$


• PROBLEM 394

Calculate the solubility product constant of pure PbSO_4 in water. The solubility of PbSO_4 in water at 25°C is 1.25×10^{-4} moles/liter.

Solution: The solubility of a substance is the number of moles of that substance which will dissolve in one liter. Since PbSO_4 contains one Pb^{2+} ion and one SO_4^{2-} ion per molecule in aqueous solution, the concentration of each of these ions is equal to the solubility. Hence,

$$[\text{Pb}^{2+}] = [\text{SO}_4^{2-}] = 1.25 \times 10^{-4} \text{M}.$$

The solubilization reaction for PbSO_4 is



and the solubility product constant is

$$K_{sp} = [\text{Pb}^{2+}][\text{SO}_4^{2-}].$$


Substituting the values for $[\text{Pb}^{2+}]$ and $[\text{SO}_4^{2-}]$ gives

$$\begin{aligned} K_{sp} &= (1.25 \times 10^{-4} \text{M}) \times (1.25 \times 10^{-4} \text{M}) \\ &= 1.6 \times 10^{-8} \text{ moles}^2/\text{liter}^2. \end{aligned}$$

• PROBLEM 395

A chemist dissolves PbS in water. The K_{sp} is found to be 1×10^{-28} . What is the molarity of each ion in solution?

Solution: The key to answering this question is the determination of the equilibrium equation. Since PbS is a salt, it dissociates into ions in solution. Upon inspection of the oxidation states of the periodic table, the equilibrium will be found to be



The presence of the equilibrium derives from the fact that a saturated salt solution's ions will exist in equilibrium with any additional salt. This additional salt is called the solid phase. K_{sp} measures this equilibrium and can be expressed as

$$K_{sp} = [\text{Pb}^{2+}][\text{S}^{2-}]$$

This, equals 1×10^{-28} . Therefore, you have $1 \times 10^{-28} = [\text{Pb}^{2+}][\text{S}^{2-}]$. The question asks for the concentrations represented on the right side. From the equilibrium equation, you see that whatever the concentration of S^{2-} , it will be equal to Pb^{2+} . Therefore, you can represent both by x . As such, you now have

$$1 \times 10^{-28} = x \cdot x$$

or

$$1 \times 10^{-28} = x^2$$

Solving for x you obtain $1 \times 10^{-14}\text{M}$. This means, therefore, that the concentration of each ion in solution is $1 \times 10^{-14}\text{M}$.

• PROBLEM 396

A chemist has a saturated solution of CaSO_4 in $.0100\text{M}$ Na_2SO_4 . The K_{sp} of CaSO_4 is 2.4×10^{-5} , calculate the concentration of Ca^{2+} ion in this saturated solution.

Solution: The K_{sp} is the solubility constant of a substance. For the general reaction, $\text{AB}_{(s)} \rightleftharpoons \text{A}^+ + \text{B}^-$, $K_{sp} = [\text{A}^+][\text{B}^-]$. The K_{sp} = the product of the concentrations of the ions. In this problem, $K_{sp} = [\text{Ca}^{++}][\text{SO}_4^{--}]$. You are given that $K_{sp} = 2.4 \times 10^{-5}$. Equating, you obtain $2.4 \times 10^{-5} = [\text{Ca}^{++}][\text{SO}_4^{--}]$. To solve this problem, you must determine the value of $[\text{Ca}^{++}]$. Let x = the moles/liter of CaSO_4 . Since $\text{CaSO}_4 \rightleftharpoons \text{Ca}^{++} + \text{SO}_4^{--}$, this x also is the concentration of Ca^{++} . The concentration of SO_4^{--} , however, has two sources. SO_4^{--} comes from CaSO_4 AND Na_2SO_4 . The concentration of Na_2SO_4 was given as $.01$. Therefore, the total concentration of SO_4^{--} is $x + .01$, where x represents the amount of SO_4^{--} contributed by CaSO_4 . Thus,


$$[x][.01 + x] = 2.4 \times 10^{-5}$$

Solving for x you obtain,

$$x = 2.0 \times 10^{-3} = [\text{Ca}^{++}]$$

How many grams of $\text{Ca}(\text{C}_2\text{O}_4)$ (calcium oxalate) will dissolve in water to form 1.0 liter of saturated solution? The K_{sp} of $\text{Ca}(\text{C}_2\text{O}_4)$ is $2.5 \times 10^{-9} \text{ mole}^2 / \text{liter}^2$.

Solution: This problem is an application of the solubility product constant (K_{sp}) expression for dissociation of a solid. In general, if a solid, A , dissociates into ions B, C, D, \dots according to the equation




the ion product constant (I.P.) is given by

$$\text{I.P.} = [\text{B}]^b [\text{C}]^c [\text{D}]^d \dots$$

At the point where addition of solid will result in precipitation, the ion product constant is equal to the solubility product constant. (When $\text{I.P.} > K_{sp}$, precipitation will occur.) Thus, for a solution just on the verge of precipitation,

$$K_{sp} = [\text{B}]^b [\text{C}]^c [\text{D}]^d \dots$$

The dissociation of calcium oxalate is given by the equation



and the K_{sp} is

$$K_{sp} = 2.5 \times 10^{-9} \text{ mole}^2 / \text{liter}^2 = [\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}]$$

Let the concentration of the calcium ion be x . Then, since one $\text{C}_2\text{O}_4^{2-}$ ion is produced for every Ca^{2+} ion produced, $[\text{Ca}^{2+}] = [\text{C}_2\text{O}_4^{2-}] = x$.

Substituting into the expression for K_{sp} gives

$$[\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}] = 2.5 \times 10^{-9} \text{ mole}^2 / \text{liter}^2$$

$$(x)(x) = 2.5 \times 10^{-9} \text{ mole}^2 / \text{liter}^2$$

$$x = (2.5 \times 10^{-9} \text{ mole}^2 / \text{liter}^2)^{1/2}$$

$$x = 5.0 \times 10^{-5} \text{ mole/liter} = 5.0 \times 10^{-5} \text{ M}$$

Thus, the concentration of $\text{Ca}(\text{C}_2\text{O}_4)$ will be $5.0 \times 10^{-5} \text{ M}$. To convert to mass, we use the relationship (mass = concentration \times volume \times molecular weight). The molecular weight of $\text{Ca}(\text{C}_2\text{O}_4)$ is 128g/mole. Hence, the required mass of $\text{Ca}(\text{C}_2\text{O}_4)$ is, for one liter of solution,

$$\text{mass} = \text{concentration} \times \text{volume} \times \text{molecular weight}$$

$$= 5.0 \times 10^{-5} \text{ M} \times 1 \text{ L} \times 128 \text{ g/mole}$$


$$= 0.0064 \text{ g}$$

Calculate the concentration of calcium ion and fluoride ion in a concentrated solution of calcium fluoride, CaF_2 ($K_{sp} = 4.00 \times 10^{-11}$).

Solution: Problems of this type are solved by making use of the fact that the K_{sp} , which is known, is equal to the product of the concentrations of Ca^{2+} and F^- . If we know the concentration of one ion then, from the proportions in which the ions occur in the molecule,

we can determine the concentrations of all the ions formed by dissociation of the molecule.

When calcium fluoride dissociates,



one ion of Ca^{2+} and two ions of F^- form per molecule of CaF_2 . Hence, if we let x denote the concentration of calcium ions

$([\text{Ca}^{2+}] = x)$, the concentration of fluoride ions is twice as great, $([\text{F}^-] = 2x)$. The expression for the solubility product constant is

$$K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2 = 4.00 \times 10^{-11}$$

Substituting $[\text{Ca}^{2+}] = x$ and $[\text{F}^-] = 2x$ gives

$$(x)(2x)^2 = 4.00 \times 10^{-11}$$

$$(x)(4x^2) = 4.00 \times 10^{-11}$$

$$4x^3 = 4.00 \times 10^{-11}$$

or,


$$x = \left(\frac{4.00 \times 10^{-11}}{4} \right)^{1/3} = 2.15 \times 10^{-4}$$

Hence, $[\text{Ca}^{2+}] = x = 2.15 \times 10^{-4}$ and $[\text{F}^-] = 2x = 2 \times 2.15 \times 10^{-4}$
 $= 4.3 \times 10^{-4}$

● PROBLEM 399

You are given .01 moles of AgCl . Assuming the K_{eq} for $\text{Ag}(\text{NH}_3)_2^+$ is 1.74×10^7 , how much 1M ammonia solution is required to dissolve this AgCl ? Assume, also, K_{sp} of AgCl is 2.8×10^{-10} .

Solution: The first thing to do is to write the overall reaction that occurs when ammonia is mixed with AgCl . From this, write an equilibrium expression and equate this expression with the equilibrium constant. Next, calculate the concentrations of species in solution. After this point, there is a volume determination. Following this procedure: when AgCl is dissolved in a solution, it dissociates into ions, since it is a salt. The reaction is $\text{AgCl}(s) \rightleftharpoons \text{Ag}^+ + \text{Cl}^-$. The $K_{sp} = 2.8 \times 10^{-10}$. The silver ion, Ag^+ , produced will react with ammonia (NH_3) to produce the silver ammonia complex, $\text{Ag}(\text{NH}_3)_2^+$ via the reaction $\text{Ag}^+ + 2\text{NH}_3 \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+$ with $K = 1.74 \times 10^7$. The overall reaction is the sum of these two. The K for this will be the product of the K 's for the two reactions just written. Thus, overall reaction is:



with

$$K = (2.8 \times 10^{-10})(1.74 \times 10^7) = 4.9 \times 10^{-3}$$

The equilibrium expression indicates the ratio of products to reactants, each raised to the power of their coefficients in the overall reaction. Thus,

$$K = 4.9 \times 10^{-3} = \frac{[\text{Ag}(\text{NH}_3)_2^+][\text{Cl}^-]}{[\text{NH}_3]^2}$$

One must now find these concentrations. Let $x = [\text{Ag}(\text{NH}_3)_2^+]$, thus $x = [\text{Cl}^-]$, since the overall reaction indicates they are formed in equimolar amounts. Since 2 moles of NH_3 is consumed per mole of complex, $[\text{NH}_3] = 1 - 2x$, where 1 is the original molarity of the ammonia solution. Substituting and solving for x ,

$$4.9 \times 10^{-3} = \frac{x \cdot x}{1 - 2x};$$

thus,

$$x = .065\text{M}.$$

From the initial .01 moles of AgCl , .01 moles of Cl^- must be produced. This concentration of Cl^- is .065M. Since $M = \text{molarity} = \text{moles/volume}$, the volume required to dissolve the AgCl is

$$\frac{.01 \text{ moles}}{.065 \text{ moles/liter}} = .154 \text{ liters} = 154 \text{ ml}.$$

• PROBLEM 400

Calculate the minimum concentration of Br^- ion necessary to bring about precipitation of AgBr from a solution in which the concentration of Ag^+ ion is 1×10^{-5} mole per liter.

$$K_{sp} \text{ for } \text{AgBr} = 4 \times 10^{-13}.$$

Solution: The product of the concentrations of ions in a saturated solution of a relatively insoluble salt, such as AgBr , at a given temperature is constant. This constant is called the solubility product constant (K_{sp}). For AgBr , K_{sp} can be stated as

$$K_{sp} = [\text{Ag}^+][\text{Br}^-] = 4 \times 10^{-13}.$$

In this problem, we are given the concentration of Ag^+ as $1 \times 10^{-5}\text{M}$, therefore $[\text{Br}^-]$ can be found by using the equation for K_{sp} .

$$\begin{aligned} [\text{Ag}^+][\text{Br}^-] &= 4 \times 10^{-13} \\ [1 \times 10^{-5}][\text{Br}^-] &= 4 \times 10^{-13} \\ [\text{Br}^-] &= \frac{4 \times 10^{-13}}{1 \times 10^{-5}} = 4 \times 10^{-8}\text{M}. \end{aligned}$$

If the concentration of Br^- ions is raised to $4 \times 10^{-8}\text{M}$, AgBr will precipitate out, for the addition of any more Br^- ions could not be supported by the solubility of the ions in solution, as the K_{sp} indicates.

• PROBLEM 401

If you mix 50ml of $5.0 \times 10^{-4}\text{M}$ $\text{Ca}(\text{NO}_3)_2$ and 50ml of $2.0 \times 10^{-4}\text{M}$ NaF to give 100ml of solution, will precipitation occur? The K_{sp} of CaF_2 is 1.7×10^{-10} .

Solution: Whether or not precipitation will occur, when these two solutions are mixed, depends upon the ion product. If two solutions

containing the ions of a salt are mixed, and if the ion product exceeds the K_{sp} , then precipitation will occur.

You need to determine the concentrations of Ca^{2+} and F^- and to see if the product exceeds the given K_{sp} of 1.7×10^{-10} for CaF_2 . The equation for this reaction is $\text{CaF}_2(s) \rightleftharpoons \text{Ca}^{2+} + 2\text{F}^-$. Thus, the $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$. You are told that a 50ml solution of $\text{Ca}(\text{NO}_3)_2$ has a molarity of $5 \times 10^{-4}\text{M}$. Each $\text{Ca}(\text{NO}_3)_2$ dissociates to yield one Ca^{++} ion which means the molarity of Ca^{++} is also $5 \times 10^{-4}\text{M}$. However, it is $5 \times 10^{-4}\text{M}$ in 50ml, and the mixture is 100ml. Therefore, the molarity must be divided by two. The molarity of NaF , 2.0×10^{-4} , and therefore Na^+ , because of its dissociation, must also be divided by two for the same reasons. As such, in the mixture,

$$[\text{Ca}^{2+}] = 2.5 \times 10^{-4}\text{M}$$

and

$$[\text{F}^-] = 1.0 \times 10^{-4}\text{M}.$$

Recalling that $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$, you have

$$K_{sp} = (2.5 \times 10^{-4})(1.0 \times 10^{-4})^2 = 2.5 \times 10^{-12}.$$

You were given that the K_{sp} for a saturated solution of CaF_2 is 1.7×10^{-10} . Since 2.5×10^{-12} is less than 1.7×10^{-10} , the mixture is unsaturated, and precipitation should NOT occur.

• PROBLEM 402

Will precipitation occur if .01 mole of Ba^{+2} is added to a liter of solution containing .05 mole of SO_4^{2-} ? K_{sp} for $\text{BaSO}_4 = 1 \times 10^{-10}$.

Solution: K_{sp} is the solubility product constant. It is defined as the minimum K_{sp} product of the concentration of the ions needed to bring about the formation of a precipitate (i.e., a solid compound). The general form of the K_{sp} equation for compound A_nB_m is:

$$K_{sp} = [\text{A}]^n[\text{B}]^m$$

Therefore, for BaSO_4 the K_{sp} equation is:

$$K_{sp} = [\text{Ba}^{+2}][\text{SO}_4^{-2}] = 1 \times 10^{-10}.$$


To find out if precipitation occurs, substitute in the given concentration values. If the calculated value is greater than the given value for K_{sp} , then precipitation occurs. If it is lower, then no precipitation occurs. Thus,

$$\begin{aligned} K_{sp} &= [\text{Ba}^{+2}][\text{SO}_4^{-2}] \\ &= [1 \times 10^{-2}][5 \times 10^{-2}] \\ &= 5 \times 10^{-4}. \end{aligned}$$

Since this value is greater than the given K_{sp} value, precipitation occurs.

A chemist has a solution which contains Zn^{2+} and Cu^{2+} , each at .02M. If the solution is made 1M in H_3O^+ and H_2S is bubbled in until the solution is saturated, will a precipitate form?
 K_{sp} of $\text{ZnS} = 1 \times 10^{-22}$, K_{sp} of $\text{CuS} = 8 \times 10^{-37}$, and K_{eq} for H_2S ; when saturated, $= 1 \times 10^{-22}$.

Solution: A precipitate will form only when the ion product exceeds the experimentally determined solubility product constant (K_{sp}). Therefore, compute the ion product of ZnS and CuS to determine if they exceed the stated K_{sp} 's, 1×10^{-22} and 8×10^{-37} , respectively. The equation for the reaction between H_2S and H_2O is



In a saturated solution, the ion product is equal to

$$[\text{H}_3\text{O}^+]^2[\text{S}^{2-}] = 1 \times 10^{-22} = K.$$

You are told that the H_3O^+ concentration is 1M. Substituting this into that expression, and solving for $[\text{S}^{2-}]$, you obtain

$$[\text{S}^{2-}] = 1 \times 10^{-22} \text{M}.$$

You are given the Zn^{2+} and Cu^{2+} concentrations, which means that the ion products can be computed. For Zn^{2+} ,

$$\begin{aligned} [\text{Zn}^{2+}][\text{S}^{2-}] &= (.02)(1 \times 10^{-22}) = 2 \times 10^{-24} \\ &= K_{sp} \text{ for ZnS.} \end{aligned}$$

$$\begin{aligned} [\text{Cu}^{2+}][\text{S}^{2-}] &= (.02)(1 \times 10^{-22}) = 2 \times 10^{-24} \\ &= K_{sp} \text{ for CuS.} \end{aligned}$$

Since the ion product of ZnS does not exceed the K_{sp} of ZnS , 1×10^{-22} , there will be no precipitation. However, the ion product of CuS does exceed the K_{sp} of CuS , 8×10^{-37} , which means you will see precipitation of CuS .

• PROBLEM 404

A chemist mixes equal volumes of .01M $\text{Na}_2\text{C}_2\text{O}_4$ and .001M BaCl_2 together. Assuming the K_{sp} of BaC_2O_4 is 1.2×10^{-7} , will a precipitate form?

Solution: A precipitate will only form when the ion concentration product for the particular substances exceed their given K_{sp} . In other words, a precipitate will form if $[\text{Ba}^{2+}][\text{C}_2\text{O}_4^{2-}]$ is greater than the $K_{sp} = 1.2 \times 10^{-7}$. Thus, to obtain this ion product, calculate the concentrations of Ba^{2+} and $\text{C}_2\text{O}_4^{2-}$ in the mixture. One is given that the initial concentrations of $\text{C}_2\text{O}_4^{2-}$ and Ba^{2+} in the solution are .01M $\text{Na}_2\text{C}_2\text{O}_4$ and .001M BaCl_2 . When they are mixed, however, the volume doubles. Thus, in the mixture, the con-

centrations of BaCl_2 and $\text{Na}_2\text{C}_2\text{O}_4$ are halved. (Remember, $M =$ moles/liter, so that if the volume increases, $M =$ concentration decreases). Therefore, after mixing, the concentrations are .005M of $\text{Na}_2\text{C}_2\text{O}_4$ and .0005M of BaCl_2 . As such, the ion product of

$$[\text{Ba}^{2+}][\text{C}_2\text{O}_4^{2-}] = (.0005)(.005) = 2.5 \times 10^{-6}.$$

2.5×10^{-6} is greater than the K_{sp} of 1.2×10^{-7} , which indicates that a precipitate will form.

• PROBLEM 405

0.01 liter of 0.3M Na_2SO_4 is mixed with 0.02 liters of a solution that is initially 0.1M Ca^{++} and 0.1M Sr^{++} . Given that $K_{sp} = 2.4 \times 10^{-5}$ for CaSO_4 and $K_{sp} = 7.6 \times 10^{-7}$ for SrSO_4 , what is the final concentration of Ca^{++} , Sr^{++} , and SO_4^{--} in this solution?

Solution: Solve this problem in four steps:

- (A) Find the relationships involving the final concentrations of Ca^{++} , SO_4^{--} , and Sr^{++} .
- (B) Find the initial concentrations of the three species.
- (C) Relate the final concentrations to the initial concentrations.
- (D) Solve for $[\text{Ca}^{++}]$, $[\text{SO}_4^{--}]$, and $[\text{Sr}^{++}]$.

(A) K_{sp} , the solubility product constant, equals the product of the concentrations of the ions in a saturated solution. Thus, $K_{sp} \text{ CaSO}_4 = [\text{Ca}^{++}][\text{SO}_4^{--}]$ and $K_{sp} \text{ SrSO}_4 = [\text{Sr}^{++}][\text{SO}_4^{--}]$. Combine this with the given:

$$(i) \quad [\text{Ca}^{++}][\text{SO}_4^{--}] = 2.4 \times 10^{-5}$$

$$(ii) \quad [\text{Sr}^{++}][\text{SO}_4^{--}] = 7.6 \times 10^{-7}$$

Equation (i) relates $[\text{Ca}^{++}]$ and $[\text{SO}_4^{--}]$; equation (ii) relates $[\text{Sr}^{++}]$ and $[\text{SO}_4^{--}]$. To find an equation relating $[\text{Ca}^{++}]$ and $[\text{Sr}^{++}]$, solve (i) for $[\text{SO}_4^{--}]$:

$$[\text{SO}_4^{--}] = \frac{2.4 \times 10^{-5}}{[\text{Ca}^{++}]}$$

Substitute this into (ii); and solve for $[\text{Sr}^{++}]$.

$$(iii) \quad [\text{Sr}^{++}] = \frac{7.6 \times 10^{-7}}{2.4 \times 10^{-5}} [\text{Ca}^{++}] = 0.032 [\text{Ca}^{++}].$$

(B) 0.01 liters of 0.3M Na_2SO_4 was initially mixed. The number of moles of Na_2SO_4 present is, therefore, (molarity)(volume) = $(0.3)(0.01) = 0.003$ moles of Na_2SO_4 . Since the salt dissociates completely, there are 0.003 moles of SO_4^{--} in the solution. The new volume is $(0.02 + 0.01)$ liters = 0.03L, after adding the Ca^{++} and Sr^{++} solution. Thus, the initial molarity of SO_4^{--} is:

$$\frac{0.003 \text{ moles}}{0.03 \text{ liter}} = 0.1 \text{ M.}$$

By similar analysis, it follows that the initial concentration of Sr^{++} = initial concentration of Ca^{++} = 0.0667M.

(C) Let the final concentration of Ca^{++} equal x . Then the final concentration of Sr^{++} , according to equation (iii), must be 0.032 $[\text{Ca}^{++}]$ = .032. The change in Ca^{++} concentration is $0.0667 - x$; the change in Sr^{++} is $0.0667 - 0.032x$.

To find the final $[\text{SO}_4^{--}]$, note that one SO_4^{--} is lost for each Ca^{++} that reacts and one SO_4^{--} is lost for each Sr^{++} that reacts. Thus the change in $[\text{SO}_4^{--}]$ is the sum of the changes in $[\text{Ca}^{++}]$ and $[\text{Sr}^{++}]$. Thus, change in $[\text{SO}_4^{--}] = (0.0667 - x) + (0.0667 - .032x) = (0.1334 - 1.032x)$. Since the final $[\text{SO}_4^{--}]$ = initial $[\text{SO}_4^{--}]$ minus change in $[\text{SO}_4^{--}]$, the final $[\text{SO}_4^{--}]$ equal $0.10 - (0.1334 - 1.032x)$ or

$$(iv) \quad [\text{SO}_4^{--}] = 1.032x - 0.0334$$

(D) Substitute the final values for $[\text{Ca}^{++}]$ and $[\text{SO}_4^{--}]$ in equation (i) and solve for x :

$$(v) \quad [\text{Ca}^{++}][\text{SO}_4^{--}] = 2.9 \times 10^{-5}$$

$$(x) \quad (1.032x - 0.0334) = 2.4 \times 10^{-5}$$

$$(vi) \quad 1.032x^2 - 0.0334x - 2.4 \times 10^{-5} = 0.$$

Use the quadratic formula

$$(vii) \quad x = \frac{0.0334 \pm \sqrt{(0.0334)^2 - 4(-2.4 \times 10^{-5})(1.032)}}{2(1.032)}$$

Thus, $x = 3.30 \times 10^{-2} \text{ M} = [\text{Ca}^{++}]$. Substitute this value in equation (iii) to obtain

$$(viii) \quad [\text{Sr}^{++}] = .032 [\text{Ca}^{++}] = .032(3.30 \times 10^{-2}) \\ = 1.045 \times 10^{-3} \text{ M.}$$

Use this value in equation (ii)

$$(ix) \quad [\text{SO}_4^{--}] = \frac{7.6 \times 10^{-7}}{[\text{Sr}^{++}]} = \frac{7.6 \times 10^{-7}}{1.045 \times 10^{-3}} = 7.15 \times 10^{-4} \text{ M.}$$

• PROBLEM 406

The solubility products of $\text{Fe}(\text{OH})_2$ and $\text{Fe}(\text{OH})_3$ are 10^{-17} and 10^{-38} , respectively. If the concentrations of Fe^{2+} and Fe^{3+} are each 10^{-5} M , at what pH will each hydroxide just begin to precipitate?

Solution: To solve this problem, set up the solubility product equations for $\text{Fe}(\text{OH})_2$ and $\text{Fe}(\text{OH})_3$. Once this is done, substitute the values of Fe^{2+} and Fe^{3+} concentrations and solve for the OH^- ion concentrations. Since $\text{pH} + \text{pOH} = 14$, then $\text{pH} = 14 - \text{pOH}$, where $\text{pOH} = -\log[\text{OH}^-]$. One has, therefore,

$$[\text{Fe}^{2+}][\text{OH}^-]^2 = 10^{-17} \text{ and}$$

$$[\text{Fe}^{3+}][\text{OH}^-]^3 = 10^{-33}.$$

For precipitation to occur in these solutions $[\text{Fe}^{2+}][\text{OH}^-]^2 \geq 10^{-17}$

and $[\text{Fe}^{3+}][\text{OH}^-]^3 \geq 10^{-33}$. When the product of the concentrations is equal to zero the concentrations are just sufficient for precipitation to occur. Solving for the pH at these concentrations:

For $\text{Fe}(\text{OH})_2$,

$$[\text{OH}^-]^2 = \frac{10^{-17}}{[\text{Fe}^{2+}]} = \frac{10^{-17}}{10^{-5}} = 10^{-12},$$

so that

$$[\text{OH}^-] = 10^{-6}$$

therefore, $\text{pOH} = 6$ and thus $\text{pH} = 8$.

For $\text{Fe}(\text{OH})_3$,

$$[\text{OH}^-]^3 = \frac{10^{-33}}{[\text{Fe}^{3+}]} = \frac{10^{-33}}{10^{-5}} = 10^{-33},$$

so that

$$[\text{OH}^-] = 10^{-11}.$$

Consequently,

$$\text{pOH} = 11 \text{ and } \text{pH} = 3.$$

• PROBLEM 407

Calculate the solubility of carbon dioxide in water at 25°C , where the partial pressure of CO_2 over the solution is 760 torr., using Henry's law constant, K_{CO_2} at $25^\circ\text{C} = 1.25 \times 10^6$. Assume that a liter of solution contains 1000 grams of water.

Solution: Raoult's law states that the partial pressure, P_2 , of the component present at lower concentration is directly proportional to its mole fraction, X_2 , for dilute solutions. Those that don't usually obey Henry's law,

$$P_2 = X_2 K_2.$$

The subscript 2 indicates that the solute (the component at lower concentration) is being considered. The constant K_2 is referred to as Henry's law constant. For nonideal solutions, Henry's law holds for the solute in the same range where Raoult's law holds for the solvent. For ideal solutions $K_2 = P_2^\circ$ and Henry's law becomes identical with Raoult's law.

To solve this problem one must know three things:

- 1) the value of K_{CO_2} , equal to 1.25×10^6 ;
- 2) the partial pressure P_2 , equal to 760 torr;
- 3) the mole fraction of CO_2 in water, which has to be determined.

The mole fraction of CO_2 in water is

$$x_{\text{CO}_2} = \frac{\text{number of moles of } \text{CO}_2}{\text{number of moles of } \text{CO}_2 + \text{number of moles of } \text{H}_2\text{O}}$$

The number of moles of H_2O is $\frac{1000\text{g}}{18.02\text{g/mole}} = 55.49 \text{ moles } \text{H}_2\text{O}$.

The number of moles of CO_2 has yet to be determined. Thus substituting into Henry's law

$$x_2 = \frac{P_2}{K_2} = \frac{760 \text{ torr}}{1.25 \times 10^6} = \frac{(\text{moles } \text{CO}_2)}{(\text{moles } \text{CO}_2) + (55.49 \text{ moles } \text{H}_2\text{O})}$$

However, since the number of moles of CO_2 may be considered negligible in comparison with the number of moles of water, then,

$$\frac{760 \text{ torr}}{1.25 \times 10^6} = \frac{\text{moles } \text{CO}_2}{55.49 \text{ moles } \text{H}_2\text{O}}$$

Rewriting and solving

$$\text{no. of moles of } \text{CO}_2 = \frac{(55.49)(760)}{1.25 \times 10^6} = 3.37 \times 10^{-2}$$

There are 1000g of H_2O , or 1 liter of solvent, and therefore, the solubility of

$$\text{CO}_2 = \frac{3.37 \times 10^{-2} \text{ moles}}{1 \text{ liter}} = 3.37 \times 10^{-2} \text{ M}$$

• PROBLEM 408

What is the equilibrium concentration of oxygen gas in fresh water at 25°C when exposed to air at a pressure of 760 torr? The Henry's law constant for oxygen in water at 25°C is 3.30×10^7 torrs. Express your answer in milligrams of oxygen per liter.

Solution: Solving this problem requires the use of Henry's law. It is stated

$$P_2 = x_2 K_2$$

where P_2 is the partial pressure, x_2 is the mole fraction, and K_2 is the Henry's law constant. The subscript, 2, indicates the solute (or component at lower concentration).

Since $P_2 = 760 \text{ torr}$ and $K_2 = 3.30 \times 10^7 \text{ torrs}$ are known, we can solve for x_2 or the mole fraction of O_2 in water. Thus,

$$x_2 = \frac{P_2}{K_2} = \frac{760 \text{ torr}}{3.3 \times 10^7 \text{ torr}} = \frac{\text{moles of } \text{O}_2}{(\text{moles of } \text{O}_2) + (\text{moles of } \text{H}_2\text{O})}$$

The volume is assumed to be 1 liter or 1000 grams of H_2O . The number of moles of H_2O is $\frac{1000\text{g}}{18.02\text{g/mole}} = 55.49 \text{ moles } \text{H}_2\text{O}$. In addition, the number of moles of O_2 is negligible in comparison with the number of moles of H_2O , thus the number of moles of H_2O is

$$\frac{(760 \text{ torr})(55.49 \text{ moles H}_2\text{O})}{3.3 \times 10^7 \text{ torr}} = 1.28 \times 10^{-3} \text{ moles O}_2 .$$

The weight in milligrams of O_2 is

$$(1.28 \times 10^{-3} \text{ moles})(32 \text{ g/mole})(10^3 \text{ mg/g}) = 41 \text{ mg}.$$

Since there is 1 liter of H_2O , then the concentration of O_2 is $41 \text{ mg/l} .$

• PROBLEM 409

The solubility of iodine in pure water is 0.0279g per 100g of water. In a solution that is originally 0.10M KI, it is possible to dissolve, at the maximum, 1.14g of iodine per 100g of solution. What is K for the reaction $\text{I}_2 + \text{I}^- \rightleftharpoons \text{I}_3^-$?

Solution: The concentration of iodine in the given solution is greater than can be dissolved. However, the wording of the problem indicates that there is no precipitate. Thus, the excess iodine must be used up in the production of I_3^- which is soluble. With this in mind, we can solve for K .

The equilibrium constant K can be solved from the following equation,

$$(1) \quad K = \frac{[\text{I}_3^-]}{[\text{I}_2][\text{I}^-]} .$$

First, solve for the concentrations of I_3^- , I_2 , and I^- .

The I^- species comes entirely from the dissociation of KI .

Since a salt such as KI totally dissociates in aqueous solution, initial molarity of I^- = molarity of KI = 0.10M .

The initial amount of I_2 added is 1.14g per 100g of water or the number of moles added equals

$$\frac{\text{grams of I}_2}{\text{mol.wt. of I}_2} = \frac{1.14 \text{ g}}{254 \text{ g/mole}} = 4.49 \times 10^{-3} \text{ moles}$$

per 100g of water. However, the water can only hold 0.0279g of iodine per 100g of water, or

$$\frac{0.0279 \text{ g}}{254 \text{ g/mole}} = 1.1 \times 10^{-4} \text{ moles per 100g of water}.$$

Since there is to be no precipitate, the excess I_2 must be used up in the production of I_3^- . Since one mole I_2 yields one mole of I_3^- , we have for the final concentrations:

- (1) concentration of I_2 = maximum concentration of I_2 that can be dissolved = 1.1×10^{-4} moles per 100g of water. Since 1000g of $\text{H}_2\text{O} = 1 \text{ l}$, 1.1×10^{-4} moles/100g $\text{H}_2\text{O} = \frac{1.1 \times 10^{-3} \text{ moles}}{1000 \text{ g H}_2\text{O}} =$

$$\frac{1.1 \times 10^{-3} \text{ moles}}{1 \text{ liter H}_2\text{O}} \quad . \quad \text{Thus, } [I_2] = 1.1 \times 10^{-3} \text{ M} .$$

- (2) concentration of I_3^- = concentration of I_2 reacted. Since there were originally 4.49×10^{-3} moles of $I_2/100\text{g H}_2\text{O}$ and the final concentration of I_2 is 1.1×10^{-4} moles/100g H_2O , the concentration of I_2 that reacted is the difference

$$\frac{4.49 \times 10^{-3} \text{ moles}}{100\text{g H}_2\text{O}} - \frac{1.1 \times 10^{-4} \text{ moles}}{100\text{g H}_2\text{O}} = \frac{4.38 \times 10^{-3} \text{ moles}}{100\text{g H}_2\text{O}}$$

$$= \frac{4.38 \times 10^{-2} \text{ moles}}{1 \text{ liter H}_2\text{O}} \quad . \quad \text{Thus, } [I_3^-] = 4.38 \times 10^{-2} \text{ M} .$$

- (3) concentration of I^- = initial concentration I^- minus (concentration of I^- needed to react with the I_2). From (2), the concentration of reacted I_2 was found to be $4.38 \times 10^{-2} \text{ M}$.

Since one mole of I^- is required to react with each mole of I_2 , the concentration of I^- needed to react with the I_2 is also $4.38 \times 10^{-2} \text{ M}$.

$$\text{Thus, } [I^-] = 0.10\text{M} - 4.38 \times 10^{-2} \text{ M} = .0562\text{M} .$$

To find K, substitute these values in equation (1):

$$K = \frac{(0.0438)}{(0.0011)(0.0562)} = 710 .$$

CHAPTER 12

CALCULATIONS USING pH AND DISSOCIATION CONSTANT

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 385 to 421 for step-by-step solutions to problems.

The principles of equilibrium that have been discussed in the preceding chapters (Chapters 9, 10, and 11) can be applied directly to solve the problems in this chapter. The dissociation constant is simply the equilibrium constant for a dissociation reaction. When the H^+ ion concentration is determined from an equilibrium calculation, the pH can be calculated directly from Equation 10-3.

$$pH = -\log_{10}(H^+) \quad 12-1 \text{ (also 10-3)}$$

There are also problems in Chapter 12 that ask for values of pK , pK_a , pK_b , etc. The key to calculating these quantities lies in the definition of pK , which is analogous to the definition of pH.

$$pK_i = -\log_{10}(K_i) \quad 12-2$$

Dissociation is the ionization or dissolution of an electrolyte and is applied most often to weak electrolytes such as organic acids. Several of the problems in this chapter involve calculation of the concentration of H^+ ions resulting from the dissociation of acetic acid. The equilibrium dissociation of acetic acid was outlined in some detail in Equations 10-10 through 10-12 when discussing buffered solutions. Another example of acetic acid dissociation without buffering will illustrate the essential technique for solving such problems. Let us solve for the pH in a 0.1 molar solution of acetic acid.



$$K_a = 1.8 \times 10^{-5} = (H^+)(Ac^-)/(HAc) \quad 12-4$$

The key to solving this problem lies in making two observations. First, the stoichiometry of Equation 12-3 dictates that, in the absence of other sources of H^+ and Ac^- ions, the concentrations of these two ions will be equal. Second, since the equilibrium constant is a small number, the concentration of undissociated HAc will not be appreciably reduced by the dissociation reaction. We assume that the concentration, (HAc) , is equal to the total HAc before dissociation. (We will rework this problem without making this assumption to see if the assumption is valid.)

The concentrations of H^+ and Ac^- can be assigned a new variable, x . Equation 12-4 becomes

$$1.8 \times 10^{-5} = (x)(x)/(0.1) = 10x^2 \quad 12-5$$

$$x = (1.8 \times 10^{-6})^{1/2} = 0.00134 = (H^+) = (Ac^-) \quad 12-6$$

$$pH = -\log_{10}(0.00134) = 2.87 \quad 12-7$$

Note that the pH would be 1 $[-\log(0.1)]$ had dissociation been complete.

Now let us illustrate the same calculation **without** making the assumption that the concentration of undissociated HAc will not be appreciably reduced by the dissociation reaction. In this case, if $x = (H^+) = (Ac^-)$, the concentration, $(HAc) = (0.1 - x)$.

$$1.8 \times 10^{-5} = (x)(x)/(0.1 - x) \quad 12-8$$

This is a quadratic equation. When arranged in the standard quadratic form, $ax^2 + bx + c = 0$, it appears as follows.

$$x^2 + (1.8 \times 10^{-5})x - 1.8 \times 10^{-6} = 0 \quad 12-9$$

$$x = \{-(1.8 \times 10^{-5}) + [(1.8 \times 10^{-5})^2 - 4(1)(-1.8 \times 10^{-6})]^{1/2}\}/2 \quad 12-10$$

$$(H^+) = 0.00133 : pH = 2.87 \quad 12-11$$

The obvious conclusion is that the simplifying assumption is justified.

Step-by-Step Solutions to Problems in this Chapter, "Calculations Using pH and the Dissociation Constant"

• PROBLEM 410

Find K_b and pK_b for the acetate ion CH_3COO^- . The ionization constant of CH_3COOH is $K_a = 1.75 \times 10^{-5}$; $K_w = 1.00 \times 10^{-14}$.

Solution: This problem is concerned with conjugate acid-base pairs. In the general reaction,



HA is the conjugate acid of A^- , or conversely, A^- is the conjugate base of HA. H_2O has acted as a weak base. The acid and base ionization constants for a conjugate pair, as illustrated above, are K_a and K_b .

$$K_a K_b = K_w \text{ which means } pK_a + pK_b = pK_w.$$

Thus, if given the ionization constant for an acid or base, one can calculate the ionization constant for its conjugate base or acid. Also, remember that the K and pK values are related by the general equation,

$$pK = -\log K.$$

Therefore, the values of K_b and pK_b are

$$K_b = \frac{K_w}{K_a} = \frac{1.00 \times 10^{-14}}{1.75 \times 10^{-5}} = 5.71 \times 10^{-10}.$$




$$\begin{aligned} pK_b &= -\log K_b = -\log 5.71 \times 10^{-10} \\ &= -[-10 + \log 5.71] = 9.244. \end{aligned}$$

A chemist mixes .5 moles of acetic acid ($\text{HC}_2\text{H}_3\text{O}_2$) and .5 moles of HCN with enough water to make a one liter solution. Calculate the final concentrations of H_3O^+ and OH^- . Assume the following constants:


$$K_{\text{HC}_2\text{H}_3\text{O}_2} = 1.8 \times 10^{-5}, \quad K_{\text{HCN}} = 4 \times 10^{-10}, \quad K_w = 1.0 \times 10^{-14}.$$

Solution: To solve this problem, note the simultaneous equilibria. Once this is recognized, you can set up equilibrium constant expressions which measure these equilibria in terms of the concentrations of the species involved.

There are three simultaneous equilibria in the final solution. Two derive from the acids HCN and $\text{HC}_2\text{H}_3\text{O}_2$ donating their protons to form hydronium ions in water. The third stems from the ionization of water to hydronium and hydroxyl ions. You have:



Each reaction contributes H_3O^+ ions. However, only the acetic acid produces an appreciable concentration of H_3O^+ . This can be determined by inspection of the dissociation constants. The larger a dissociation constant, the greater the dissociation of the species. The dissociation constant of acetic acid, 1.8×10^{-5} , is the largest, and, therefore, its $[\text{H}_3\text{O}^+]$ contribution is the greatest. As such, you can neglect the H_3O^+ contribution from HCN and water. Let the H_3O^+ contribution from acetic acid = x . If you started with a concentration of .5M for acetic acid, then, at equilibrium, it becomes $.5 - x$. From the dissociation of acetic acid in water, it also becomes evident that the concentration of $\text{C}_2\text{H}_3\text{O}_2^-$ can be represented by x , since the equation states H_3O^+ and $\text{C}_2\text{H}_3\text{O}_2^-$ will be produced in equimolar quantities. You have



thus,

$$K_{\text{HOAc}} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]} = \frac{x \cdot x}{.5 - x} = 1.8 \times 10^{-5}$$


Solving for x , using the quadratic equation, you obtain

$$x = 3.0 \times 10^{-3} \text{M} = [\text{H}_3\text{O}^+]$$

To find $[\text{OH}^-]$, use the fact that $K_w = [\text{OH}^-][\text{H}_3\text{O}^+] = 1 \times 10^{-14}$.


Since you know $[\text{H}_3\text{O}^+]$, you can substitute to find $[\text{OH}^-]$. Thus, $[\text{OH}^-] = 3.3 \times 10^{-12} \text{M}$.

Pentobarbital, sometimes prescribed for the treatment of chronic insomnia under the trade name Nembutal, has the following structure:




pentobarbital

It is toxic in doses of 0.5 g and lethal in doses of 1.5 g. Barbituric acid, denoted by H Bar, and which may be used in the synthesis of pentobarbital, has the following structure:



barbituric acid

H Bar dissociates according to the following equation:



It is a weak acid, having a pK_a of only 4.01. Approximately what percent of H Bar molecules are dissociated in a 0.10 N solution?

Solution: We will solve this problem by determining the concentrations of H^+ and Bar^- formed from the dissociation of H Bar.

If we let x denote the equilibrium concentration of H^+ , then, since one mole of Bar^- is formed per mole of H^+ formed, the equilibrium concentration of Bar^- is also x . Furthermore, the equilibrium concentration of H Bar is $0.10 - x$. Then

$$K_a = \frac{[H^+][\text{Bar}^-]}{[\text{H Bar}]} = \frac{x^2}{0.10 - x}$$

Note that we have neglected the contribution to $[H^+]$ from the dissociation of water. Assuming that x is much smaller than 0.10 (a justifiable assumption, since H Bar is so weak an acid), the $0.10 - x \approx 0.10$ and we have

$$K_a = \frac{x^2}{0.10 - x} \approx \frac{x^2}{0.10}$$

$$\text{or, } x^2 = 0.10 \times K_a$$

$$x = (0.10 \times K_a)^{1/2}$$

The value of K_a is determined from the pK_a . By definition, $pK_a = -\log K_a$. Hence $K_a = 10^{-pK_a} = 10^{-4.01}$, and

$$\begin{aligned} x &= (0.10 \times K_a)^{1/2} = (0.10 \times 10^{-4.01})^{1/2} = (10^{-5.01})^{1/2} \\ &= 10^{-2.50} \\ &\approx 3.2 \times 10^{-3} \end{aligned}$$

The equilibrium concentrations of H^+ , Bar^- , and H Bar are then $[H^+] = x = 3.2 \times 10^{-3} \text{ M}$, $[\text{Bar}^-] = x = 3.2 \times 10^{-3} \text{ M}$, and $[\text{H Bar}] = 0.10 - x = 0.10 - 3.2 \times 10^{-3} \approx 0.10 \text{ M}$.

The percent dissociation of H Bar is given by

$$\begin{aligned} \% \text{ dissociation} &= \frac{[H^+]}{[\text{H Bar}]} \times 100 \% = \frac{[\text{Bar}^-]}{[\text{H Bar}]} \times 100 \% \\ &= \frac{3.2 \times 10^{-3} \text{ M}}{0.10 \text{ M}} \times 100 \% \\ &\approx 3 \% \end{aligned}$$

• PROBLEM 413

Assuming $K_w = 1.0 \times 10^{-14}$ and $K_a = 4.0 \times 10^{-10}$, find the pH of a 1.0 M NaCN solution.

Solution: NaCN is a salt that exists in water as Na^+ and CN^- . As such, this problem requires considering the hydrolysis of a salt of a weak acid and strong base. Since water is a weak acid or weak base (which dissociates to H^+ and OH^- ions), it reacts with ions from either weak acids or bases.



Inherent in this equation is the ionization of water, that is, $\text{H}_2\text{O} \rightleftharpoons \text{HO}^- + \text{H}^+$ so that it is actually, $\text{CN}^- + \text{OH}^- + \text{H}^+ \rightleftharpoons \text{HCN} + \text{OH}^-$.

The hydrolysis of the NaCN salt can be expressed by K_{hyd} , a hydrolysis constant. $K_{\text{hyd}} = K_w/K_a$, where K_a is the ionization of the acid formed and K_w is the ionization constant of water to $\text{OH}^- + \text{H}^+$, as previously mentioned.

$$K_{\text{hyd}} = \frac{K_w}{K_a} = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]} = \frac{1.0 \times 10^{-14}}{4.0 \times 10^{-10}}.$$

It is given that $K_a = 4.0 \times 10^{-10}$. K_w is always 1.0×10^{-14} , since a pure water solution must be neutral. As such, the molar concentrations of H^+ and OH^- are both 1.0×10^{-7} moles/liter. This stems from the fact that $[\text{H}^+] = [\text{OH}^-]$ with $[\text{H}^+][\text{OH}^-] = K_w = 1.0 \times 10^{-14}$. Therefore,

$$[\text{H}^+]^2 = 1.0 \times 10^{-14} \quad \text{or} \quad [\text{H}^+] = 1.0 \times 10^{-7}.$$

One now has:

$$\frac{1.0 \times 10^{-14}}{4.0 \times 10^{-10}} = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]}$$

The object is to calculate $[\text{OH}^-]$. From $[\text{OH}^-]$, one can determine pOH, since $\text{pOH} = -\log [\text{OH}^-]$. Once one knows pOH, one can find pH, since $14 = \text{pH} + \text{pOH}$. To find $[\text{OH}^-]$, let the concentration of OH^- be x . If this is the case, then $[\text{HCN}]$ can also be represented by x . Recall, that $\text{CN}^- + \text{H}_2\text{O} \rightleftharpoons \text{HCN} + \text{OH}^-$. For every mole of OH^- generated, a mole of HCN must also be produced. The concentration of $[\text{CN}^-]$ can be represented as $1 - x$. It is given that the original concentration of NaCN or CN^- , is 1.0 M. When placed in water, some of CN^- becomes HCN. If HCN is x , then the original concentration of CN^- must be reduced by x . Therefore, $[\text{CN}^-]$ has a concentration of $1 - x$. It follows that

$$\frac{1.0 \times 10^{-14}}{4.0 \times 10^{-10}} = \frac{x \cdot x}{1 - x}, \text{ if one substitutes the values}$$

$$\text{in the equation } \frac{1.0 \times 10^{-14}}{4.0 \times 10^{-10}} = \frac{[\text{HCN}][\text{OH}^-]}{[\text{CN}^-]}.$$

The procedure now is to solve for x .

$$x = 5.0 \times 10^{-3} \text{ M} = [\text{OH}^-].$$

As previously stated, $\text{pOH} = -\log [\text{OH}^-]$.

$$\text{pOH} = 2.30 \quad \text{and} \quad \text{pH} = 14 - \text{pOH} = 11.70.$$

A laboratory technician has a solution of 0.1 M $\text{NaC}_2\text{H}_3\text{O}_2$. Find its pH and the percent hydrolysis. Assume $K_w = 1.0 \times 10^{-14}$ and $K_{\text{diss}} = 1.8 \times 10^{-5}$.

Solution: $\text{NaC}_2\text{H}_3\text{O}_2$ is a salt that exists as ions Na^+ and $\text{C}_2\text{H}_3\text{O}_2^-$ in water. Since water is a weak acid and weak base, which dissociates to H^+ and OH^- ions, it can react with ions from weak acids and bases. Such a process, termed hydrolysis, occurs in this problem. Because the problem asks for pH, one must determine the concentrations of the H_3O^+ or OH^- in this problem. This means that the hydrolysis reaction must be written and an equilibrium constant set up. The net hydrolysis reaction may be written as



The K_{hyd} for this reaction is

$$\frac{[\text{OH}^-][\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]}$$

However, K_{hyd} is defined as K_w/K_{diss} , where K_w is the dissociation constant for water and K_{diss} is the dissociation constant for the acid formed. One can equate the two to obtain

$$\frac{[\text{OH}^-][\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]} = \frac{K_w}{K_{\text{diss}}}.$$

But the value of K_w and K_{diss} is known. Therefore,

$$\frac{[\text{OH}^-][\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]} = \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-5}} = 5.6 \times 10^{-10}.$$

If one can find $[\text{OH}^-]$, one can determine $[\text{H}_3\text{O}^+]$ from $[\text{H}_3\text{O}^+][\text{OH}^-] = K_w = 1.0 \times 10^{-14}$. Therefore, let $x =$ moles/liter of OH^- . From the hydrolysis reaction, one can see that for each mole of OH^- generated, a mole of $\text{HC}_2\text{H}_3\text{O}_2$ is also generated. Therefore, let $x = [\text{HC}_2\text{H}_3\text{O}_2]$ also. If one starts with 0.1 M of $\text{C}_2\text{H}_3\text{O}_2^-$, and x moles/liter reacts to form OH^- (and $\text{C}_2\text{H}_3\text{O}_2^-$), then, at equilibrium, one has $0.1 - x$ moles/liter of $\text{C}_2\text{H}_3\text{O}_2^-$ left. Substituting these values into the previous expression:

$$\frac{x \cdot x}{0.1 - x} = 5.6 \times 10^{-6}. \text{ Solving for } x,$$

$$x = 7.5 \times 10^{-4} \text{ M.}$$

(assuming in this calculation that x is small, therefore approximating $0.1 - x$ as 0.1.) To find $[\text{H}^+]$, solve

$$[\text{H}_3\text{O}^+] = \frac{K_w}{[\text{OH}^-]} = \frac{1.0 \times 10^{-14}}{7.5 \times 10^{-4}} = 1.33 \times 10^{-11}.$$

$$\text{Since } \text{pH} = -\log [\text{H}_3\text{O}^+]$$

$$\text{pH} = -\log [1.33 \times 10^{-11}] = 10.88.$$

The percent hydrolysis of this reaction is the number of moles of hydrolyzed acetate ion, divided by the number of moles available for hydrolysis, times 100.

$$\begin{aligned} \% \text{ Hydrolysis} &= \frac{\text{Moles } \text{C}_2\text{H}_3\text{O}_2^- \text{ hydrolyzed}}{\text{Moles } \text{C}_2\text{H}_3\text{O}_2^- \text{ available}} \times 100 \\ &= \frac{7.5 \times 10^{-4}}{0.1} \times 100 = 0.75\%. \end{aligned}$$


• PROBLEM 415

What is the pH of a 0.001 M aqueous solution of the strong electrolyte KOH? The dissociation constant for water is $K_w = 10^{-14}$ mole²/liter².

Solution: We need to find $[\text{H}_3\text{O}^+]$ in order to find the pH. $[\text{H}_3\text{O}^+]$ can be determined by substituting into the expression $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$, or,

$$[\text{H}_3\text{O}^+] = \frac{K_w}{[\text{OH}^-]}$$

The hydroxyl concentration, $[\text{OH}^-]$, is contributed by the dissociation of KOH. This equation for this reaction is



Since KOH is a strong electrolyte, we assume that it dissociates completely. Hence, the concentrations of OH^- and of K^+ are equal to the initial concentration of KOH, or $[\text{OH}^-] = 0.001 \text{ M} = 10^{-3} \text{ M}$. The hydronium ion concentration is then

$$\begin{aligned} [\text{H}_3\text{O}^+] &= \frac{K_w}{[\text{OH}^-]} = \frac{10^{-14} \text{ mole}^2/\text{liter}^2}{10^{-3} \text{ M}} \\ &= \frac{10^{-14} \text{ mole}^2/\text{liter}^2}{10^{-3} \text{ mole/liter}} = 10^{-11} \text{ mole/liter} = 10^{-11} \text{ M}. \end{aligned}$$

The pH is defined as $\text{pH} = -\log [\text{H}_3\text{O}^+]$. Hence

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log (10^{-11}) = -(-11) = 11.$$

Find the pH of a 0.2 M solution of formic acid. $K_a = 1.76 \times 10^{-4}$.

Solution: To solve the problem, first note that the acid dissociates according to the general equation



where HA is the acid and H^+ and A^- are the dissociation products. To find the pH, calculate $[\text{H}^+]$.

One can see from the equation that equal concentrations of H^+ and A^- form when HA dissociates. At equilibrium, x moles of HA dissociate into x moles of H^+ and x moles of A^- .



Before: 0.2 0 0

After: 0.2-x x x

Now, set up the equilibrium constant equation, which is

$$K_a = \frac{[\text{products}]}{[\text{reactants}]} = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} = 1.76 \times 10^{-4}.$$

Before substituting in the values, simplify the calculation by assuming that $0.2 - x \approx 0.2$. This is a valid assumption since the acid is very weak and only a very small percentage dissociates. Thus, one has

$$K_a = \frac{(x)(x)}{(0.2)} = 1.76 \times 10^{-4}. \quad \text{Solving:}$$

$$x = [\text{H}^+] = 5.93 \times 10^{-3}.$$

H^+ concentration is then converted to pH according to the equation,

$$\begin{aligned} \text{pH} &= -\log [\text{H}^+] = -\log 5.93 \times 10^{-3} \\ &= -[-3 + \log 5.93] = 3 - 0.77 = 2.23. \end{aligned}$$

Lemon juice is very acidic, having a pH of 2.1. If we assume that the only acid in lemon juice is citric acid (represented by HCit), that HCit is monoprotic and that no citrate salts are present, what is the concentration of citric acid in lemon juice? The dissociation constant

for citric acid



is $K_a = 8.4 \times 10^{-4}$ mole/liter.

Solution: The solution to this problem involves determination of $[\text{HCit}]$ from the expression for K_a of the reaction



By definition,

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{Cit}^-]}{[\text{HCit}]}, \text{ or, } [\text{HCit}] = \frac{[\text{H}_3\text{O}^+][\text{Cit}^-]}{K_a}$$

Since one mole of Cit^- is produced per mole of H_3O^+ produced, $[\text{Cit}^-] = [\text{H}_3\text{O}^+]$. $[\text{H}_3\text{O}^+]$ can be determined from the pH. By definition, $\text{pH} = -\log [\text{H}_3\text{O}^+]$, hence

$$[\text{H}_3\text{O}^+] = 10^{-\text{pH}} = 10^{-2.1} \text{ mole/liter. Therefore,}$$


$$[\text{H}_3\text{O}^+] = [\text{Cit}^-] = 10^{-2.1} \text{ mole/liter and we have}$$

$$\begin{aligned} [\text{HCit}] &= \frac{[\text{H}_3\text{O}^+][\text{Cit}^-]}{K_a} \\ &= \frac{10^{-2.1} \text{ mole/liter} \times 10^{-2.1} \text{ mole/liter}}{8.4 \times 10^{-4} \text{ mole/liter}} \\ &= \frac{(10^{-2.1})^2}{8.4 \times 10^{-4}} \text{ mole/liter} = \frac{10^{-4.2}}{8.4 \times 10^{-4}} \text{ mole/liter} \\ &\approx \frac{6.3 \times 10^{-5}}{8.4 \times 10^{-4}} \text{ mole/liter} \\ &= 7.5 \times 10^{-2} \text{ mole/liter} = 7.5 \times 10^{-2} \text{ M.} \end{aligned}$$

• PROBLEM 418

Find the pH of a solution of 0.3 M acetic acid. Assume a K_a value of 1.8×10^{-5} .

Solution: The dissociation of acetic acid into positive and negative ions proceeds as follows:



For this particular dissociation the equilibrium

constant, K_a , is written as

$$K_a = \frac{[H^+][C_2H_3O_2^-]}{[HC_2H_3O_2]} = 1.8 \times 10^{-5},$$

where the brackets represent the concentrations of the substances within them.

The pH of a solution is defined as the negative log of the hydrogen ion concentration. Mathematically, it is written as

$$pH = -\log [H^+].$$

Thus, to answer this problem, the $[H^+]$ in this solution must be calculated from the equilibrium constant expression and then substituted into the pH equation.

From the stoichiometry of the dissociation reaction, for every x molar concentration of H^+ formed there are x molar concentration of $C_2H_3O_2^-$ also formed. Therefore, let

$$x = [H^+] = [C_2H_3O_2^-].$$

The x molar concentration of these ions comes from the dissociation of the 0.3 M acetic acid. Thus, the exact amount of acetic acid in solution at equilibrium is $0.3 - x$. However, x is a very small amount as compared with 0.3 M of the acetic acid and is, therefore, approximated as 0.3 M. Substituting these values into the equilibrium expression

$$1.8 \times 10^{-5} = \frac{x \cdot x}{0.3 - x} \approx \frac{x^2}{0.3}$$

$$\text{Solving for } x; \quad x = [H^+] = 2.3 \times 10^{-3}$$


Substituting this value of $[H^+]$ into the pH equation:

$$pH = -\log (2.3 \times 10^{-3}) = 2.63.$$


• PROBLEM 419

For HF, $K_{diss} = 6.7 \times 10^{-4}$, what is the H_3O^+ and OH^- concentrations of a 0.10M HF solution? Assume $K_w = 1 \times 10^{-14}$.

Solution: To answer this question, you must set up the equations for the equilibria involved. HF is an acid, which means it donates protons (H^+). The only base present is H_2O . They react as shown in the following equation:



This is not the only source of H_3O^+ . Recall that water also can dissociate into ions.



This means that in the consideration of the H_3O^+ concentration you must look at both sources. Because you are asked the concentrations, you must employ equilibrium constants, K_{diss} and K_w . Let x = the number of moles/liter of H_3O^+ from the reaction of HF with H_2O , then x is also equal to the number of moles/liter of F^- produced. They have the same mole ratio, as indicated by the equilibria. Since HF is the stronger acid, some of its protons ionize and then join with H_2O to form H_3O^+ . Thus, the concentration of HF is $(0.1 - x)$. If you let y = moles/liter of OH^- , then y = moles/liter of H_3O^+ formed by the dissociation of water. These variables that represent the concentrations at equilibrium must satisfy the two equilibrium conditions:

$$K_{\text{diss}} = \frac{[\text{H}_3\text{O}^+][\text{F}^-]}{[\text{HF}]} = 6.7 \times 10^{-4} = \frac{(x+y)(x)}{0.1-x}$$

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} = (x+y)(y).$$

For both conditions, the H_3O^+ concentration is represented by $(x+y)$ since, at equilibrium, the concentration must be the sum from both sources of H_3O^+ production. Solve these equations simultaneously for x and y , you can avoid cumbersome calculations by noting which contribution is the dominant one. In this way, the other can be neglected to give a fairly good approximation. A general rule for determining which is dominant is to compare the dissociation constants. The larger the dissociation constant, the more it dominates the final equilibrium state. From inspection of these constants, you find that the contribution of H_3O^+ from HF dominates over that from water. You need consider only $\text{HF} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{F}^-$, therefore,

$$K_{\text{diss}} = \frac{[\text{H}_3\text{O}^+][\text{F}^-]}{[\text{HF}]} = 6.7 \times 10^{-4} = \frac{(x+y)(x)}{0.1-x}.$$

Solving, you obtain $x = 7.7 \times 10^{-3} \text{ M} = [\text{H}_3\text{O}^+]$. To find y , go back to $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} = (x+y)(y)$ and substitute in your value for x . Assuming that y is negligible in comparison to x , the answer becomes

$$\frac{1.0 \times 10^{-14}}{7.7 \times 10^{-3}} = 1.3 \times 10^{-12} \text{ M} = [\text{OH}^-].$$

• PROBLEM 420

There exists the equilibrium $\text{CO}_2 + 2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HCO}_3^-$. The $K_a = 4.3 \times 10^{-7}$ and $\text{pH} = 7.40$. Find the ratio of concentrations of bicarbonate (HCO_3^-) to carbon dioxide (CO_2).

Solution: The equilibrium constant expression, which indicates the ratio of the products to reactants, each

raised to the power of their respective coefficients, can be written as follows:

$$K_a = \frac{[H_3O^+][HCO_3^-]}{[CO_2]}$$

Note, $[H_2O]$ is omitted, since it is considered to be constant. Since $K_a = 4.3 \times 10^{-7}$, then

$$4.3 \times 10^{-7} = \frac{[H_3O^+][HCO_3^-]}{[CO_2]}$$

One is solving for $[HCO_3^-]/[CO_2]$. If one knows $[H_3O^+]$, then one can calculate this ratio.

$[H_3O^+]$ can be found, since, as given, the $pH = 7.40$.

$$pH = -\log [H_3O^+] \quad \text{or}$$

$$\begin{aligned} [H_3O^+] &= \text{antilog} (-pH) = \text{antilog} (-7.40) \\ &= 4.0 \times 10^{-8} \text{ M.} \end{aligned}$$


$$\text{Thus, } 4.3 \times 10^{-7} = \frac{4.0 \times 10^{-8} [HCO_3^-]}{[CO_2]}, \quad \text{and}$$

$$\frac{[HCO_3^-]}{[CO_2]} = \frac{4.3 \times 10^{-7}}{4.0 \times 10^{-8}} = 10.75.$$

• PROBLEM 421

100 ml of 0.1 M NaOH is mixed with 100 ml of 0.15 M HOAC. $K_a = 1.75 \times 10^{-5}$ for acetic acid, what is the pH of this mixture?

Solution: To find the pH of the mixture, first determine the H_3O^+ concentration, since $pH = -\log [H_3O^+]$. To find H_3O^+ , consider what happens when acetic acid is added to NaOH, a base. When an acid is added to a base, a neutralization reaction occurs such that water and a salt are produced. Thus,



This reaction does not show the presence of any H_3O^+ ions, which may lead one to incorrectly conclude that its concentration was zero. There are 100 ml of 0.15 M HOAC. $M = \text{Molarity} = \text{moles/liter}$. Thus, $0.15 = \text{moles}/0.10 \text{ liters}$ ($1 \text{ liter} = 1000 \text{ ml}$). Therefore, $(0.15)(0.10) = 0.015 \text{ moles}$ of HOAC. For NaOH, $M = 0.10 = \text{moles}/0.10 \text{ liters}$. Thus,

the number of moles of NaOH = (0.10)(0.10) = 0.010. From the previously written neutralization equation, one mole of acid reacts for every mole of base. Only 0.01 moles of HOAC could react with NaOH. Thus, 0.015 - 0.010 = 0.005 moles of HOAC was NOT neutralized by the base, NaOH. This excess 0.005 moles of HOAC is, in the presence of its salt NaOAC. Thus, it is a buffer solution. Solutions that contain a weak acid, such as HOAC, and its salt are called buffers. Therefore, consider the dissociation of acetic acid; it is in excess and is the source of the hydronium ion concentration.



The equilibrium constant for this reaction measures the ratio of the concentration of the products to reactants, each raised to a power based on its coefficient in the chemical equation. Thus,

$$K_a, \text{ the equilibrium constant, } = \frac{[\text{H}_3\text{O}^+][\text{OAC}^-]}{[\text{HOAC}]}.$$

Water is not included since it is a constant. One is solving for $[\text{H}_3\text{O}^+]$ to find pH, let $x = [\text{H}_3\text{O}^+]$. From the dissociation equation $x = [\text{OAC}^-]$ also, since they are formed in equimolar amounts. Recall, however, that the neutralization reaction produced OAC^- also. There were 0.010 moles each of NaOH and HOAC which reacted. From the equation, 0.010 moles of OAC^- should have been produced. The volume was 200 ml. Thus,

$$M = \text{concentration} = \frac{0.01 \text{ moles}}{\frac{200 \text{ ml}}{1000 \text{ ml/l}}} = 0.05 \text{ M.}$$

The total concentration of OAC^- , from the acid dissociation and neutralization, is $x + .05$. The acetic acid concentration is the initial amount minus the amount that dissociated. Recall, there were 0.005 moles of acetic acid in 200 ml. to give

$$\frac{0.005 \text{ moles}}{\frac{200 \text{ ml}}{1000 \text{ ml/l}}} = 0.025 \text{ M.}$$

x moles/liter of it dissociated. Thus, the final concentration is $(0.025 - x)$.

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{OAC}^-]}{[\text{HOAC}]}.$$

$$1.75 \times 10^{-5} = \frac{[\text{H}_3\text{O}^+][\text{OAC}^-]}{[\text{HOAC}]}.$$

Substituting final concentrations,

$$1.75 \times 10^{-5} = \frac{(x)(x + 0.05)}{(0.025 - x)}.$$

The excess OAC^- , from the dissociation of HOAC , x , is small. Thus,

$$\frac{(x)(x + 0.05)}{(0.025 - x)} \approx \frac{(0.050 x)}{(0.025)} = 1.75 \times 10^{-5}.$$

Solving for x , one finds

$$[\text{H}_3\text{O}^+] = x = 8.8 \times 10^{-6} \text{ M}.$$

Thus, $\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log [8.8 \times 10^{-6}] = 5.06.$

• PROBLEM 422


Allow exactly 100 ml of 1.5 N NaOH solution to be mixed with 100 ml 3.0 N H_3PO_4 solution and allow them to reach equilibrium. (1) Determine what species will be present at equilibrium. (2) Find the pH of the solution. $K_2 = 6.2 \times 10^{-8}.$

Solution: To solve this problem, first consider the type of reaction, if any, that occurs and to what extent it has proceeded. Once this is done, the answers to (1) and (2) follow directly. One is adding a base of a given amount to an acid of a given amount. This is a neutralization; i.e., the reaction of an acid and base to produce a salt and water. One is told there are 100 ml of 1.5 N NaOH and 100 ml of 3.0 N H_3PO_4 solution. N = normality, which is defined as equivalents/liter. An equivalent = the number of hydronium ions that an acid can produce or hydroxide ions that a base can produce per mole of that substance. In a neutral solution the number of equivalents of base is equal to the number of equivalents of acid. Thus, to see to what extent the reaction has proceeded, calculate the equivalents of base and acid. If they are equal, the reaction went to completion. Recall, N = equivalents/liter. Thus, equivalents NaOH = $(1.5 \text{ equiv/l}) 100 \text{ ml} / 1000 \text{ m/l} = 0.15 \text{ equiv}.$ Equivalents $\text{H}_3\text{PO}_4 = (3.0 \text{ equiv/l}) 100 \text{ ml} / 1000 \text{ ml} = 0.30 \text{ equiv}.$


The equivalents of acid do NOT equal those of the base. The H_3PO_4 has $0.30 - 0.15 = 0.15$ equivalents that do not react. The rest has been neutralized by the base to form a salt and water. H_3PO_4 is a polyprotic acid, which means it can donate 3 protons when a mole of it exists. Since only one-half of the equivalents are neutralized, only one-half of the protons were used. This means that the reaction




goes to completion, but



will only be half complete. With this information, (1) can be answered. One started with 1.5 N of 100 ml NaOH solution, which indicates that 0.15 moles of Na^+ is present. NaOH can react with only 1 mole of hydronium ion. Thus, its equivalents correspond to its moles. 0.30 equivalents of H_3PO_4 correspond to 0.10 moles, since each mole generates 3 hydronium ions. In



there is a total 0.1 moles of H_2PO_4^- generated, since H_3PO_4 is the limiting reagent with only 0.1 moles compared with OH^- 's 0.15 moles. This means that for the reaction;



0.05 moles of OH^- is left to react with the 0.1 moles of

H_2PO_4^- generated previously. Thus, only 0.05 moles of H_2PO_4^- is consumed in producing 0.05 moles of HPO_4^{2-} . Thus, an unreacted amount of 0.05 moles of H_2PO_4^- is left. In summary, the species in solution are 0.05 moles each of H_2PO_4^- and HPO_4^{2-} with small amounts of H_3O^+ , H_3PO_4 , PO_4^{3-} , OH^- and 0.15 moles Na^+ . One can now calculate the pH of this. There exists some H_2PO_4^- in solution. This is the source of hydronium ions, whose pH is measured by $\text{pH} = -\log [\text{H}_3\text{O}^+]$. The reaction is



The equilibrium constant,

$$K_2 = \frac{[\text{H}_3\text{O}^+][\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} = 6.2 \times 10^{-8}$$

One can calculate $[\text{HPO}_4^{2-}]$ and $[\text{H}_2\text{PO}_4^-]$, since one calculated the mole amounts. The volume of the solution is 0.2 liters, and the concentration in the brackets, are in moles/liter. Substituting these values,

$$6.2 \times 10^{-8} = \frac{[\text{H}_3\text{O}^+](0.05/.2)}{(0.05/.2)}$$

Solving for $[\text{H}_3\text{O}^+]$, one obtains

$$[\text{H}_3\text{O}^+] = 6.2 \times 10^{-8} \text{ M. pH} = -\log [\text{H}_3\text{O}^+]. \text{ Thus,}$$


$$\text{pH} = -\log [6.2 \times 10^{-8}] = 7.21.$$

• PROBLEM 423

0.001 mol of NaOH is added to 100 ml. of a solution that is 0.5 M $\text{HC}_2\text{H}_3\text{O}_2$ and 0.5 M $\text{NaC}_2\text{H}_3\text{O}_2$. Determine the pH of this solution; the K_{diss} of acetic acid is 1.8×10^{-5} .

Solution: This problem requires an understanding of the concept of buffer solution. In some processes, a solution of constant pH is desired. This constancy is maintained by the buffering action of an acid-base equilibrium. A buffer contains both an acid and a base and responds to the addition of either H_3O^+ or OH^- to maintain the pH. In this problem, the 0.5 M solutions of $\text{HC}_2\text{H}_3\text{O}_2$ and $\text{NaC}_2\text{H}_3\text{O}_2$ act as a buffer that must respond to the addition of base, NaOH . One has equal numbers of moles of $\text{HC}_2\text{H}_3\text{O}_2$ and $\text{NaC}_2\text{H}_3\text{O}_2$ in the buffer, which means that the

$\frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]}$ ratio is one. In solution, one has



indicates that each mole/liter of $\text{HC}_2\text{H}_3\text{O}_2$ must produce the same number of mole/liter of $\text{C}_2\text{H}_3\text{O}_2^-$. For this reaction,

$$K = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_3\text{O}_2^-]}{[\text{HC}_2\text{H}_3\text{O}_2]} \quad \text{or} \quad [\text{H}_3\text{O}^+] = \frac{K [\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]},$$

which is an expression for the concentration of hydronium ions in the buffer. Recalling that $\text{pH} = -\log [\text{H}_3\text{O}^+]$ and that $\text{pK} = -\log K$, by taking the negative log of both sides of the above equation, one can write

$$-\log [\text{H}_3\text{O}^+] = -\log K - \log \frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]}.$$

$$\text{pH} = \text{pK} - \log \frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]}$$

For this solution, it is given that $K = 1.8 \times 10^{-5}$ and that $[\text{HC}_2\text{H}_3\text{O}_2]/[\text{C}_2\text{H}_3\text{O}_2^-]$ is equal to one. Thus,

$$\text{pH} = \text{pK} - \log \frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]}$$

$$\text{pH} = (-\log 1.8 \times 10^{-5}) - \log (1)$$

$$\text{pH} = 4.74 - 0 = 4.74$$

One adds 0.001 moles of NaOH . This base will convert an equal number of moles of the acid, $\text{HC}_2\text{H}_3\text{O}_2$ to $\text{C}_2\text{H}_3\text{O}_2^-$, by a neutralization reaction. If one started with 0.5 M $\text{HC}_2\text{H}_3\text{O}_2$ in 100 ml, one had 0.05 mole of it. Upon addition of NaOH , however, 0.001 moles of it is converted to $\text{C}_2\text{H}_3\text{O}_2^-$. This means one has $0.05 - 0.001 = 0.049$ moles of $\text{HC}_2\text{H}_3\text{O}_2$ left. There were 0.05 moles of $\text{NaC}_2\text{H}_3\text{O}_2$ to start. $\text{NaC}_2\text{H}_3\text{O}_2$ exists as the ions Na^+ and $\text{C}_2\text{H}_3\text{O}_2^-$. There were 0.001 moles of $\text{C}_2\text{H}_3\text{O}_2^-$ produced upon addition of NaOH , which means one has a total of $0.001 + 0.05 = 0.051$ moles of $\text{C}_2\text{H}_3\text{O}_2^-$. The volume of the solution remained at 100 ml. Therefore, these mole amounts are 0.49 M concentration of $\text{HC}_2\text{H}_3\text{O}_2$ and 0.51 M concentration of $\text{C}_2\text{H}_3\text{O}_2^-$. To find the new pH, one need only substitute these values into

$$\text{pH} = \text{pK} - \log \frac{[\text{HC}_2\text{H}_3\text{O}_2]}{[\text{C}_2\text{H}_3\text{O}_2^-]} .$$

$$\text{pH} = -\log (1.8 \times 10^{-5}) - \log \frac{.49}{.51} = 4.74 + 0.017 = 4.76.$$


• PROBLEM 424

Find the pH of a 0.1 M solution of ammonia, NH_3 . $\text{pK}_b = 1.76 \times 10^{-5}$.

Solution: This problem involves the dissociation of a weak base, which proceeds by the following general equation:



where B is the base and BH^+ and OH^- are the dissociation products. x moles of B dissociate into x moles of BH^+ and x moles of OH^- . This is more clearly seen below:



Before: 0.1 M 0 0

After: 0.1 - x x x

If x moles of NH_3 dissociate, then, at equilibrium, one has 0.1 - x left, since 0.1 is the initial concentration. NH_3 is a weak base and very little dissociation occurs, which means x is extremely small. This allows one to approximate 0.1 - x as 0.1. By now, it should be evident that $[\text{H}^+]$ does not appear in the equilibrium equation. The strategy will be to first find $[\text{OH}^-]$, then calculating the pOH, and then using the equation $\text{pK} = \text{pOH} + \text{pH} = 14$ to find pH.

In order to calculate the value of $[\text{OH}^-]$, set up the equilibrium constant equation for which $K_b = 1.76 \times 10^{-5}$.

Thus, $K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]}$. Substituting,

$$K_b = \frac{(x)(x)}{0.1} = 1.76 \times 10^{-5}, \text{ so that}$$

$$x = [\text{OH}^-] = 1.33 \times 10^{-3}.$$

Then, one can calculate pOH by:

$$\begin{aligned} \text{pOH} &= -\log [\text{OH}^-] = -\log [1.33 \times 10^{-3}] \\ &= 3 - \log 1.33 = 2.88. \end{aligned}$$

Therefore, $\text{pH} = 14 - \text{pOH}$

$$= 14 - 2.88 = 11.12.$$

• PROBLEM 425

Find the pH of 0.15 M H_2SO_4 solution, assuming $K_2 = 1.26 \times 10^{-2}$.

Solution: Begin this problem by noting that $\text{pH} = -\log [\text{H}^+]$. Thus, to find pH, calculate the $[\text{H}^+]$ in 0.15 M H_2SO_4 . H_2SO_4 is an acid and dissociates into H^+ ($= \text{H}_3\text{O}^+$) ions. The general reaction for acid dissociation can be written as



H_2SO_4 undergoes this reaction, as



But note, HSO_4^- is also an acid, it still possesses a hydrogen that can dissociate. Thus, one also has



In other words, there exist two dissociation reactions. As given, the dissociation constant for the second reaction, K_2 , measures the ratio of the concentrations of products to reactants. It is from this expression that $[\text{H}^+]$ can be determined. H_2SO_4 is a very strong acid in its first dissociation reaction. This means it is completely ionized; i.e. 100% of the H^+ comes off (the first H^+ only). Thus, if there are 0.15 M of H_2SO_4 , then, there are also 0.15 = $[\text{H}^+]$ for the first dissociation.

$[\text{H}^+] = [\text{HSO}_4^-]$, as can be seen from the first dissociation reaction and are formed in equimolar amounts. Also, $[\text{HSO}_4^-] = 0.15$ (initially). For the second ionization, one is given $K_2 = 1.26 \times 10^{-2}$, indicating dissociation is not complete. From the prior explanation of what K_2 indicates, one can write

$$K_2 = 1.26 \times 10^{-2} = \frac{[\text{H}_3\text{O}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]}.$$

Note: Water is not included since it is assumed to be a constant. From the first dissociation, it is known that $[\text{H}_3\text{O}^+] = 0.15$. If x moles/liter of H^+ is produced in the second dissociation, the total $[\text{H}_3\text{O}^+] = 0.15 + x$. $[\text{SO}_4^{2-}] = [\text{H}_3\text{O}^+]$ from the second dissociation, since here they are formed in equimolar amounts. Thus, $[\text{SO}_4^{2-}] = x$. If one started with $[\text{HSO}_4^-] = 0.15\text{M}$ (from the first dissociation) and x moles/liter dissociate in the second, one has $[\text{HSO}_4^-] = 0.15 - x$ left.

Substituting these values,

$$1.26 \times 10^{-2} = \frac{[\text{H}_3\text{O}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} = \frac{(0.15 + x)(x)}{(0.15 - x)}.$$

Solving for x (using the quadratic formula), $x = .011$. Thus, $[\text{H}^+] = 0.15 + 0.011 = 0.161$. $\text{pH} = -\log [\text{H}^+]$, so that $\text{pH} = -\log [0.161] = 0.79$.

• PROBLEM 426

Find the pH of 0.10 M HOAc solution that has 0.20 M NaOAc dissolved in it. The dissociation constant of HOAc is 1.75×10^{-5} .

Solution: To find the pH of this solution, calculate the H_3O^+ concentration; pH is defined as $-\log [\text{H}_3\text{O}^+]$. This necessitates considering the dissociation of acetic acid, HOAc, since the only source of H_3O^+ is from this dissociation. The dissociation reaction may be written



To find the H_3O^+ concentration, the equilibrium constant expression must be written. This expression measures the ratio of the concentrations of products to reactants, each raised to the power of their respective coefficients in the chemical equation. Thus,

$$K_a = \frac{[\text{OAc}^-][\text{H}_3\text{O}^+]}{[\text{HOAc}]}.$$

(Note: water is excluded since it is considered a constant.)

Let $x = [\text{H}_3\text{O}^+]$ at equilibrium. The $[\text{OAc}^-]$ at equilibrium will be from two sources: the dissociation of HOAc and the presence of NaOAc, which exists as ions Na^+ and OAc^- . From the dissociation, the concentration is x , since the reaction indicates that H_3O^+ and OAc^- are formed in equimolar amounts. Since one starts with 0.2 M NaOAc, one has a concentration of 0.2 M for OAc^- . The total is $x + 0.2$. If one starts with 0.1 M HOAc and x M of H_3O^+ or OAc^- form, then, at equilibrium, one has $0.1 - x$ left of HOAc. Substituting these values into the equilibrium constant expression, and noting that $K_a = 1.75 \times 10^{-5}$ (given),

$$K_a = \frac{x(0.2 + x)}{(0.1 - x)} \approx \frac{0.20x}{0.1} = 1.7 \times 10^{-5}.$$


The acetic acid is weak; its dissociation to H_3O^+ is low, thus, one can make the above approximations. Solving for x ,

$$x = 8.5 \times 10^{-6} \text{ M} = [\text{H}_3\text{O}^+].$$

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log [8.5 \times 10^{-6}] = 5.07.$$

Assuming $pD = -\log [D_3O^+]$ in analogy to pH, what is pD of pure D_2O ? $K = 2 \times 10^{-15}$.

Solution: This question can be solved with the knowledge that D_2O undergoes dissociation in a similar manner to H_2O . The only difference is that D_2O is composed of deuterium instead of hydrogen. Therefore, one can write:



$$K = [D_3O^+][OD^-] = 2 \times 10^{-15}.$$


$[D_2O]$ is a constant and does not appear in the above equilibrium expression. From the reaction, it becomes apparent that the concentrations of D_3O^+ and OD^- are equal. Therefore, both concentrations can be represented by x , $x \cdot x = 2 \times 10^{-15}$. Solving for x ,

$$x = 4.47 \times 10^{-8}.$$

This, then, represents the concentration of D_3O^+ (and OD^-).

$$pD = -\log [D_3O^+] = -\log 4.47 \times 10^{-8} = 7.35.$$

Calculate the pH of a 0.2 M NH_3 solution for which $K_b = 1.8 \times 10^{-5}$ at 25°C. The equation for the reaction is



Solution: pH is defined in terms of $[H^+]$, ($pH = -\log [H^+]$), but this reaction shows the production of OH^- . However, $[H^+]$ and $[OH^-]$ are related by the definition, $[H^+] = 10^{-14}/[OH^-]$. Therefore, one can find $[H^+]$ and pH after solving for $[OH^-]$. One solves for $[OH^-]$ by using the fact that K_b is equal to $\frac{[NH_4^+][OH^-]}{[NH_3]}$

$[H_2O]$ is excluded from this expression because it is assumed to be constant.

In solving for $[OH^-]$, one can assume that 1 l of the 0.2 M NH_3 solution is present. This means that there were originally 0.2 moles of NH_3 present. For each NH_3 that dissociates, 1 mole of OH^- and 1 mole of NH_4^+ is formed. Let $x = [OH^-] = [NH_4^+]$ and $0.2 - x = [NH_3]$

$$K_b = \frac{[NH_4^+][OH^-]}{[NH_3]} = 1.8 \times 10^{-5} = \frac{(x)(x)}{(0.2 - x)}$$

One can assume that x is negligible compared to 0.2. Therefore, $0.2 - x = 0.2$. Using this assumption, one can now solve for x .

$$1.8 \times 10^{-5} = \frac{x^2}{0.2}$$

$$x^2 = 3.6 \times 10^{-6}; \quad x = 1.9 \times 10^{-3}.$$

$$[\text{NH}_4^+] = 1.9 \times 10^{-3} \text{ M}, \quad [\text{OH}^-] = 1.9 \times 10^{-3} \text{ M} \quad \text{and}$$

$$[\text{NH}_3] = 0.2 - 1.9 \times 10^{-3} \approx 0.2.$$

One can now solve for $[\text{H}^+]$ using $[\text{OH}^-]$.

$$[\text{H}^+] = \frac{10^{-14}}{[\text{OH}^-]} = \frac{10^{-14}}{1.9 \times 10^{-3}} = 5.26 \times 10^{-12} \text{ M}$$


Using this value one can solve for pH.

$$\begin{aligned} \text{pH} &= -\log [\text{H}^+] = -\log (5.26 \times 10^{-12}) \\ &= -(.72 - 12) = 11.28. \end{aligned}$$

• PROBLEM 429

What is the pH of 0.500 M NaHSO_3 ? The K_{eq} for NaHSO_3 is 1.26×10^{-2} .

Solution: The pH is defined as $-\log [\text{H}^+]$, where $[\text{H}^+]$ is the concentration of H^+ ions. NaHSO_3 ionizes as shown in the equation:



Thus, for each NaHSO_3 that ionizes, one H^+ and one HSO_3^- are formed. Therefore, $[\text{H}^+] = [\text{HSO}_3^-]$. The equilibrium constant (K_{eq}) for NaHSO_3 is 1.26×10^{-2} . The equation for K_{eq} is

$$K_{\text{eq}} = \frac{[\text{H}^+][\text{SO}_3^{2-}]}{[\text{NaHSO}_3]} = 1.26 \times 10^{-2}$$

One knows that $[\text{NaHSO}_3] = 0.500$ and that for each H^+ and NaSO_3^- formed, one NaHSO_3 is ionized. Therefore, at equilibrium, the concentration of NaHSO_3 is $.500 - [\text{H}^+]$. One can now solve for $[\text{H}^+]$, using the equation for the equilibrium constant.

$$K_{\text{eq}} = \frac{[\text{H}^+][\text{HSO}_3^-]}{[\text{NaHSO}_3]} = 1.26 \times 10^{-2}$$

and because $[H^+] = [SO_3^{2-}]$, you have

$$K_{eq} = \frac{[H^+][H^+]}{(0.500 - [H^+])} = 1.26 \times 10^{-2}$$

$$[H^+]^2 = (0.500 - [H^+]) \times 1.26 \times 10^{-2}$$

$$[H^+]^2 = (6.3 \times 10^{-3}) - 1.26 \times 10^{-2}[H^+]$$

$$\text{Rewriting, } [H^+]^2 + 1.26 \times 10^{-2}[H^+] - 6.3 \times 10^{-3} = 0.$$

Using the quadratic formula:

$$ax^2 + bx + c = 0$$

$$a = 1$$

$$b = 1.26 \times 10^{-2}$$

$$c = -6.3 \times 10^{-3}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-1.26 \times 10^{-2} \pm \sqrt{(1.26 \times 10^{-2})^2 - [4 \times 1 \times (-6.3 \times 10^{-3})]}}{2 \times 1}$$

$$x = \frac{-1.26 \times 10^{-2} \pm \sqrt{1.59 \times 10^{-4} + (2.52 \times 10^{-2})}}{2}$$

$$x = \frac{-1.26 \times 10^{-2}}{2} \pm \frac{\sqrt{2.54 \times 10^{-2}}}{2}$$

$$x = -6.3 \times 10^{-3} \pm \frac{1.59 \times 10^{-1}}{2}$$

$$x = -6.3 \times 10^{-3} \pm 7.97 \times 10^{-2}$$

$$x = -6.3 \times 10^{-3} + 7.97 \times 10^{-2} = 7.34 \times 10^{-2} \quad \text{or}$$

$$x = -6.3 \times 10^{-3} - 7.97 \times 10^{-2} = -8.60 \times 10^{-2}$$

x must equal 7.34×10^{-2} because $[H^+]$ cannot be negative. One can now find pH.

$$\text{pH} = -\log [H^+] \quad [H^+] = 0.734$$

$$\text{pH} = -\log 0.734$$

$$\text{pH} = 1.13.$$

• PROBLEM 430

At normal body temperature, 37°C (98.6°F), the ionization constant of water, K_w , is 2.42×10^{-14} moles²/liter².

A physician injects a neutral saline solution into a patient. What will be the pH of this solution when it has come into thermalequilibrium with the patient's body?

Solution: To solve this problem we must employ the definition of K_w , i.e. $K_w = [H_3O^+][OH^-]$. At 37°C, we are given

$$K_w = 2.42 \times 10^{-14} \text{ mole}^2/\text{liter}^2 = [H_3O^+][OH^-].$$

Since, for neutral solution, $[H_3O^+] = [OH^-]$, we can set $x = [H_3O^+] = [OH^-]$. Then,

$$x^2 = [H_3O^+][OH^-] = 2.42 \times 10^{-14} \text{ mole}^2/\text{liter}^2 \quad \text{or}$$

$$x = (2.42 \times 10^{-14} \text{ mole}^2/\text{liter}^2)^{1/2} = 1.56 \times 10^{-7} \text{ mole/liter}.$$

Hence, $[H_3O^+] = x = 1.56 \times 10^{-7} \text{ mole/liter}$ and the pH of the solution is

$$pH = -\log [H_3O^+] = -\log (1.56 \times 10^{-7})$$

$$= -(-6.807) = 6.807 \approx 6.8.$$

● PROBLEM 431

K_{diss} of water for $2H_2O \rightleftharpoons H_3O^+ + OH^-$ changes from 1.0×10^{-14} at 25°C to 9.62×10^{-14} at 60°C. Does the pH of water or its neutrality change when the temperature is increased from 25°C to 60°C?

Solution: To answer this question, it is necessary to find $[H^+]$, since $pH = -\log [H^+]$. To do this, write out the equilibrium dissociation expression for water, which equates the dissociation constant, K_{diss} , with the ratio of concentrations of products to reactants, each raised to the power of its coefficient in the chemical reaction. Thus, $K_{\text{diss}} = [H_3O^+][OH^-]$. The concentration of water is omitted; it is assumed to be constant. For the chemical reaction one can see that OH^- and H_3O^+ are formed in equimolar amounts, which means $[OH^-] = [H_3O^+]$. Thus, when $K_{\text{diss}} = 1.0 \times 10^{-14}$, one has

$$1.0 \times 10^{-14} = [H_3O^+][OH^-] = [H_3O^+]^2 = [OH^-]^2. \text{ Solving}$$


for either, $[H_3O^+] = [OH^-] = 1.0 \times 10^{-7}$. Thus, $pH = -\log [H_3O^+] = 7$. When $K_{\text{diss}} = 9.62 \times 10^{-14}$ at 60°C,

$$9.62 \times 10^{-14} = [H_3O^+][OH^-] = [H_3O^+]^2 = [OH^-]^2. \text{ Solving}$$

for either, $[H_3O^+] = [OH^-] = 3.1 \times 10^{-7}$. Thus, $pH = -\log [H_3O^+] = 6.51$. Thus, the pH does change. What about neutrality? Neutrality occurs when $[H^+] = [OH^-]$. In both cases, such a condition exists, so that the neutrality was not altered when the temperature was changed.

A solution of hydrogen peroxide is 30% by weight H_2O_2 . Assuming a density of 1.11 g/cm^3 and a dissociation constant of 1.0×10^{-12} for H_2O_2 , what is the pH of the solution?

Solution: To obtain the pH of any solution, calculate $[\text{H}^+]$, since $\text{pH} = -\log [\text{H}^+]$. To find $[\text{H}^+]$, write a balanced chemical equation that expresses the reaction, and from this, set up a dissociation constant expression. Begin by writing the reaction.



The dissociation constant for this reaction, K , equals 1.0×10^{-12} . K also = $[\text{H}^+][\text{HO}_2^-]/[\text{H}_2\text{O}_2]$. Equating, one obtains

$$\frac{[\text{H}^+][\text{HO}_2^-]}{[\text{H}_2\text{O}_2]} = 1.0 \times 10^{-12}$$

Solving this expression for $[\text{H}^+]$, one obtains the pH. To do that it is necessary to know $[\text{H}_2\text{O}_2]$. Given the density, one can assume a volume of 1 liter solution. This means that the mass of the solution is $1.11 \text{ g/cm}^3 \times 1000 \text{ ml (or 1 liter)} = 1110 \text{ g/l}$. However, the percent by weight of H_2O_2 is 30%. Therefore, there is only 0.30 1110 g/l of H_2O_2 or 333 g/l . The molecular weight of H_2O_2 is 34 g/mole . The molarity of the H_2O_2 solution is, then,

$$\frac{333 \text{ g/liter}}{34 \text{ g/mole}} = 9.79 \text{ M } \text{H}_2\text{O}_2$$

since molarity equals moles per liter. If one lets $x = [\text{H}^+]$, then $[\text{H}_2\text{O}_2] = 9.79 - x$, since, at equilibrium, the original amount of H_2O_2 must be decreased by the amount of H^+ (or O_2H^-) formed. The equation,



indicates that H^+ will be formed in equal mole amounts with HO_2^- . Thus, $[\text{HO}_2^-]$ can equal x . Substituting these values into the expression:

$$\frac{[\text{H}^+][\text{HO}_2^-]}{[\text{H}_2\text{O}_2]} = 1.0 \times 10^{-12}$$

$$\frac{x \cdot x}{9.79 - x} = 1.0 \times 10^{-12} \text{ . Solving for } x,$$

$$x = 3.13 \times 10^{-6} = [\text{H}^+]$$

Since $\text{pH} = -\log [\text{H}^+]$, $\text{pH} = 5.5$.

• PROBLEM 433

Calculate the pH at 25°C of a solution containing 0.10 M sodium acetate and 0.03 M acetic acid. The apparent pK for acetic acid at this ionic strength is 4.57.

Solution: This problem involves the calculation of $[H^+]$ ($pH = -\log [H^+]$) in a mixture of a monoprotic acid (acetic acid) and its completely dissociated salt (sodium acetate). Because the addition of a salt represses the ionization of the acid, the concentration of undissociated acid is approximately equal to the molar concentration of added weak acid (C_a). The addition of the acid represses the hydrolysis of the salt so that the concentration of anions of the weak acid is approximately equal to the molar concentration of the salt C_s .

Using several approximations, the pK for a weak acid is the pH of a solution containing equimolar quantities of salt and acid. However, if the solution does not contain equimolar quantities of salt and acid, then the following approximation holds:

$$pH = pK + \log \frac{C_s}{C_a} ,$$

where C_s and C_a are the concentrations of the salt and the acid, respectively.

Thus, in this problem,

$$C_s = 0.10 \text{ M}$$

$$C_a = 0.03 \text{ M}$$

$$pK = 4.57$$

$$\text{and } pH = 4.57 + \log \frac{0.10}{0.03} = 4.57 + 0.52 = 5.09.$$


• PROBLEM 434

10 ml of 0.200 M $HC_2H_3O_2$ is added gradually to 25 ml of 0.200 M NaOH. Calculate the pH of the initial 0.200 M NaOH solution and after each successive addition of 5 ml of acid. $K_w = 10^{-14}$.

Solution: This is a titration problem. One is asked to find the pH of the 0.200 M NaOH at different concentration levels. First one wants to calculate the pH before any acid is added. To do this, consider the fact that the equilibrium constant for the autodissociation of water, K_w , is defined as

$$K_w = [H_3O^+][OH^-].$$

Since, $pH = -\log [H_3O^+]$, one needs to calculate $[H_3O^+]$, which can be determined from K_w , once $[OH^-]$ is known. One knows $[OH^-]$ from the fact that one has a 0.2 M NaOH solution. M = molarity = moles/liter. In solution, NaOH dissociates according to



Thus, $[OH^-] = [NaOH]$. Since $[NaOH] = 0.2$, $[OH^-] = 0.2$, which allows for substitution into the equilibrium expression,

$$K_w = 1.0 \times 10^{-14} = [H_3O^+](0.2), \quad \text{or}$$

$$[H_3O^+] = \frac{1.0 \times 10^{-14}}{0.2} = 5.0 \times 10^{-14}, \text{ which means}$$

$$pH = -\log [5.0 \times 10^{-14}] = 13.3.$$

What is the pH when 5 ml of acid ($HC_2H_3O_2$) is added? To answer this, note that a neutralizing reaction occurs, (i.e. a reaction between an acid and base to produce a salt and water) when the acid is added to the NaOH solution.



It is necessary to find how much base is consumed, since this will reflect the remaining $[OH^-]$. From this, return to the expression for K_w and substitute this new value for $[OH^-]$ to determine $[H_3O^+]$ from which the pH can be calculated. To find the amount of NaOH that was consumed, use Molarity = moles/liter.

If the acid is 0.200 M and one adds in 5 ml or 0.005 l (1000 ml = 1 l), one has $(0.2)(0.005) = 0.001$ moles of acid. Amount of base present in moles equals;

$$\text{molarity} \times \text{volume} = (0.2)(0.025) = (0.005) \text{ moles.}$$


According to the reaction, they will react in equimolar amounts. Thus, since only 0.001 moles of acid is present, only 0.001 moles of base react. This leaves 0.004 moles of base (NaOH), and as such 0.004 moles of OH^- . Since

$$\text{volume} = \text{base (l)} + \text{acid (l)} = 0.03,$$

$$M = \frac{0.004}{0.03} = 0.13 \text{ M.}$$

Recalling, $K_w = 1.0 \times 10^{-14} = [H_3O^+][OH^-]$ and that $[OH^-] = 0.13$, such that

$$[H_3O^+] = \frac{1.0 \times 10^{-14}}{0.13} = 7.5 \times 10^{-14}.$$



pH is defined as being equal to $-\log [\text{H}_3\text{O}^+]$. Thus, to solve for pH, one must first determine $[\text{H}_3\text{O}^+]$. This can be done by setting up the equation for the equilibrium of the reaction

$$\text{CH}_3\text{COOH} + \text{H}_2\text{O} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}_3\text{O}^+$$

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} = 1.75 \times 10^{-5}.$$

$$1.75 = \frac{(x)(0.05 + x)}{(0.05 - x)}$$

Since CH_3COOH is a weak acid, there will be little dissociation and x will be very small. Thus one can approximate $0.05 + x \approx 0.05$ and $0.05 - x \approx 0.05$.

• Solving:

$$\frac{x(0.5)}{(0.5)} = 1.75 \times 10^{-5}$$

$$x = [\text{H}_3\text{O}^+] = 1.75 \times 10^{-5}$$

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log (1.75 \times 10^{-5}) = 4.76.$$

Part (b):

A buffer solution contains an acid and its conjugate base, both are moderately weak. What this type of solution does is to prevent a large change in pH when more acid or base is added. In a nonbuffered solution, the pH will change drastically. If a strong acid is added to a buffered solution, it reacts with the base A^- to produce a weak acid.



When a strong base is added the same effect occurs. It reacts with the acid to produce a weak base.



The acid-base ratio is altered somewhat but not to any great extent, thus allowing the pH to stay relatively stable. However, if the number of moles of acid or base added is greater than the number of moles of base or acid present, then the pH will change, thereby destroying the buffer.

It is given that 0.1 M HCl is added. Since HCl is a strong acid, and dissociates completely, the H_3O^+ and Cl^- formed each have an initial concentration of 0.1 M. However, the H_3O^+ will react with the CH_3COO^- and drive the equilibrium to the left by Le Châtelier's principle and produce a weak acid. Therefore, the buffer has converted the acid ions so that they cannot change the pH.

Le Châtelier's principle states that when a stress is brought to bear on a system at equilibrium, the system tends to change so as to relieve the stress. The stress is relieved by driving the reaction to the other side of the equilibrium sign to prevent excesses of ions from building up. The net effect can be seen below.



Before: $0.5-x$ x $0.5 + x$

After add of	$0.5 + 0.1 - x$	x	$0.5 - 0.1 + x$
$0.1 \text{ M H}_3\text{O}^+$	$(0.6 - x)$		$(0.4 + x)$

At the new equilibrium, there will be x moles of H_3O^+ still in solution, which were formed from the redissociation of CH_3COOH . x moles of CH_3COO^- are also formed resulting in the new concentrations of


$$0.6 - x \qquad x \qquad 0.4 + x$$

Upon setting up the equilibrium constant equation, one obtains $[H_3O^+]$.

$$K_a = \frac{[H_3O^+][CH_3COO^-]}{[CH_3COOH]} = 1.75 \times 10^{-5}$$

$$1.75 \times 10^{-5} = \frac{(x)(0.4 + x)}{(0.6 - x)}$$

Using the assumption outlined above, $0.6 - x \approx 0.6$,
and $0.4 + x \approx 0.4$.

Solving:


$$1.75 \times 10^{-5} = \frac{x(0.4)}{(0.6)}$$

$$x = 2.63 \times 10^{-5} = [\text{H}_3\text{O}^+]$$

Since, $\text{pH} = -\log [\text{H}_3\text{O}^+]$, one finds by substitution,

$$\text{pH} = -\log 2.63 \times 10^{-5} = 4.58.$$

Calculate the number of grams of NH_4Cl that must be added to 2 l of 0.1 M NH_3 to prepare a solution of $\text{pH} = 11.3$ at 25°C . The K_b for $\text{NH}_3 = 1.8 \times 10^{-5}$. The equilibrium equation



Solution: To determine how many grams of NH_4Cl must be added to this solution to maintain a pH of 11.3, one must first calculate the $[\text{NH}_4^+]$ needed, because $[\text{NH}_4^+] = [\text{NH}_4\text{Cl}]$ in the solution. This is done by using the expression that describes the dissociation,



The dissociation constant (K_b) for this reaction is 1.8×10^{-5} . K_b is equal to the product of the concentrations of the substances formed divided by the concentration of the reactants. These concentrations are raised to the power of their coefficients in the equation for the reaction. $[\text{H}_2\text{O}]$ is not included because it is assumed to remain constant.

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.8 \times 10^{-5}.$$

To solve for $[\text{NH}_4^+]$, one must know $[\text{OH}^-]$ and $[\text{NH}_3]$ first. One is given that NH_3 solution is .1 M, thus $[\text{NH}_3] = .1$. One can find $[\text{OH}^-]$ by using the definition of pOH . $\text{pOH} = 14.0 - \text{pH} = -\log [\text{OH}^-]$. The pH of this solution is 11.3, therefore $\text{pOH} = 14.0 - 11.3 = 2.7$. One solves for OH^- by using the relation $\text{pOH} = -\log [\text{OH}^-]$. Solving for $[\text{OH}^-]$

$$2.7 = -\log [\text{OH}^-]$$

$$10^{-2.7} = [\text{OH}^-]$$

$$2.0 \times 10^{-3} = [\text{OH}^-]$$

One can now use the expression for K_b to solve for $[\text{NH}_4^+]$

$$1.8 \times 10^{-5} = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = \frac{[\text{NH}_4^+][2.0 \times 10^{-3}]}{[.1]}$$

$$[\text{NH}_4^+] = \frac{(.1)(1.8 \times 10^{-5})}{(2.0 \times 10^{-3})} = 9.0 \times 10^{-4} \text{ M}$$

One is given that 2 l of this solution has been made up. Because, Molarity = number of moles/volume in liters;

$$\text{number of moles} = \text{molarity} \times \text{volume}$$

$$= \left(9.0 \times 10^{-4} \frac{\text{moles}}{\text{liter}} \right) (2 \text{ liters})$$

$$= 18 \times 10^{-4} \text{ moles.}$$

18×10^{-4} moles of NH_4Cl must be added to the solution.
Because moles = weight in grams/molecular weight

$$\text{weight in grams} = \text{moles} \times \text{M.W. (MW of } \text{NH}_4\text{Cl} = 53.5)$$

$$= 18 \times 10^{-4} \text{ moles (53.5 g/mole)}$$


$$= 0.096 \text{ g of } \text{NH}_4\text{Cl} \text{ to be added.}$$

• PROBLEM 437

Calculate the pH of a 0.25 M solution of the salt ethylamine hydrochloride, $\text{C}_2\text{H}_5\text{NH}_3\text{Cl}$. The dissociation constant for the base, ethylamine ($\text{C}_2\text{H}_5\text{NH}_2$) is $K_b = 5.6 \times 10^{-4}$.

Solution: The pH will be determined by three processes occurring simultaneously: dissociation of ethylamine, ionization of ethylamine hydrochloride, and hydrolysis of the ethylammonium cation, $\text{C}_2\text{H}_5\text{NH}_3^+$. All three species are present in an aqueous solution of the salt.


Ethylamine hydrochloride ionizes according to the equation



giving rise to the ethylammonium cation. The ethylammonium cation hydrolysis according to the equation



forming the ethylamine base. Ethylamine base dissociates according to



Let K_w denote the water constant, $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$, and K_b the dissociation constant of ethylamine,

$$K_b = [\text{C}_2\text{H}_5\text{NH}_3^+][\text{OH}^-]/[\text{C}_2\text{H}_5\text{NH}_2] = 5.6 \times 10^{-4}. \text{ Then,}$$

$$\frac{K_w}{K_b} = \frac{[\text{H}_3\text{O}^+][\text{OH}^-]}{[\text{C}_2\text{H}_5\text{NH}_3^+][\text{OH}^-]/[\text{C}_2\text{H}_5\text{NH}_2]} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_5\text{NH}_2]}{[\text{C}_2\text{H}_5\text{NH}_3^+]}$$

$$\frac{10^{-14}}{5.6 \times 10^{-4}} = 1.8 \times 10^{-11} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_5\text{NH}_2]}{[\text{C}_2\text{H}_5\text{NH}_3^+]}$$

Note that this is the equilibrium constant for hydrolysis of the ethylammonium cation.

Let $x = [\text{C}_2\text{H}_5\text{NH}_2]$ be the number of moles per liter of ethylamine formed by hydrolysis of the ethylammonium cation. Then $[\text{H}_3\text{O}^+] = x$, since the balanced equation for hydrolysis states that for every mole of ethylamine formed, one mole of H_3O^+ is formed and one mole of ethylammonium cation is consumed. If we assume that ethylamine hydrochloride dissociates completely, then the number of moles per liter of ethylammonium chloride not hydrolyzed is 0.25 M (from the salt) minus the number of moles per liter which is hydrolyzed (x), so that $[\text{C}_2\text{H}_5\text{NH}_3^+] = 0.25 - x$. Hence,

$$1.8 \times 10^{-11} = \frac{[\text{H}_3\text{O}^+][\text{C}_2\text{H}_5\text{NH}_2]}{[\text{C}_2\text{H}_5\text{NH}_3^+]} = \frac{x \cdot x}{0.25 - x}.$$

Assuming that the number of moles hydrolyzed is much less than 0.25 M, then $0.25 - x \approx 0.25$ and the algebra simplifies to

$$1.8 \times 10^{-11} = \frac{x \cdot x}{0.25} = \frac{x^2}{0.25}, \text{ or}$$

$$x = (0.25 \times 1.8 \times 10^{-11})^{1/2} = 2.1 \times 10^{-6}$$

(which is much smaller than 0.25, justifying our assumption). But $x = [\text{H}_3\text{O}^+] = 2.1 \times 10^{-6}$, hence

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log 2.1 \times 10^{-6} = 5.7.$$


The pH of 0.25 M ethylamine hydrochloride is thus 5.7.

• PROBLEM 438

Calculate the pH of a 0.10 M solution of sodium acetate, NaOAc. The equilibrium constant for the dissociation of acetic acid is 1.8×10^{-5} .


Solution: The pH will be governed by two processes: the ionization of NaOAc, and the hydrolysis of acetate ion, OAc^- .

NaOAc dissociates according to the equation



If we assume that dissociation is complete (a reasonable assumption for dilute salt solutions), the concentrations of Na^+ and OAc^- are equal to the initial concentration of NaOAc: $[\text{Na}^+] = [\text{OAc}^-] = 0.10 \text{ M}$.

Hydrolysis of acetate ion proceeds according to the equation



Let x denote the concentration of undissociated HOAc formed by hydrolysis. Then, since one mole of OH^- is formed and one mole of OAc^- is consumed per mole of HOAc formed, the concentration of hydroxide ion formed is x , and the concentration of acetate ion remaining is $0.10 - x$. That is,

$$[\text{OH}^-] = x, \quad [\text{OAc}^-] = 0.10 - x.$$

These concentrations can be substituted into the expression for the hydrolysis constant,

$$K_h = \frac{[\text{HOAc}][\text{OH}^-]}{[\text{OAc}^-]}$$

to find $x = [\text{OH}^-]$. K_h can be found by using the water constant, $K_w = 1.0 \times 10^{-14} = [\text{H}_3\text{O}^+][\text{OH}^-]$ and the equilibrium constant for dissociation of acetic acid (HOAc),

$$K_a = 1.8 \times 10^{-5} = \frac{[\text{H}_3\text{O}^+][\text{OAc}^-]}{[\text{HOAc}]} . \quad \text{Then,}$$

$$\begin{aligned} K_h &= \frac{[\text{HOAc}][\text{OH}^-]}{[\text{OAc}^-]} = [\text{H}_3\text{O}^+][\text{OH}^-] \times \frac{[\text{HOAc}]}{[\text{H}_3\text{O}^+][\text{OAc}^-]} = \frac{K_w}{K_a} \\ &= \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-5}} = 5.6 \times 10^{-10}. \end{aligned}$$

Substituting the unknown concentrations into the expression gives

$$\begin{aligned} K_h &= 5.6 \times 10^{-10} = \frac{[\text{HOAc}][\text{OH}^-]}{[\text{OAc}^-]} \\ &= \frac{(x)(x)}{0.10-x} \approx \frac{x^2}{0.10} = 10 x^2, \end{aligned}$$

where we have assumed that x is much smaller than 0.10 (hence $0.10 - x \approx 0.10$). Then

$$10 x^2 = 5.6 \times 10^{-10}$$

$$x = (5.6 \times 10^{-11})^{1/2} = 7.5 \times 10^{-6}.$$

Therefore, $[\text{OH}^-] = x = 7.5 \times 10^{-6}$. Using the water constant,

$$[\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14}$$

$$[\text{H}_3\text{O}^+] = \frac{1.0 \times 10^{-14}}{[\text{OH}^-]} = \frac{1.0 \times 10^{-14}}{7.5 \times 10^{-6}} = 1.3 \times 10^{-9}$$


and the pH of the solution is

$$\text{pH} = -\log [\text{H}_3\text{O}^+] = -\log (1.3 \times 10^{-9}) = 8.9.$$

• PROBLEM 439

A chemist mixes together 100 ml of 0.5 M sodium acetate, 100 ml of 0.25 M hydrochloric acid (HCl), and 100 ml of a 1.0 M salt solution. She dilutes this to 1000 ml. Determine the concentrations of all the ions present, undissociated acetic acid, and the final pH of the solution, using K_a of acetic acid = 1.7×10^{-5} and $K_w = 1.0 \times 10^{-14}$.

Solution: To start, determine if any reaction took place in this mixture. If no reaction took place, then the molar concentrations will be numerically the same in 1 liter as the computed moles from the initial concentrations. However, a reaction does take place. Namely,



The H_3O^+ (from HCl) reacts with the acetate ion (OAc^-) from the NaOAc (sodium acetate) to produce acetic acid and water. Start the analysis of the reaction by computing the number of moles involved. M = molarity = moles/liter. Thus, in 100 ml of 0.5 M NaOAc, there are $0.5 (100/1000) = 0.05$ moles of NaOAc. Similar calculations show that 0.025 moles of HCl and 0.1 moles of NaCl (the salt solution) are present. The problem asks one to find the pH, the concentration of the ions present, and the concentration of undissociated acetic acid. This necessitates writing the equilibrium constant expression for this reaction, from which these concentrations can be calculated. Thus, write that K_a , the equilibrium constant, equals

$$\frac{[\text{H}_3\text{O}^+][\text{OAc}^-]}{[\text{HOAc}]}.$$

Suppose the reaction went to completion. If it did, 0.25 moles each of HOAc and excess OAc^- would have been produced. One starts with 0.5 moles and only 1 mole of OAc^- can react per mole of H_3O^+ . If there exist 0.25 moles H_3O^+ , then only 0.25 moles of the OAc^- can react, leaving 0.25 moles in excess. If the reaction, however, stops x moles short of completion, x moles of H_3O^+ would remain and the excess of OAc^- would increase from 0.25 to $0.25 + x$. The HOAc would be reduced to $(0.25 - x)$ moles. Since they are in 1000 ml or 1 liter, these become concentrations that can be substituted into

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{OAc}^-]}{[\text{HOAc}]}, \text{ to give } K_a = \frac{x (0.25 + x)}{(0.25 - x)}.$$

It is given that $K_a = 1.75 \times 10^{-5}$. Equating,

$$\frac{x(0.025 + x)}{(0.025 - x)} = 1.75 \times 10^{-5}.$$

When the K_a for an acid is small, one knows that dissociation is minimal and therefore the concentration of H_3O^+ is very small compared to the concentration of OAc^- and $HOAc$. This means that one can assume that their concentrations will not be appreciably changed by the dissociation. Thus,

$$0.025 - x \approx 0.025 \quad \text{and} \quad 0.025 + x \approx 0.025.$$

The equation for the K_a can be rewritten as

$$K_a = \frac{x(0.025)}{0.025} = 1.75 \times 10^{-5}$$

$$\frac{x(0.025)}{(0.025)} = 1.75 \times 10^{-5}$$

$$x = 1.75 \times 10^{-5}.$$

Therefore, $[H_3O^+] = 1.75 \times 10^{-5} \text{ M}$.

Since, $pH = -\log [H_3O^+]$,

$$pH = -\log [1.75 \times 10^{-5}] = 4.76.$$

$$\begin{aligned} [OAc^-] &= 0.025 + x = 0.025 + 1.75 \times 10^{-5} \\ &\approx 0.025. \end{aligned}$$

$$\begin{aligned} [HOAc] &= 0.025 - x = 0.025 - 1.75 \times 10^{-5} \\ &\approx 0.025. \end{aligned}$$

The other ions: $[Na^+] = 0.150 \text{ M}$ from $NaOAc$ and $NaCl$.
 $[Cl^-] = 0.125 \text{ M}$ from HCl and $NaCl$. Since

$$[H_3O^+][OH^-] = K_w = 1.0 \times 10^{-14}$$

and $[H_3O^+] = 1.7 \times 10^{-5}$,

$$[OH^-] = \frac{1.0 \times 10^{-14}}{1.7 \times 10^{-5}} = 5.9 \times 10^{-10} \text{ M}.$$

• PROBLEM 440

(a) A chemist prepares a 0.01 M solution of $NaC_2H_3O_2$. Find its pH . $K_a = 1.8 \times 10^{-5}$. (b) Determine the pH of 0.1 M solution of NH_4Cl . $K_b = 1.8 \times 10^{-5}$.

Solution: Both parts of this problem involve the

hydrolysis of a salt of a weak acid or base to form an acidic or basic solution. This occurs because water can dissociate to H^+ and OH^- ions, and these can react with ions from weak acids and bases. This necessitates a consideration of K_{hyd} , the hydrolysis constant. Because both $NaC_2H_3O_2$ and NH_4Cl are salts, they exist as ions in aqueous solution. The situation in part (a) is



The K_{hyd} for this is

$$\frac{[HC_2H_3O_2][OH^-]}{[C_2H_3O_2^-]} .$$

K_{hyd} is also defined as K_w/K_a . Therefore,

$$K_{hyd} = \frac{K_w}{K_a} = \frac{[HC_2H_3O_2][OH^-]}{[C_2H_3O_2^-]} .$$

From the equation that depicts the reaction one can see that the $[OH^-] = [HC_2H_3O_2]$, since for every molecule of OH^- , a molecule of $HC_2H_3O_2$ must be generated. One can say

$$\frac{K_w}{K_a} = \frac{[OH^-][OH^-]}{[C_2H_3O_2^-]} \quad \text{or} \quad [OH^-]^2 = \frac{K_w}{K_a} [C_2H_3O_2^-] .$$

K_w/K_a is small enough so that $[C_2H_3O_2^-]$ at equilibrium will have a value as if hydrolysis had not occurred.

Letting B represent $[C_2H_3O_2^-]$ and taking the square root;

$$[OH^-] = \sqrt{\frac{K_w}{K_a} B} .$$

Next take the negative log of the equation. (Remember that $pOH = -\log [OH^-]$, and that K_w is the ionization constant of water, which is 1.0×10^{-14}).

$$pOH = \frac{1}{2} pK_w - \frac{1}{2} pK_a + \frac{1}{2} pB$$

However, $\frac{1}{2} pK_w = 7$ and $pOH = 14 - pH$. As stated,

$$K_w = 1.0 \times 10^{-14} = [H^+][OH^-] \text{ and } [H^+] = [OH^-] .$$

Therefore, $[OH^-]^2$ (or $[H^+]^2$) = 1.0×10^{-14} ,

$$[OH^-] = 1.0 \times 10^{-7}, \text{ and } [H^+] = 1.0 \times 10^{-7}$$

$$pOH = -\log [1 \times 10^{-7}] = 7 \quad pOH + pH = 14 .$$

$$pH = -\log [1 \times 10^{-7}] = 7 . \text{ Therefore, } pOH = 14 - pH .$$

One can now substitute to obtain

$$\text{pH} = 14 - (7 - \frac{1}{2}\text{pK}_a + \frac{1}{2}\text{pB}) \quad \text{or}$$

$$\text{pH} = 7 + \frac{1}{2}\text{pK}_a - \frac{1}{2}\text{pB}.$$

This is the equation wanted. It is known that $\beta = [\text{C}_2\text{H}_3\text{O}_2^-]$. Only when this union is at equilibrium \xrightarrow{B} can this procedure be used (i.e. the K_a of the acid must be $\gg 10^{-10}$). The K_a is given as 1.8×10^{-5} and meets these requirements. The $\text{pK}_a = -\log [1.8 \times 10^{-5}] = 4.74$. $B = 0.01 \text{ M}$ (given) and $\text{pB} = 2.00$.

Now go back to $\text{pH} = 7 + \frac{1}{2}\text{pK}_a - \frac{1}{2}\text{pB}$ and substitute these values to obtain:

$$\text{pH} = 7 + \frac{1}{2}(4.74) - \frac{1}{2}(2.00)$$

$$\text{pH} = 8.37$$

Part (b) is worked out in exactly the same way. One ends up having

$$\begin{aligned}\text{pH} &= 7 - \frac{1}{2}\text{pK}_b + \frac{1}{2}\text{pB} \\ &= 7 - \frac{1}{2}(4.74) + \frac{1}{2}(1.00)\end{aligned}$$

$$\text{pH} = 5.13$$

where $\frac{1}{2}\text{pB} = -\frac{1}{2}\log [0.1] = \frac{1}{2}(1.00)$.

CHAPTER 13



CHEMICAL KINETICS

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 423 to 457 for step-by-step solutions to problems.

The rate of a homogeneous chemical reaction typically depends on the temperature and the concentrations of the species which enter the reaction. For a typical general homogeneous reaction,



experimental results have shown that the rate of disappearance of the reactant A , defined as $(-d[A]/dt)$, can be written as

$$\text{rate} = -k[A]^n[B]^m \quad 13-2$$

In this expression, k , the specific reaction rate constant, is a function of the temperature, but not of the concentrations, and n and m are called the order of the reaction with respect to species A and B respectively. It is important to understand that, while the order and the stoichiometric coefficients for a reaction may be the same, the order is **not** necessarily equal to the stoichiometric coefficient. However, the order is equal to the stoichiometric coefficient for each species in an elementary reaction — a reaction that represents the actual molecular path (or mechanism) the reaction follows. This is discussed briefly under chain reactions.

The Arrhenius equation is the most commonly used equation to express how the specific reaction rate constant, k , varies with temperature.

$$k = k_0 \exp(-E/RT) \{(t^{-1})(\text{conc})^{(1-\text{order})}\} \quad 13-3$$

In this equation, k_0 (the pre-exponential constant) has the same number of units as k and E (the activation energy) has units of energy (e.g., joules). R is

the universal gas constant and T is the absolute temperature measured in kelvins. A common error is the failure to use an absolute temperature for T ; this mistake can lead to very serious errors in reaction rate calculations.

To calculate the order of a reaction, it is important to examine the data. If rate data vs. concentrations are available (presumably at a constant temperature), simply try different orders until Equation 13-2 is satisfied. For example, if the rate at which A is consumed, $-r_A$, is tabulated as a function of the concentration of A , the different values of n can be tried until the correct one is determined.

$$-r_A = k [A]^n \quad 13-4$$

However, it is more common to measure the concentration as a function of time. It is then necessary to integrate Equation 13-2 to obtain an expression, from the rate equation, giving concentration as a function of time (again, presumably at a constant value of temperature).

$$-d[A]/dt = k [A]^n \quad 13-5$$

In order to integrate Equation 13-5, the value of n must be known. The procedure, therefore, is to guess a value of n , say $n = 1$, integrate the equation, and compare the result to the data. If $n = 1$,

$$[A] = [A_0] e^{(-kt)} \quad 13-6$$

where $[A_0]$ is the concentration of A when time, t , equals zero. Equation 13-6 can also be written

$$-\ln \{ [A]/[A_0] \} = kt \quad 13-7$$

$$\text{or } k = (-\ln \{ [A]/[A_0] \})/t \quad 13-8$$

If k is the same for each data point, then the guess that $n = 1$ was correct. If not, other values of n must be tried. If $n \neq 1$, the integrated form of Equation 13-5 is

$$[A]^{(1-n)} = [A_0]^{(1-n)} + kt \quad 13-9$$

A plot of $[A]^{(1-n)}$ vs. t is a straight line of slope k .

If the half-life is measured, these data can also be used to determine the order and specific rate constant of a reaction. The half-life is defined as the time for one half of a reactant to disappear. For first-order reactions, it is

independent of the initial concentration; hence the half-life is a direct measure of the reaction rate constant.

$$k = (\ln 2)/t_{1/2} \text{ (for first-order reactions)} \quad 13-10$$

Equation 13-10 is easily derived from Equation 13-8 where $[A] = [A]_0/2$ at $t = t_{1/2}$. For reactions of any other order (i.e., $n \neq 1$), the half-life depends on the initial concentration and the rate constant and is given by the equation.

$$k = \{2^{(n-1)} - 1\} / \{(n-1)[A]_0^{(n-1)} t_{1/2}\} \quad 13-11$$

The activation energy of a reaction, which can be loosely interpreted as the energy the reactants must possess to react, can be determined by measuring the rate constant at several different temperatures. If the Arrhenius equation (Equation 13-3) is written differently,

$$\ln k = \ln k_0 - E/RT \quad 13-12$$

it becomes apparent that a plot of $\ln k$ vs. $1/T$ will be a straight line with a slope of $-E/R$. If the specific rate constant is known at two temperatures, T_1 and T_2 , E can be calculated from the equation

$$E = R \ln(k_1/k_2) / (1/T_2 - 1/T_1) \quad 13-13$$

A common "rule of thumb" states that reaction rates double with a 10°C increase in temperature. For this rule to apply for reactions at or near room temperature, let us calculate what the activation energy must be. For the rate to double between 25°C and 35°C , the specific reaction rate constant must double.

$$k_{308\text{K}} = 2 k_{298\text{K}} \quad 13-14$$

$$k_0 e^{-[E/308R]} = 2 k_0 e^{-[E/298R]} \quad 13-15$$

$$-E/(308R) = (\ln 2) - E/(298R) \quad 13-16$$

$$E = R (\ln 2) / \{1/298 - 1/308\} = 52.87 \text{ KJ} \quad 13-17$$

Problem 469 asks you to repeat the calculation above for different temperatures; its solution follows the procedure above with different values of temperature. Problem 468 asks what activation energy will result if a tripling of the rate occurs for a 10°C increase in temperature. The solution is identical to that just completed except that 3 replaces 2 in Equation 13-14 and is carried through to the subsequent equations.

Chain reactions are reactions that take place in sequence. Two types are discussed in the problems of Chapter 13. The first, illustrated in Problem 474, is two consecutive first-order reactions. The quantitative solutions for the concentrations of A , B , and C are given here, which expands on the qualitative answers requested in the problem.



$$A = [A_0] e^{(-k_1 t)} \quad 13-19$$

$$B = \{k_1 / (k_2 - k_1)\} [A_0] \{e^{(-k_1 t)} - e^{(-k_2 t)}\} \quad 13-20$$

$$C = \{A_0 / (k_2 - k_1)\} \{k_2 [1 - e^{(-k_1 t)}] - k_1 [1 - e^{(-k_2 t)}]\} \quad 13-21$$

Note also that

$$C = A_0 - A - B. \quad 13-22$$

The second type of chain reaction discussed is a reaction mechanism. Mechanisms are series of elementary reactions that represent the actual path followed on a molecular scale in an overall reaction. The key to solving the problems is to assume, for short-lived intermediates, that the concentration does not change with time (i.e., $dC/dt = 0$ for short-lived intermediates). This is called the steady state assumption and permits the rate equations written for any step in the mechanism containing a short-lived species to be greatly simplified.

Step-by-Step Solutions to Problems in this Chapter, "Chemical Kinetics"

THE RATE LAW

• PROBLEM 441

A group of mountain climbers set up camp at a 3 km altitude and experience a barometric pressure of 0.69 atm. They discover that pure water boils at 90°C and that it takes 300 minutes of cooking to make a "three-minute" egg. What is the ratio of the rate constant $k_{100^\circ\text{C}}$ and $k_{90^\circ\text{C}}$?

Solution: Since we do not know the rate expression for cooking an egg, we will assume one of the form

$$\text{rate} = k [A]^m [B]^n \dots$$

where k is the rate constant, A , B , ... are the reactants, and the overall order of the reaction is $m + n + \dots$. We will write the rate equations at the normal boiling point of water (100°C) and at 90°C as

$$\text{rate}_{100^\circ\text{C}} = k_{100^\circ\text{C}} [A]^m [B]^n \dots \quad \text{and}$$

$$\text{rate}_{90^\circ\text{C}} = k_{90^\circ\text{C}} [A]^m [B]^n \dots$$

Dividing the first of these by the second gives

$$\frac{\text{rate}_{100^\circ\text{C}}}{\text{rate}_{90^\circ\text{C}}} = \frac{k_{100^\circ\text{C}} [A]^m [B]^n \dots}{k_{90^\circ\text{C}} [A]^m [B]^n \dots} = \frac{k_{100^\circ\text{C}}}{k_{90^\circ\text{C}}}$$


Since the egg cooks 100 times faster at 100°C than at 90°C (300 min/3 min = 100), $\text{rate}_{100^\circ\text{C}}/\text{rate}_{90^\circ\text{C}} = 100$.

Hence,

$$\frac{\text{rate}_{100^{\circ}\text{C}}}{\text{rate}_{90^{\circ}\text{C}}} = 100 = \frac{k_{100^{\circ}\text{C}}}{k_{90^{\circ}\text{C}}}, \text{ or, } \frac{k_{100^{\circ}\text{C}}}{k_{90^{\circ}\text{C}}} = 100.$$

• PROBLEM 442

The reaction



was studied kinetically and the following data was obtained.

Experiment	A	B	Rate (mole/liter - min)
1	1.0 M	1.0 M	0.15
2	2.0 M	1.0 M	0.30
3	1.0 M	2.0 M	0.15

Determine the rate expression.

Solution: The rate of reaction is equal to some rate constant, k , multiplied by the concentrations of A and B raised to the appropriate powers. That is,

$$\text{rate} = k [\text{A}]^m [\text{B}]^n$$

where the exponents m and n are to be determined.


Comparing experiments 2 and 1, we see that holding $[\text{B}]$ constant while doubling $[\text{A}]$ doubles the rate of reaction (from 0.15 to 0.30). Hence, the rate is directly proportional to $[\text{A}]$ and $m = 1$. Thus, if we hold $[\text{B}]$ constant and triple $[\text{A}]$, the rate triples.

Comparing experiments 3 and 1, we see that holding $[\text{A}]$ constant and changing $[\text{B}]$ (from 1.0M to 2.0M) has no effect on the rate. Hence, the rate is independent of $[\text{B}]$ and $n = 0$ (so that $[\text{B}]^0 = 1$ and $[\text{B}]$ does not appear in the rate expression).

Substituting $m = 1$ and $n = 0$ into the rate expression gives

$$\text{rate} = k [\text{A}]^1 [\text{B}]^0, \quad \text{or,} \quad \text{rate} = k [\text{A}].$$

Two different molecules, A and B, were reacted to give products according to the equation



The rate of reaction was measured for fixed concentrations of A and B and the following data was obtained:

Experiment	[A] (moles/liter)	[B] (moles/liter)	Rate (moles ² /liter ² -sec)
1	1.0	1.0	0.05
2	1.0	2.0	0.10
3	3.0	1.0	0.45

Write an expression for the rate of reaction.

Solution: The rate of reaction is equal to some rate constant, k , multiplied by the concentrations of A and B raised to appropriate powers. That is,

$$\text{rate} = k [A]^m [B]^n$$

where the exponents m and n are to be determined.

Comparing experiment 2 with experiment 1, we see that when the concentration of B is doubled from 1.0 M to 2.0 M, the concentration of A being held constant at 1.0 M, the rate of reaction doubles from 0.05 to 0.10 moles²/liter²-sec. Hence, the rate of reaction is directly proportional to the concentration of B, and the exponent of [B] in the rate expression is $n = 1$. If we were to triple [B], holding [A] constant, the rate would triple, and so on.

Comparing experiment 3 with experiment 1, we see that when the concentration of A is tripled from 1.0 M to 3.0 M, the concentration of B being held constant at 1.0 M, the rate of reaction is multiplied by 9 ($9 \times 0.05 = 0.45$ moles²/liter² - sec). Since the factor by which the rate is multiplied (9) is the square of the factor by which the concentration of A is multiplied (3), the exponent of [A] in the rate expression must be $m = 2$. Thus if [A] is doubled, [B] being held constant, the rate is multiplied by 4.

Substituting the exponents of [A] and [B] determined above into the rate expression gives

$$\text{rate} = k [A]^2 [B].$$

For the reaction



at 1100°K, data, as shown in the following table, were obtained. Find the rate law and the numerical value of the specific rate constant.

<i>Initial Pressure of NO, atm</i>	<i>Initial Pressure of H₂, atm</i>	<i>Initial rate of pressure decrease, atm/min</i>
0.150	0.400	0.020
0.075	0.400	0.005
0.150	0.200	0.010

Solution: The rate of a homogeneous reaction, i.e. one which occurs in only one phase, depends on the concentration of reactants in that phase. This reaction is a homogeneous reaction in the gaseous phase. It can be investigated kinetically by following the change in pressure of the gaseous mixture as the reaction proceeds. The pressure drops because 3 moles of gaseous reactants are converted to two moles of gaseous products. Since the reactants are being used up during the course of the reaction, their concentrations and their rate of reaction are constantly changing. The concentrations and rates listed are those at the very beginning of the reaction, when little change has occurred. From the data given, one can see that when the initial pressure of NO is halved with the pressure of H₂ remaining constant, the initial rate is quartered. When the pressure of H₂ is halved and the NO remains constant, the initial rate is halved. When the pressure of H₂ is doubled and NO remains constant, the rate doubles. We conclude that the rate is proportional to [H₂]. When the H₂ pressure is kept constant and the NO concentration is doubled, the rate is quadrupled, this means that the rate is proportional to [NO]². The equation for the rate can thus be written

$$\text{Rate} = k [\text{H}_2][\text{NO}]^2$$

where k is the rate constant.

Using the values in the first trial of the table one can solve for k .

$$0.020 \text{ atm/min} = k (0.400 \text{ atm})(0.150 \text{ atm})^2$$

$$\frac{0.020 \text{ atm/min}}{(0.400 \text{ atm})(0.150 \text{ atm})^2} = k$$

$$k = 2.22 \text{ atm}^{-2} \text{ min}^{-1}.$$

• PROBLEM 445

Assume that an A molecule reacts with two B molecules in a one-step process to give AB_2 . (a) Write a rate law for this reaction. (b) If the initial rate of formation of AB_2 is 2.0×10^{-5} M/sec and the initial concentrations of A and B are 0.30 M, what is the value of the specific rate constant?

Solution: (a) The overall equation for this reaction is



Since no other information is provided about the reaction, the rate law for the reaction is assumed to be written

$$\text{Rate} = k [A][B]^2,$$

where k is the rate constant and $[\]$ indicates concentration.

(b) One can solve for k using the rate law when the rate, $[A]$ and $[B]$ are given as they are in this problem.

$$\text{Rate} = k [A][B]^2$$

$$2.0 \times 10^{-5} \text{ M/sec} = k (0.30 \text{ M}) (0.30 \text{ M})^2$$

$$\frac{2.0 \times 10^{-5} \text{ M/sec}}{(0.30 \text{ M}) (0.30 \text{ M})^2} = k$$

$$k = 7.41 \times 10^{-4} \text{ M}^{-2} \text{ sec}^{-1}.$$

• PROBLEM 446

A reacts with B in a one-step reaction to give C. The rate constant for the reaction is $2.0 \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1}$. If 0.50 mole of A and 0.30 mole of B are placed in a 0.50 liter-box, what is the initial rate of the reaction?

Solution: The equation for this reaction can be written:



From this, one can write the rate law assuming that the reaction is first order in both A and B. When a reaction is first order in a particular reactant, it means that the rate is proportional to the concentration of the reactant. Thus, the rate law is written

$$\text{Rate} = k[A][B],$$


where k is the rate constant and $[\]$ indicates concentration. In this problem, one is given k , $[A]$ and $[B]$. Thus,

$$\begin{aligned}\text{Rate} &= 2.0 \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1} \\ &\times (1 \text{ M})(.60 \text{ M}) \\ &= 1.20 \times 10^{-3} \text{ M/sec}\end{aligned}$$

$$\begin{aligned}k &= 2.0 \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1} \\ [\text{A}] &= 0.50 \text{ mole}/0.5 \text{ liter} \\ &= 1 \text{ M} \\ [\text{B}] &= 0.30 \text{ mole}/0.5 \text{ liter} \\ &= .60 \text{ M}.\end{aligned}$$

• PROBLEM 447

In studying the kinetics of the reaction



at 800°K, the data in the following table were observed. (a) Write the rate law for this reaction. (b) What is the numerical value of the specific rate constant? (c) What would be the initial rate of Z formation starting with 0.15 M X and 0.15 M Y? (d) How would the rate in (c) be changed if, after the reaction had just begun, the volume of the container were abruptly doubled?

<i>Initial concentration of X, M</i>	<i>Initial concentration of Y, M</i>	<i>Initial rate of formation of Z, M/min</i>
0.10	0.10	0.030
0.20	0.20	0.240
0.20	0.10	0.120

Solution: (a) All of the components in this system are in the gaseous phase, thus the system is homogeneous. The rate of a reaction in a gaseous homogeneous system is dependent upon the concentration of the reactants. In a gaseous system, concentration is a function of the pressure of the system. From the data given in the table, one can see that when the concentration of Y is held constant and the concentration of X is doubled, the rate is quadrupled. Therefore, the rate is proportional to $[\text{X}]^2$. When the concentration of X is held constant and the concentration of Y is doubled, the rate is doubled and is thus proportional to $[\text{Y}]$. The rate law for this reaction can thus be written $\text{Rate} = k [\text{X}]^2 [\text{Y}]$ where k is the rate constant.

(b) The numerical value of k is found by substituting in values for $[\text{X}]$, $[\text{Y}]$ and the rate into the rate law. Concentration values for X and Y may be obtained from any one line of the table. The values obtained from the first line of the table are

$$0.030 \text{ M/min} = k (0.10 \text{ M})^2 (0.10 \text{ M})$$

$$\frac{0.030 \text{ M/min}}{(0.10 \text{ M})^2 (0.10 \text{ M})} = k$$

$$30.0 \text{ M}^{-2} \text{ min}^{-1} = k$$

(c) One can solve for the initial rate of the formation of Z by using the rate law and k when [X] and [Y] are given. Thus,

$$\text{Rate} = k[X]^2[Y]$$

$$\begin{aligned} \text{Rate} &= (30.0 \text{ M}^{-2} \text{ min}^{-1}) (0.15 \text{ M})^2 (0.15 \text{ M}) \\ &= .101 \text{ M/min.} \end{aligned}$$

(d) According to Boyle's Law the pressure is inversely proportional to the volume. Therefore, if the volume is doubled the pressure is halved. Because the reactants in this system are gases, when the pressure is halved, the concentration is halved. One can find the rate of the reaction described in (c) by using the rate law with the concentrations of the reactants halved

$$\begin{aligned} \text{Rate} &= 30.0 \text{ M}^{-2} \text{ min}^{-1} (\frac{1}{2}(0.15 \text{ M}))^2 (\frac{1}{2}(0.15 \text{ M})) \\ &\neq 30.0 \text{ M}^{-2} \text{ min}^{-1} (.075 \text{ M})^2 (.075 \text{ M}) \\ &= 1.27 \times 10^{-2} \text{ M/min} \end{aligned}$$


The percent of the first rate that the second rate is equal to is found by dividing the second rate by the first and then multiplying the quotient by 100.

$$\frac{\text{Rate}_2}{\text{Rate}_1} \times 100 = \frac{1.27 \times 10^{-2}}{1.01 \times 10^{-1}} \times 100 = 12.5\% \text{ or } 1/8.$$

The second rate is, therefore, 1/8 of the first.

● PROBLEM 448

Photochemical smogs are thought to be partially initiated by the photolytic decomposition of NO₂ to form NO and O. When the subsequent reaction



is studied kinetically, it is found that when the initial concentration of O₂ is doubled and that of NO held constant, the initial reaction rate doubles; and when the initial concentration of NO is doubled and that of O₂ held constant, the initial reaction rate quadruples. Write the rate expression for this reaction.

Solution: The rate of reaction is equal to some rate constant k multiplied by the concentrations of NO and O_2 raised to the appropriate powers. That is,

$$\text{rate} = k [\text{NO}]^m [\text{O}_2]^n$$

where the exponents m and n are to be determined.

When $[\text{NO}]$ is held constant and $[\text{O}_2]$ is doubled, the rate is doubled. Hence, the rate is directly proportional to $[\text{O}_2]$ and $n = 1$. Thus, if we were to hold $[\text{NO}]$ constant and triple $[\text{O}_2]$, the rate is tripled.

When $[\text{O}_2]$ is held constant and $[\text{NO}]$ is doubled, the reaction rate is quadrupled. Hence, the rate is second order in $[\text{NO}]$ and $m = 2$. Thus, if we were to hold $[\text{O}_2]$ constant and triple $[\text{NO}]$, the rate is multiplied by $3^2 = 9$.

Substituting $m = 2$ and $n = 1$ into the rate expression above gives

$$\text{rate} = k [\text{NO}]^2 [\text{O}_2]^1, \quad \text{or, } \text{rate} = k [\text{NO}]^2 [\text{O}_2].$$

• PROBLEM 449

The reaction



was studied kinetically and the following data was obtained.

Experiment	[A] (moles/liter)	[B] (moles/liter)	Rate (mole/liter - sec)
1	2.0×10^{-3}	4.0×10^{-3}	5.7×10^{-7}
2	2.0×10^{-3}	8.0×10^{-3}	11.4×10^{-7}
3	4.0×10^{-3}	4.0×10^{-3}	22.8×10^{-7}

Determine the rate law for this reaction.

Solution: The rate of reaction is equal to some rate constant, k , multiplied by the concentrations of A and B raised to the appropriate powers. That is,

$$\text{rate} = k [\text{A}]^m [\text{B}]^n$$

where the exponents m and n are to be determined. The presence of the coefficient 2 in the reaction equation



in no way affects the form of the rate equation.

Comparing experiments 2 and 1 we see that doubling [B] while holding [A] constant doubles the rate of reaction ($2 \times 5.7 \times 10^{-7} = 11.4 \times 10^{-7}$). Hence, the rate is directly proportional to [B] and $n = 1$.

Comparing experiments 3 and 1 we see that doubling [A] while holding [B] constant quadruples the rate ($4 \times 5.7 \times 10^{-7} = 22.8 \times 10^{-7}$). Thus, the rate has a square dependence on [A] ($2^2 = 4$), that is, it is second order in [A], and $m = 2$.

Substituting $m = 2$ and $n = 1$ into the rate expression gives

$$\text{rate} = k [A]^2 [B]^1, \text{ or } \text{rate} = k [A]^2 [B].$$

• PROBLEM 450

You have the general reaction: $A + 2B + C = AB + BC$. You collect the following data:

	Concentration(moles/liter)			
	[A]	[B]	[C]	$-dA/dt$ (moles/liter-sec)
(1)	1.00	1.00	2.00	1.00
(2)	2.00	1.00	2.00	2.00
(3)	2.00	2.00	2.00	8.00
(4)	2.00	2.00	4.00	8.00

(a) Determine the experimental rate law expression. (b) Find the specific reaction rate constant. (c) Calculate the rate of reaction, if the [A], [B], and [C] are, respectively, 1.0, 2.0, and 3.0: (d) Speculate on the rate-controlling step, i.e., the slow step.

Solution: First write a general rate law for the reaction, and then write a specific law based on the given data. The specific rate constant can be found from a given set of data. The rate with given concentrations is obtainable from the specific rate law. From this, a rate-controlling step can be guessed at. Proceed as follows:

(a) To write the rate law, note that, in general, the rate $= r = k[A]^x[B]^y[C]^z$, where the brackets indicate concentrations of A, B, and C, k = specific rate constant and the exponents (x , y , and z) are the orders of the reaction with respect to each reactant. The order of a chemical reaction is the number of particles needed to form the activated complex or transition state. To find the actual law, you must determine these exponents. To do this, consult the data. You are told the actual rate of the re-

action, since you are given $-dA/dt$, which is an indication of rate. Thus, to find the order, see what happens when one of the concentrations is changed. In going from (1) to (2), the concentration of A is doubled from 1.00 to 2.00. Notice: $-dA/dt$ also doubles, from 1.00 to 2.00, which means the reaction is first order, $x = 1$, with respect to A. In going from (2) to (3), [B] goes from 1.00 to 2.00 (doubles). But notice, $-dA/dt$ quadruples going from 2.00 to 8.00. Thus, the reaction is second order, $y = 2$, with respect to B. In going from (3) to (4), [C] is doubled going from 2.00 to 4.00, while $-dA/dt$ remains constant at 8.00. Thus, the reaction is zero order, $z = 0$, with respect to C, i.e., rate doesn't depend on [C]. You then have, $r = k[A][B]^2[C]^0$ or $r = k[A][B]^2$ for the experimental rate law expression.

(b) To find k , substitute any of the four sets of data in the experimental rate law expression.

$$\text{rate} = \frac{-dA}{dt} = k[A][B]^2$$

For example, let us take the data from line (1). You have 1.00 mole/liter-sec = $k[1.00 \text{ mole/liter}][1.00 \text{ mole/liter}]^2$, or $k = 1.00$. Thus, the value of the specific rate constant k is 1.00 liter²/mole² - sec.


(c) To find the rate of the reaction at the concentrations [A] = 1, [B] = 2, and [C] = 3, substitute these values into the rate law, $r = k[A][B]^2$. Thus, $r = (1)[1][2]^2 = 4.0$ moles/liter-sec.

(d) You can use previous information gained in this problem to speculate about what the rate-controlling step must be. From part (a) you know the rate does not depend on [C]. Thus, C cannot be involved in the slow step. The rate of the reaction is influenced by [A] and [B], however. The rate-controlling step is probably the combination of B₂ and A, i.e., $B_2 + A \xrightarrow{\text{slow}} AB_2$.

THE ORDER OF REACTIONS

• PROBLEM 451

Under certain conditions, the rate equation for the formation of water



is given by

$$\text{rate} = k [\text{H}_2]^2 [\text{H}_2\text{O}]$$

where k is the rate constant. What is the overall order of this rate equation?

Solution: The overall order of a rate equation is equal to the sum of the exponents to which the concentrations are raised. In the equation

$$\text{rate} = k [\text{H}_2]^2 [\text{H}_2\text{O}],$$

$[\text{H}_2]$ is raised to the second power and $[\text{H}_2\text{O}]$ is raised to the first power. Hence, the rate is second order in $[\text{H}_2]$, first order in $[\text{H}_2\text{O}]$, and $2 + 1 = 3$, or third order overall.

• PROBLEM 452

You have the general reaction: $\text{A} + \text{B} \rightarrow \text{products}$. Consider the following reaction rate data and then determine the rate law for each case: (a) When $[\text{A}]$ is doubled, the initial rates doubles; when $[\text{B}]$ is doubled, the initial rate doubles, (b) When $[\text{A}]$ doubles, initial rate doubles, when $[\text{B}]$ doubles, the initial rate is halved, (c) Doubling $[\text{A}]$, doubles initial rate; but doubling $[\text{B}]$ leaves the rate unchanged.

Solution: The given reaction is one with two different molecules being converted to products. The rate of any such reaction is:

$$r = \text{rate} = k [\text{A}]^x [\text{B}]^y, \quad \text{where } k = \text{a rate}$$

constant and $[\text{A}]$ = concentration of A and $[\text{B}]$ = concentration of B. x and y are the number of each A and B molecules, respectively, that are present in the activated complex or transition state. The determination of these exponents is the key to writing the rate law for each case. You proceed as follows:

(a) You are told that when A or B is doubled, the rate is doubled. Thus, when $\text{A} \rightarrow 2\text{A}$, $r \rightarrow 2r$, which means $2r = k [2\text{A}]^x [\text{B}]^y$. Now, if $x = 1$, then $[2\text{A}]^x = [2\text{A}]$ and dividing by 2 you come back to $r = k [\text{A}][\text{B}]^y$. In other words, $r \rightarrow 2r$ if $\text{A} \rightarrow 2\text{A}$, implies r varies directly as A; that is, $x = 1$. The same type of situation exists with B, so that $y = 1$. Thus, the rate law becomes $r = k [\text{A}][\text{B}]$.

(b) Here, when $\text{A} \rightarrow 2\text{A}$, $r \rightarrow 2\text{A}$, which means, again, A varies directly as r , and so $x = 1$. When $\text{B} \rightarrow 2\text{B}$, $r \rightarrow \frac{1}{2}r$, however, in this case, $\frac{1}{2}r = [\text{A}][2\text{B}]^y$. This can only come about if $y = -1$. If $y = -1$, then $[2\text{B}]^{-1} = 1/[2\text{B}]$, so that $\frac{1}{2}r = [\text{A}] 1/[2\text{B}]$. The halves can be cancelled to return

a direction proportion. Thus, the rate law becomes $r = k[A]/[B]$.

(c) Here, when $[A] \rightarrow [2A]$, $r \rightarrow 2r$ but when $[B] \rightarrow [2B]$, $r \rightarrow r$. For A: you have, again, a direction proportion between A and r , which requires $x = 1$. For B: when B's concentration is doubled, nothing happens to r ; it remains constant. This means r does not depend upon the $[B]$. For r not to depend upon $[B]^y$, y must equal zero, thus, $r = k \frac{[A]}{[B]^0} = k [A](1)$. As such, the rate law becomes $r = k [A]$.

• PROBLEM 453

From collision theory, it is found that when the rate-determining step involves collision of two A molecules, the rate will be proportional to the square of the A concentration. Explain why.


Solution: Collision theory makes the assumption that for a chemical reaction to take place, particles must collide. The rate will depend upon the number of collisions per unit time between the particles involved and the fraction of these collisions that are effective. If you increase $[A]$, there are more A molecules which can be hit. Also, the average A molecule gets hit more often. As such, the number of A with A collisions varies as the square of concentration due to this double effect.

• PROBLEM 454

For the general reaction $A + B \rightarrow C + D$, what is the effect on the number of collisions between A and B of (a) tripling the concentration of each, (b) quadrupling the concentration of each?

Solution: For any given chemical reaction, the number of collisions is proportional to the rate. The Law of Mass Action states that the speed of a chemical reaction is proportional to the product of the concentrations of the reacting molecules. The number of collisions in the above reaction depends upon the concentration of A and B. If the concentration of A is tripled the number of collisions is tripled, then if the concentration of B is tripled the number of collisions will triple again. Thus the number of collisions is increased 3×3 times or 9 times. When the concentrations of A and B are quadrupled a similar method is used. The number of collisions is increased 4×4 or 16 times.

The following data was collected for the reaction



Time, min	$[\text{CH}_3\text{COOCH}_2\text{CH}_3]$	$[\text{OH}^-]$
0	.02000	.02000
5	.01280	.01280
15	.00766	.00766
25	.00540	.00540
35	.00426	.00426
55	.00289	.00289
120	.00137	.00137

(a) Determine whether the reaction is first or second order. (b) Write the rate law expression.

Solution: The order of a reaction is the number of particles needed to form the transition state. One can tell whether the reaction is first or second order by an investigation into the constancy of the half-life of this reaction over time. It is a characteristic feature of first order reactions that the half-life, the time necessary for half the particular reactant present initially to decompose, is constant over time. Thus, by determining whether the half-life is constant or not, the order of this reaction can be found.

From the data, half the material is decomposed between 5 and 15 minutes. In the period, 5 to 15 minutes, the concentration of both $[\text{CH}_3\text{COOCH}_2\text{CH}_3]$ and $[\text{OH}^-]$ goes from .0128 to .00766, which means half has decomposed. One can approximate that the half-life is about 10 minutes. To see if this time of half-life remains constant, notice the time needed for the concentration to go from .0054 to .00270, which is another $\frac{1}{2}$ being decomposed. This occurs between $t = 25$ minutes to approximately $t = 55$ minutes, a difference of 30 minutes. Thus, the half-life is not constant. It changed from 10 to 30 minutes. Thus, the reaction is not first order. It must be second order.

(b) To solve this part, use the information from (a). If the reaction is second order, the rate depends upon both the concentration of $\text{CH}_3\text{COOCH}_2\text{CH}_3$ and OH^- . The rate law expression is, thus,

$$\text{rate} = -\frac{d[\text{CH}_3\text{COOCH}_2\text{CH}_3]}{dt} = -\frac{d[\text{OH}^-]}{dt} = k[\text{OH}^-][\text{CH}_3\text{COOCH}_2\text{CH}_3],$$

where k = rate constant. This equation is, by definition, the rate law expression for a second order reaction with 2 different molecules.

The reaction $\text{C}_2\text{H}_6\text{O} + \text{CH}_4 + \text{CO}$ is a first-order reaction with a specific rate constant of $.0123 \text{ min}^{-1}$ at 415°C . Calculate the percent of original $\text{C}_2\text{H}_6\text{O}$ present that will be decomposed after heating at 415°C for 1 hour.

Solution: First find the percentage of starting material left (as compared to the initial amount C_0), $C/C_0 = (100)$; then the percent decomposition equals $1 - C/C_0 (100)$. To find $C/C_0 (100)$, employ the fact that $2.303 \log C_0/C = kt$, where C_0 = initial concentration, C = existing concentration, k = specific rate constant, and t = time in minutes for a first order reaction. The formula can be rewritten as $C/C_0 = \text{antilog} (-kt/2.303)$.

Substitute the values for k and t to obtain

$$\frac{C}{C_0} = \text{antilog} \left(\frac{(-.0123)(60)}{2.303} \right) = \text{antilog} (-.320) = .479.$$

Thus, $C/C_0(100) = 100 \times .479 = 47.9\%$. That is, $.479$ or 47.9% of the original concentration remains after 60 minutes, so that $1 - 47.9\%$ or 52.1% has decomposed.

The following observations were made for the reaction $\text{NH}_4\text{CNO} \rightleftharpoons \text{NH}_4^+ + \text{CNO}^- \rightarrow (\text{NH}_2)_2\text{CO}$: 1) Addition on an equimolar concentration of KCNO with respect to NH_4CNO doubled the initial rate of the reaction. 2) Addition of an equimolar concentration of NH_4Cl doubled the initial rate of reaction. From these observations, (a) Determine the order of the reaction. (b) Write the kinetic rate law expression for the reaction. (c) Discuss the influence, if any, of the following equilibrium upon the reaction kinetics: $\text{NH}_4^+ + \text{CNO}^- \rightleftharpoons \text{NH}_3 + \text{HCNO}$ with

$$K_{\text{equil}} = \frac{[\text{NH}_3][\text{HCNO}]}{[\text{CNO}^-][\text{NH}_4^+]} = 10^{-4}.$$

Solution: (a) The order of a chemical reaction is the number of distinct particles that must come together to form the activated complex. This factor can also be called the molecularity of the reaction. To see which particles directly determine the reaction rate, note the effect on the rate when more of a certain type of particle is added. You are told that when KCNO is added, that is, the KCNO concentration is doubled, the rate doubles. This means the reaction must be first order with respect to CNO^- (since, in solution, KCNO dissociates to CNO^-). When $[\text{NH}_4\text{Cl}]$ is doubled, the reaction rate is again doubled. Thus, the reaction is first order with respect to NH_4^+ , since, in

solution, NH_4Cl dissociates into NH_4^+ . Since doubling the amount of either NH_4^+ or CNO^- doubles the reaction rate, the reaction must be second-order overall.

(b) By definition, a second order reaction with two different molecules has rate $= -d[\text{A}]/dt = -d[\text{B}]/dt = k[\text{A}][\text{B}]$, where k = rate constant and A and B are the two different molecules in $\text{A} + \text{B} \rightarrow \text{products}$. Thus, for this reaction, the rate law becomes

$$\text{rate} = -\frac{d[\text{NH}_4\text{CNO}]}{dt} = \frac{d[(\text{NH}_2)_2\text{CO}]}{dt} = k[\text{NH}_4^+][\text{CNO}^-]$$

(c) To answer this part, notice the following: The equilibrium constant for the formation of NH_3 and HCNO is extremely small, which means that, in solution, the concentration of these species in pure NH_4CNO solution should also be small, i.e.,

$$\frac{[\text{NH}_3]}{[\text{NH}_4^+]} = \frac{[\text{HCNO}]}{[\text{CNO}^-]} = (10^{-4})^{1/2} = .01.$$

If either NH_3 or HCNO is added to the solution, the reaction is driven to the left. But, even if all of the other reactants available were consumed, the concentration of NH_4^+ and CNO^- would increase by only 1%, which means that the reaction rate would increase by 2%, which is not easily detected.

• PROBLEM 458

When mixed at 700°K , $\text{H}_2(\text{g})$ and $\text{I}_2(\text{g})$ react to produce $\text{HI}(\text{g})$. The reaction is first order in H_2 and first order in I_2 . Suppose at time $t = 0$, one mole of H_2 and one mole of I_2 are simultaneously injected into a 1-liter box. One second later, before the reaction is complete, the contents of the box are examined for the number of moles of HI . What would be the probable effect on this number if each of the following changes were made in the initial conditions? (a) Use two moles of H_2 instead of one. (b) Use two moles of I_2 instead of one. (c) Use a 2-liter box. (d) Raise the temperature to 750°K . (e) Add a platinum catalyst. (f) Add enough neon gas to double the initial pressure.

Solution: When a reaction is said to be first order in a particular reactant, it means that the rate of the reaction is proportional to the concentration of that reactant. Thus, for this reaction the rate law can be written

$$\text{Rate} = k[\text{H}_2][\text{I}_2],$$

where k is the rate constant and $[\]$ indicates concentration. When 1 mole of each H_2 and I_2 are used, one can solve for

the rate by substituting these values into the rate law.

$$\text{Rate} = k (1) (1) = 1k$$

Solving for the rate when $[H_2]$ is 2 moles

$$\text{Rate} = k[2][1] = 2k$$

Thus, when the concentration of H_2 is doubled the rate is doubled.

(b) One can solve for the rate when $[I_2] = 2$ moles by substituting into the rate law.

$$\text{Rate} = k(1)(2) = 2k$$

Thus when $[I_2]$ doubled the rate is also doubled.

(c) Boyle's law states that the pressure is inversely proportional to the volume. Thus, when this reaction which originally takes place in a 1-liter box is moved to a 2-liter box, the pressure of the reactants is halved. Solving for the rate:

$$\text{Rate} = k \left(\frac{1}{2}\right) \left(\frac{1}{2}\right) = \frac{1}{4} k \quad \text{or} \quad .25 k.$$

Thus, for this reaction, when the pressure of the reactants is halved, the rate is $\frac{1}{4}$ of the original rate.

(d) Product is formed when an atom of H collides with an atom of I, thus the rate of the reaction will increase with a rise in temperature because the atoms will move more quickly creating a greater chance for collisions.

(e) By definition, a catalyst will increase the rate of the reaction by lowering the amount of energy needed for the reaction to proceed.

(f) When neon gas is added to increase the pressure in the 1-liter box, the rate of the reaction does not change because the concentrations of H_2 and I_2 remain unchanged. When the pressure in the system is doubled by decreasing the volume by $\frac{1}{2}$ the rate is increased because the concentrations of H_2 and I_2 are increased. If one starts out with one mole per liter of H_2 at 1 atm and then increases the pressure to 2 atm by adding neon, the concentration of H_2 is still 1 mole per liter. When the pressure in a container containing 1 mole per liter of H_2 is increased from 1 to 2 atm by halving the volume of the container, the concentration of H_2 is now 2 moles per liter or 2 moles per liter. Thus when neon is added to the system in this problem, the rate is unchanged.

Interpret the following: The intensity of the deep purple color of an acid KMnO_4 solution can be used as an indicator of the extent to which oxidation has taken place. Often there is a considerable time lapse before any visual evidence of reaction is noted; but once begun, the process proceeds vigorously. If, on the other hand, a small amount of the essentially colorless MnCl_2 salt is dissolved in this solution, the process is immediately vigorous, no latency period being noted.

Solution: The presence of the Mn^{2+} in the solution is the key to the reaction. When MnCl_2 is dissolved in the solution, the reaction is vigorous since the salt immediately dissociates to produce Mn^{2+} ions.

KMnO_4 also immediately dissociates into K^+ and MnO_4^- . With KMnO_4 , however, a latency period is noticed since it takes time for Mn^{2+} to be produced from the reduction of the MnO_4^- ion.

The fact that once Mn^{2+} does exist, whether from MnO_4^- or MnCl_2 , the reaction proceeds vigorously suggests that Mn^{2+} is a catalyst. Thus, Mn^{2+} increases the rate of the reaction without being consumed. Because Mn^{2+} is also a reaction product, the reaction is autocatalytic.

HALF-LIFE

What is the half-life of an unstable substance if 75% of any given amount of the substance decomposes in one hour?

Solution: The half-life is defined as the time it takes for one half of the amount of a substance to decompose. When given the time elapsed and the percent of decomposition, one can find the half-life. One knows that $\frac{1}{2}$ of the substance decomposes in the time equal to the half-life. This leaves $\frac{1}{2}$ of the substance, $\frac{1}{2}$ of this decomposes in the next span of time elapsed equal to the half-time. This leaves $\frac{1}{4}$ of the substance. 75% of it has decomposed after two half-lives have elapsed. Thus 2 half-lives equal one hour or the half-life of the substance is $\frac{1}{2}$ hour.

For the first order decomposition of azomethane at 600°K , it takes 30 minutes for the original concentration to decrease to half its value. After 60.0 minutes have elapsed, what percentage of the azomethane originally present remains?

Solution: The decomposition is first order which means $\text{rate} = k[A]$, where k is a constant and $[A]$ is the concentration of azomethane. The half-life is independent of time, i.e., it is a constant and is defined as the time necessary for half the particular reactant present to decompose. Thus, if you knew the half-life of azomethane, you could calculate the percentage of azomethane that remains after a certain period of time. The half-life of azomethane is 30 minutes. Thus, after 30 minutes, you have half of the original material left. After 60 minutes half of this original material is left or $\frac{1}{4}$ of the original material remains.

In the reaction $\text{N}_2\text{O}_5 \rightarrow \text{N}_2\text{O}_4 + \frac{1}{2}\text{O}_2$, the N_2O_5 decomposes by a first-order mechanism. At 298°K , the half-life is 340 minutes. Find the value of the reaction rate constant. Calculate the number of minutes required for the reaction to proceed 70 percent towards completion.

Solution: Half-life may be defined as the time necessary for half the particular reactant present initially, in this case N_2O_5 , to decompose. For a first order reaction, the rate constant can be expressed in terms of half-life, $t_{\frac{1}{2}}$.

Namely, $t_{\frac{1}{2}} = \frac{.693}{k}$, where k is the rate constant. (Caution: This expression is only true for a first order reaction.) Solving this expression for k to obtain

$$k = \frac{.693}{t_{\frac{1}{2}}} = \frac{.693}{340} = 2.04 \times 10^{-3} \text{ min}^{-1}.$$

To find the amount of time required for the reaction to proceed to 70% completion, use the fact that

$$t = \frac{-2.303}{k} \log \frac{C_0}{C} \quad \text{in a first order reaction, where}$$

t = time, k = rate constant, C_0 = initial concentration and C = existing concentration.


Suppose the initial concentration is X , then at 70% completion, the existing concentration is $.30 X$. Having calculated k , substitute to find t = minutes.

$$t = \frac{-2.303}{k} \log \frac{C_0}{C} = \frac{-2.303}{2.04 \times 10^{-3}} \log \left(\frac{.30X}{X} \right)$$

$$= (1129)(.523) = 590 \text{ min.}$$

• PROBLEM 463

The half-life for the first order reaction:



is 80 minutes. In what period of time would the concentration of SO_2Cl_2 be reduced to 1.0% of the original?

Solution: If a reaction is first order, it means that the rate of the reaction is proportional to the concentration of a single reactant. For example, if substance A decomposes into the products B and C,



the rate is proportional to the concentration of A which is present at any time.

$$\text{Rate} = k [\text{A}],$$

where k is the rate constant for the reaction and [A] is the concentration of A.

The rate at which the reaction proceeds is equal to the decrease in concentration of A with change of time. This can be written in the integrated form

$$2.303 \log \frac{[\text{A}]}{[\text{A}_0]} = -kt,$$

where [A] is the concentration of A after time t has elapsed, $[\text{A}_0]$ is the original concentration of A and k is the rate constant for the reaction. In this problem, the final ratio of [A] to $[\text{A}_0]$ is given as 1.0% or 0.010. Therefore, this equation can be used to find the time it takes for this process to occur, once one has calculated the rate constant k. The rate constant can be found if one remembers the relationship between the half-life of a reaction and k. Namely,

$$t_{1/2} = \frac{0.693}{k},$$

where $t_{1/2}$ is the half life and k is the rate constant. The $t_{1/2}$ for this reaction is given as 8.0 minutes, therefore this relation can be used to obtain k.

$$k = \frac{0.693}{t_{1/2}}$$

$$k = \frac{0.693}{8.0 \text{ minutes}} = 0.087/\text{minute}$$

Once k is known, it can be used in the original rate equation to determine the time it takes for this reaction to proceed until the concentration of A is 1.0% that of the original concentration.

$$2.303 \log \frac{[A]}{[A_0]} = -kt. \quad \text{Substituting,}$$

$$2.303 \log 0.010 = -0.087/\text{minute} \times t$$

$$t = \frac{2.303 \log 0.010}{-0.087/\text{min}} = \frac{(2.303)(-2)}{-0.087/\text{min}}$$

$$= 53 \text{ minutes.}$$

• PROBLEM 464

At a certain temperature, the half-life for the decomposition of SO_2Cl_2 is 4.00 hours. (a) In what period of time would the concentration of SO_2Cl_2 be reduced to 10% of the original? (b) Starting with 100 mg of SO_2Cl_2 , what weight would be left at the end of 6.50 hours? The reaction is first order, therefore the equation $\ln[A]/[A_0] = -kt$ can be used. $[A]$ is the concentration of A at time t , $[A_0]$ is the original concentration of A and k is rate constant.

Solution: (a) When the concentration of A is 10% of the original, the ratio of $[A]/[A_0]$ equals .10. Therefore, one can use the equation given to solve for t after first determining k for the reaction. k is determined from the half-life in a first order reaction. The relation between k and $t_{1/2}$ (half-life) is

$$t_{1/2} = \frac{0.693}{k}$$

One is given the $t_{1/2}$. Solving for k :

$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{4.0 \text{ hrs}} = .173 \text{ hr}^{-1}$$

One can now solve for t :

$$\frac{\ln[A]}{[A_0]} = -kt$$

$$\ln .10 = - .173 \text{ hr}^{-1} t$$

$$t = \frac{\ln .10}{-.173 \text{ hr}^{-1}} = \frac{-2.303}{-.173 \text{ hr}^{-1}} = 13.3 \text{ hrs.}$$

(b) One can solve for the $[A]$ after 6.5 hours by use of the equation:

$$\frac{\ln[A]}{[A_0]} = -kt$$

One is given that there is originally 100 mg of A. $1 \text{ g} = 1000 \text{ mg}$, thus $100 \text{ mg} = 0.1 \text{ g}$. $[A_0]$ must be in moles/liter. One finds the number of moles of SO_2Cl_2 in 0.1 g by dividing 0.1 g by the molecular weight of SO_2Cl_2 (MW = 135).

$$[A_0] = \frac{\text{no. of moles}}{\text{liter}} = \frac{0.1 \text{ g}}{135 \text{ g/mole}} = 7.41 \times 10^{-4} \frac{\text{moles}}{\text{liter}}$$

Solving for $[A]$ using the value of k found in part a:

$$\ln \frac{[A]}{7.41 \times 10^{-4} \text{ moles/liter}} = -(.173 \text{ hr}^{-1})(6.5 \text{ hr})$$

$$\ln [A] - \ln [7.41 \times 10^{-4} \text{ moles}] = -(.173 \text{ hr}^{-1})(6.5 \text{ hr})$$

$$\ln [A] = - (1.12) + \ln (7.41 \times 10^{-4})$$

$$\ln [A] = - 1.12 - 7.21$$

$$\ln [A] = - 8.33$$


$$[A] = 2.41 \times 10^{-4} \text{ moles}$$

One can find the number of grams by multiplying the number of moles by the molecular weight.

$$\begin{aligned} \text{weight} &= 2.41 \times 10^{-4} \text{ moles} \times 135 \text{ g/mole} \\ &= .0326 \text{ g} = 32.6 \text{ mg} \end{aligned}$$

• PROBLEM 465

The ketone acid $(\text{CH}_3\text{CO}_2\text{H})_2\text{CO}$ undergoes a first-order decomposition in aqueous solution to yield acetone and carbon dioxide:



(a) Write the expression for the reaction rate. (b) The rate constant k has been determined experimentally as $5.48 \times 10^{-2}/\text{sec}$ at 60°C . Calculate $t_{1/2}$ at 60°C . (c) The rate constant at 0°C has been determined as $2.46 \times 10^{-3}/\text{sec}$. Calculate $t_{1/2}$ at 0°C . (d) Are the calculated half-lives in accord with the stated influence of temperature on reaction rate?

Solution: (a) For a chemical decomposition, the rate of the reaction is equal to the product of the rate constant (k) and the concentration of the compound decomposing. Thus,

$$\text{Rate} = k [(\text{CH}_3\text{CO}_2\text{H})_2\text{CO}]$$

(b) Because the rate is only proportional to $[(\text{CH}_3\text{CO}_2\text{H})_2\text{CO}]$, the reaction is first-order. For a first-order reaction, the half-life ($t_{1/2}$) is related to k by the following equation:

$$t_{1/2} = \frac{0.693}{k}$$

Solving for $t_{1/2}$:

$$t_{1/2} = \frac{0.693}{5.48 \times 10^{-2}/\text{sec}} = 12.65 \text{ sec.}$$

(c) One can solve for $t_{1/2}$ at 0°C using the same equation.

$$t_{1/2} = \frac{0.693}{2.46 \times 10^{-5}/\text{sec}} = 2.82 \times 10^4 \text{ sec.}$$

(d) In general, the speed of a chemical change is approximately doubled for each ten degrees rise in temperature. The temperature rises 60° , from 0°C to 60°C , therefore, the rate should double six times or the ratio of the $t_{1/2}$ at 0°C to the $t_{1/2}$ at 60°C is 2^6 . $2^6 = 64$.

$$\frac{t_{1/2}^{0^\circ}}{t_{1/2}^{60^\circ}} = \frac{2.82 \times 10^4 \text{ sec}}{12.65 \text{ sec}} = 2.23 \times 10^3$$

This is much greater than the expected ratio of 64.

• PROBLEM 466

When heated to 600°C , acetone (CH_3COCH_3) decomposes to give CO and various hydrocarbons. The reaction is found to be first order in acetone concentration with a half-life of 81 sec. Given at 600°C a 1-liter container into which acetone is injected at 0.48 atm, approximate how long would it take for the acetone pressure to drop to 0.45 atm?

Solution: When a reaction is said to be first order in a particular reactant it means that the rate of the reaction is proportional to the concentration of that reactant. Thus the rate law for this reaction can be written:

$$\text{Rate} = k [\text{CH}_3\text{COCH}_3]$$

where k is the rate constant and $[\]$ indicate concentration. The rate constant is related to the half-life ($t_{1/2}$) by the following equation:

$$t_{1/2} = \frac{0.693}{k}$$

$$\text{Solving for } k: \quad t_{1/2} = 81 \text{ sec}$$

$$81 \text{ sec} = \frac{0.693}{k}$$

$$k = 8.56 \times 10^{-3} \text{ sec}^{-1}$$

To solve for the time it takes for a reaction to proceed to a certain degree one uses the equation:

$$d \ln[A] = -k \, dt$$

where $d \ln[A]$ is the natural logarithm of the original concentration subtracted from the natural log of the final concentration of acetone, k is the rate constant and dt is the time elapsed. Because acetone is a gas in this experiment the pressure is proportional to the concentration. Therefore one can let $[A]_{\text{original}} = 0.48$ and $[A]_{\text{final}} = 0.45$. Solving for dt :

$$\begin{aligned} \ln 0.48 - \ln 0.45 &= (-8.56 \times 10^{-3} \text{ sec}^{-1})(dt) \\ (-.734) - (-.799) &= (-8.56 \times 10^{-3} \text{ sec}^{-1})(dt) \\ \frac{(-.734) - (-.799)}{-8.56 \times 10^{-3} \text{ sec}^{-1}} &= dt \\ -7.5 \text{ sec} &= dt \end{aligned}$$

Therefore 7.5 sec have elapsed since the beginning of the reaction.

THE ARRHENIUS EQUATION; RELATING TEMPERATURE AND REACTION RATE

• PROBLEM 467

Using a specific case, show that the effect of a 10°K rise in temperature will have a greater effect on the rate constant, k , at low temperatures than it does at high temperatures.

Solution: The best way to demonstrate this fact is to use the natural logarithm form of Arrhenius' equation,

which gives an indication of the effect of temperature on reaction rate. The expression may be written $\ln k = \ln A - E_a/2.303 RT$, where k is the rate constant, A = the Arrhenius constant, E_a = activation energy, R = universal gas constant ($8.314 \text{ J mol}^{-1} \text{ deg}^{-1}$) and T = temperature in Kelvin (Celsius plus 273). For a low temperature, consider room temperature ($T = 300^\circ\text{K}$). For a first-order reaction, A might be $1.0 \times 10^{14} \text{ sec}^{-1}$ and E might be 80 k J/mole . Thus, at $T = 300^\circ\text{K}$, you have $\log k = \log (1 \times 10^{14}) - 80,000/(2.303)(8.314)(300) = .07$. Thus, $k = 1.2 \text{ sec}^{-1}$. If you increase the temperature 10°K to 310°K , you find by exactly the same method of calculation, that $k = 3.3 \text{ sec}^{-1}$. Thus, at a temperature of 10°K higher (in a low temperature range), k was increased nearly three times.

Now consider a higher temperature range, say $T = 900^\circ\text{K}$. For this, $\log k = (1 \times 10^{14}) - 80,000/(2.303)(8.314)(900) = 9.358$. Solving $\log k = 9.358$, $k = 2.28 \times 10^9$. Again, let us increase the temperature by 10°K to 910°K . Using the same type of calculations, you find that k becomes 2.56×10^9 . The percent change =

$$\frac{2.56 \times 10^9 - 2.28 \times 10^9}{2.28 \times 10^9} \times 100 = 12.28\%$$

Thus, at the higher temperatures, the rate constant was increased only 12% as compared to 300% at the low temperatures.

• PROBLEM 468

What activation energy should a reaction have so that raising the temperature by 10°C at 0°C would triple the reaction rate?

Solution: The activation energy is related to the temperature by the Arrhenius equation which is stated

$$k = Ae^{-E/RT}$$

where A is a constant characteristic of the reaction; e is the base of natural logarithms, E is the activation energy, R is the gas constant ($8.314 \text{ J mol}^{-1} \text{ deg}^{-1}$) and T is the absolute temperature. Taking the natural log of each side:

$$\ln k = \ln A - E/RT$$

For a reaction that is 3 times as fast, the Arrhenius equation becomes

$$3k = Ae^{-E/R(T + 10^\circ)}$$

Taking the natural log:

$$\ln 3 + \ln k = \ln A - E/R(T + 10^\circ)$$

Subtracting the equation for the final state from the equation for the initial state:

$$\begin{array}{r} \ln k = \ln A - E/RT \\ - (\ln 3 + \ln k = \ln A - E/R(T + 10)) \\ \hline - \ln 3 = -E/RT + E/R(T + 10) \end{array}$$

Solving for E:

$$-\ln 3 = -E/RT + E/R(T + 10) \quad \begin{array}{l} R = 8.314 \text{ J/mole } ^\circ\text{K} \\ T = 0 + 273 = 273 \end{array}$$

$$- \ln 3 = -E/(8.314 \text{ J/mole-K})(273\text{K}) + E/(8.314 \text{ J/mole-K})(283\text{K})$$

$$-1.10 = -E/2269.72 \text{ J/mole} + E/2352.86 \text{ J/mole}$$

$$(2269.72 \text{ J/mole})(2352.86 \text{ J/mole}) \times -1.10 =$$

$$(-E/2269.72 \text{ J/mole} + E/2352.86 \text{ J/mole})(2269.72 \text{ J/mole})(2352.86 \text{ J/mole})$$

$$-5.874 \times 10^6 \text{ J}^2/\text{mole}^2 = (-E)(2352.86 \text{ J/mole}) + E(2269.72 \text{ J/mole})$$

$$-5.874 \times 10^6 \text{ J}^2/\text{mole}^2 = -8.314 \times 10^1 \text{ J/mole} \times E$$

$$7.06 \times 10^4 \text{ J/mole} = E$$

● PROBLEM 469

It has been suggested that a 10° rise in temperature results in a twofold response in the rate of a chemical reaction. What is the implied activation energy at 27°C ? at 1000°C ?

Solution: The activation energy, E_A , is the difference between the heat concentration of the active molecules and the inert molecules. The activation energy can be related to the rate constants of the reaction at two different temperatures by one form of the Arrhenius equation:

$$(i) \quad \log \frac{k_2}{k_1} = - \frac{E_A}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

(where R = universal gas constant; k_1 is the rate constant at temperature T_1 ; k_2 is the rate constant at T_2).

This equation can be rewritten as

$$(ii) \quad E_a = \frac{2.303 R}{\frac{1}{T_1} - \frac{1}{T_2}} \log \frac{k_2}{k_1}$$

Thus to solve for E_a , you find T_1 , T_2 , and k_2/k_1 and substitute. Let T_1 = initial temperature. If there is a 10° rise in temperature, then $T_2 = T_1 + 10^\circ\text{K}$. You are told that when the temperature is increased, the original rate is doubled. Thus, if k_1 = original rate, $k_2 = 2k_1$. Thus, $\log k_2/k_1 = \log 2 = .301$. For the case, $T_1 = 27^\circ\text{C} = 300^\circ\text{K}$, $T_2 = 310^\circ\text{K}$ you have

$$E_a = \frac{-2.303 R}{\left(\frac{1}{T_2} - \frac{1}{T_1}\right)} \log \frac{k_2}{k_1} = \frac{-2.303 (1.987) \left(\frac{1 \text{ k cal}}{1000 \text{ cal}}\right)}{\left(\frac{1}{310} - \frac{1}{300}\right)} \quad (.301)$$

$$= 12.8 \text{ k cal/mole}$$

For the case with $T_1 = 1000^\circ\text{C} = 1273^\circ\text{K}$, $T_2 = 1283^\circ\text{K}$. You have,

$$E_a = \frac{-2.303 (1.987) (1 \text{ k cal}/1000 \text{ cal})}{\left(\frac{1}{1283} - \frac{1}{1273}\right)} \quad (.301)$$

$$= 226 \text{ k cal/mole}$$

(In both cases, $1 \text{ k cal}/1000 \text{ cal}$ is a conversion factor.)

• PROBLEM 470

(1) A reaction proceeds five times as fast at 60°C as it does at 30°C . Estimate its energy of activation. (2) For a gas phase reaction with $E_a = 40,000 \text{ cal/mole}$, estimate the change in rate constant due to a temperature change from 1000°C to 2000°C .

Solution: The activation energy E_a can be related to the rate constants k_1 (at temperature T_1) and k_2 (at temperature T_2) by the Arrhenius equation:

$$\log \frac{k_2}{k_1} = - \frac{E_a}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right),$$

where R = universal gas constant.

(1) You are told a reaction proceeds five times as fast at 60°C as it does at 30°C . Therefore, if k_1 = rate constant at $30^\circ\text{C} = 303^\circ\text{K}$ with $T_1 = 303^\circ\text{K}$, then $k_2 = 5k_1$ at $60^\circ\text{C} = 333^\circ\text{K}$ with $T_2 = 333^\circ\text{K}$. You are given R . Substitute these values into the Arrhenius equation, and

solve for E_a . Rewriting and substituting,

$$E_a = \frac{-2.303 R}{\frac{1}{T_2} - \frac{1}{T_1}} \log \frac{k_2}{k_1} = \frac{(-2.303)(1.987) \left(\frac{1 \text{ k cal}}{1000 \text{ cal}} \right) \log 5}{\left(\frac{1}{333} - \frac{1}{303} \right)}$$

$$= (15.4 \text{ k cal/mole})(.699) = 10.8 \text{ k cal/mole.}$$

[Note: 1 k cal/1000 cal is a conversion factor to obtain the correct units.] To answer (2) find k_2/k_1 from the Arrhenius equation. Rewriting and substituting,

$$\frac{k_2}{k_1} = \text{antilog} \left[\frac{E_a}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right) \right]$$

$$= \text{antilog} \left[\frac{-40000}{(2.303)(1.987) \left(\frac{1}{2273} - \frac{1}{1273} \right)} \right]$$

$$= \text{antilog } 3.02 = 1.05 \times 10^3.$$

That is, the rate should be about 1050 times as great at 2000°C as at 1000°C.

• PROBLEM 471

For the gas-phase decomposition of acetaldehyde, $\text{CH}_3\text{CHO} \rightarrow \text{CH}_4 + \text{CO}$, the second-order rate constant changes from $0.105 \text{ m}^{-1} \text{ sec}^{-1}$ at 759°K to $0.343 \text{ m}^{-1} \text{ sec}^{-1}$ at 791°K . Calculate the activation energy that this corresponds to. What rate constant would you predict for 836°K ?

Solution: The rate constant is related to the activation energy by the Arrhenius equation, which is stated

$$k = Ae^{-E/RT}$$

where k is the rate constant, A is a constant characteristic to the reaction, e is the base of natural logarithms, E is the activation energy, R is the gas constant ($8.314 \text{ J/mole } ^\circ\text{K}$) natural log:

$$\ln k = \ln A - E/RT$$

One is given k , R and T in the problem for two trials.

$$\text{For } 759^\circ\text{K:} \quad k = 0.105 \text{ m}^{-1} \text{ sec}^{-1}$$

$$\text{a) } \ln(0.105) = \ln A - E/(8.314 \text{ J/mole } ^\circ\text{K})(759^\circ\text{K})$$

$$\text{For } 791^\circ\text{K:} \quad k = 0.343 \text{ m}^{-1} \text{ sec}^{-1}$$

$$\text{b) } \ln(0.343) = \ln A - E/(8.314 \text{ J/mole } ^\circ\text{K})(791^\circ\text{K})$$

Subtract equation b from equation a to solve for E.

$$\begin{aligned}
 \ln(0.105) &= \ln A - E/(8.314 \text{ J/mole}^\circ\text{K})(759^\circ\text{K}) \\
 - [\ln(0.343) &= \ln A - E/(8.314 \text{ J/mole}^\circ\text{K})(791^\circ\text{K})] \\
 \text{c) } \ln(0.105) - \ln(0.343) &= -E/(8.314 \text{ J/mole}^\circ\text{K})(759^\circ\text{K}) + \\
 &\quad E/(8.314 \text{ J/mole}^\circ\text{K})(791^\circ\text{K}) \\
 - 2.25 - (-1.07) &= -E/6310.33 \text{ J/mole} + E/6576.37 \text{ J/mole} \\
 (6310.33 \text{ J/mole})(6576.37 \text{ J/mole})(-1.18) &= E/6310.33 \text{ J/mole} \\
 + E/6576.37 \text{ J/mole}(6310.33 \text{ J/mole})(6576.37 \text{ J/mole}) \\
 - 4.8969 \times 10^7 \text{ J}^2/\text{mole}^2 &= -6576.37 E \text{ J/mole} \\
 &\quad + 6310.33 E \text{ J/mole} \\
 - 4.8969 \times 10^7 \text{ J}^2/\text{mole}^2 &= -2.6604 \times 10^2 E \text{ J/mole} \\
 E &= \frac{-4.8969 \times 10^7 \text{ J}^2/\text{mole}^2}{-2.6604 \times 10^2 \text{ J/mole}} \\
 E &= 1.8407 \times 10^5 \text{ J/mole}
 \end{aligned}$$

One can solve for k at 836°K by replacing the values of this third trial for the values found at 791°K.

$$\begin{aligned}
 \ln(0.105) - \ln k &= - \frac{1.8407 \times 10^5 \text{ J/mole}}{(8.314 \text{ J/mole}^\circ\text{K})(759^\circ\text{K})} \\
 &\quad + \frac{1.8407 \times 10^5 \text{ J/mole}}{(8.314 \text{ J/mole}^\circ\text{K})(836^\circ\text{K})} \\
 - 2.25 - \ln k &= -2.917 \times 10^1 + 2.648 \times 10^1 \\
 - \ln k &= -4.4 \times 10^{-1} \\
 k &= 1.55 \text{ m}^{-1} \text{ sec}^{-1}.
 \end{aligned}$$

• PROBLEM 472

A chemist found the value of the specific rate constant for the decomposition of nitrous oxide, $2\text{NO} + \text{N}_2 + \text{O}_2$, at two separate temperatures: $k_1 = .14 \text{ liter/mole-sec}$ at 970°K , $k_2 = 3.7 \text{ liter/mole-sec}$ at 1085°K . (a) Calculate the activation energy, E_a , for the reaction, (b) Calculate the A factor in the Arrhenius equation, (c) Calculate the specific rate constant at 800°K .

Solution: This problem deals with the effect of temperature on the kinetics of a reaction.

(a) E_a is the activation energy, the difference between the heat content of the active molecules and that of the inert molecules. To find the value of E_a for the decomposition of nitrous oxide, use the Arrhenius equation:

$$E_a = \frac{-2.303 R}{\left(\frac{1}{T_2} - \frac{1}{T_1}\right)} \log \left(\frac{k_2}{k_1}\right)$$

where T = temperature in degrees Kelvin (Celsius plus 273°), k = specific rate constant and R = universal gas constant. Substitute known values to obtain ($T_1 = 970^\circ$, $T_2 = 1085^\circ$)

$$E_a = \frac{(-2.303)(1.987)(1/1000)}{(1/1085 - 1/970)} \log \left(\frac{3.7}{.14}\right)$$

$$= 59.2 \text{ k cal/mole}$$

[Note: 1/1000 is a conversion factor of 1 k cal/1000 cal.]

(b) The Arrhenius equation can also be written in the form

$$\log k = \log A - \frac{E_a}{RT}, \quad \text{where } A \text{ is called the frequency}$$

factor. To find A , first solve the equation in terms of A .

$$\log A = \log k + \frac{E_a}{RT} \quad \text{or} \quad \log A = \log k + \frac{E_a}{2.303 RT}$$

Substitute for k and T and calculate

$$\log A = \log 3.7 + \frac{(59.2) \left(\frac{1000 \text{ cal}}{1 \text{ k cal}}\right)}{(2.303)(1.987)(1085)} = 0.57 + 11.9 = 12.5.$$

Since $\text{antilog } 12.5 = A$, $A = 3 \times 10^{12}$ liters/mole-sec. (Notice: k_2 and T_2 values were used in the calculations but k_1 and T_1 could have been used instead.)

(c) To find k at 800 solve the Arrhenius equation for k

$$\log k = \log A - \frac{E_a}{2.303 RT}$$

Substitute in known values for A , E_a , R , and T to obtain:

$$\log (3 \times 10^{12}) - \frac{(59.2) \left(\frac{1000 \text{ cal}}{1 \text{ k cal}}\right)}{2.303 (1.987)(800)} = 12.5 - 16.2 = -3.7$$


$$k = \text{antilog } (-3.7) = 2 \times 10^{-4} \text{ liters/mole-sec.}$$

CHAIN REACTIONS

• PROBLEM 473

Define the term "chain reaction". Using a specific example, distinguish between a chain-starting, a chain-propagating, and a chain-terminating step. Discuss the energy absorbed or released in the first and last of these.

Solution: In chemical kinetics, a chain reaction occurs when an intermediate species that is consumed in one step is regenerated in a later step. As a result, there is a sequence of steps which endlessly repeat themselves, like links in a chain, until the chain is ended or the starting material exhausted. A specific example of a chain reaction, is the formation of methylchloride from methane (CH_4) and chlorine gas (Cl_2). Overall, the reaction is:




The mechanism of this chain reaction is as follows:


- (1) $\text{Cl}_2 \rightarrow 2\text{Cl}\cdot$
- (2) $\text{CH}_4 + \text{Cl}\cdot \rightarrow \text{CH}_3\cdot + \text{HCl}$
- (3) $\text{CH}_3\cdot + \text{Cl}_2 \rightarrow \text{CH}_3\text{Cl} + \text{Cl}\cdot$

A chain-starting step encompasses one or more stable molecules and produces highly reactive species, such as single atoms. In this example, (1) fits this description. A stable Cl_2 molecule forms single chlorine radicals ($\text{Cl}\cdot$) that immediately react in step (2). Cl_2 does not go to $\text{Cl}\cdot$ spontaneously. Energy must be supplied in the form of heat or light. Chain-starting steps generally require energy. A chain-propagating step involves the reaction between a molecule and one of the highly reactive species, producing at least one highly reactive species. Reactions (2) and (3) fit this description. In (2), the reactive species $\text{CH}_3\cdot$, methyl radical, is produced via the reaction between the molecule CH_4 and reactive species $\text{Cl}\cdot$. In (3), this species, i.e., $\text{CH}_3\cdot$ reacts with Cl_2 to produce another reactive species, chlorine radical ($\text{Cl}\cdot$). Note: this chlorine radical can go back to react with CH_4 to repeat the entire process, i.e., (2) and (3) over and over again, until you have the chain-terminating step. Chain termination occurs when one of the highly reactive species reacts with another highly reactive species, or with the wall and no energetic species result. In our example, such a step would be the reaction of two chlorine radicals to produce chlorine gas, $\text{Cl}\cdot + \text{Cl}\cdot \rightarrow \text{Cl}_2$. Since Cl_2 is not a reactive species, the chain is terminated. Notice that this is the reverse of $\text{Cl}_2 \rightarrow 2\text{Cl}\cdot$, which required energy. Thus, a chain-terminating step will generally release energy.

Given the consecutive reaction $A \xrightarrow{k_1} B \xrightarrow{k_2} C$ with $k_1 = k_2$. Draw a graph for the time variation of the concentrations of A, B, and C.



Solution: To draw this graph, you need to determine the concentration with respect to time of each species in this reaction. This will necessitate writing rate law expressions for each of the species. The rate law states that rate is equal to a proportionality constant k times the concentration of one species or more than one species, each raised to a power. The power is equal to the number of moles reacting. You are told that



The rate at which A disappears can be written as $-\frac{d[A]}{dt} = k_1[A]$. A will disappear at a certain rate as given by the rate constant. How long it takes to disappear will reflect how much was initially present: the concentration. The rate at which C appears may be written as $\frac{d[C]}{dt} = k_2[B]$. C forms from B at a certain rate as indicated by k_2 . Thus, the amount of B initially present determines the time it takes for C to appear. What about the concentration of B? It increases with time as A disappears, yet decreases as C appears. The net increase in B will be the differences in these processes. Thus, $\frac{d[B]}{dt} = k_1[A] - k_2[B]$. If you start with pure A at $t = 0$,

$\frac{d[B]}{dt}$ is positive, since $k_1[A] \gg k_2[B]$. After a time, however, A is used up, so that $\frac{d[B]}{dt}$ becomes negative. This means the concentration of B will peak sometime along in the reaction. At that instant,

$$\frac{d[B]}{dt} = 0, \text{ so that } k_1[A] - k_2[B] = 0, \text{ which means } \frac{[A]}{[B]} = \frac{k_2}{k_1}.$$

Thus, the behavior of the system depends on k_2/k_1 . As k_1 grows small, B becomes less notable. But, you are told that $k_2 = k_1$, which indicates that $k_2/k_1 = 1$.




[C] will rise as [B] falls. Thus, you have the following graph.

If all reactions proceed to completion C will be the only species present at the conclusion of the process.

• PROBLEM 475

N_2O_5 decomposes according to the following equation:
 $N_2O_5 + 2NO_2 + \frac{1}{2}O_2$. The rate expression =

$-\frac{d[N_2O_5]}{dt} = k[N_2O_5]$. The following mechanism has been proposed:



Show the rate of O_2 formation is directly proportional to $[N_2O_5]$.

Solution: To relate O_2 formation with the concentration of N_2O_5 , investigate the mechanism of this reaction. In the mechanism, oxygen (O_2) is formed only from the second reaction, the rate being

$$\frac{d[O_2]}{dt} = k_1[NO_2][NO_3]. \text{ To relate } \frac{d[O_2]}{dt} \text{ to } [N_2O_5],$$

express $[NO_2][NO_3]$ in terms of $[N_2O_5]$.

This can be done from the first reaction, which is at equilibrium. The equilibrium constant of this reaction, k , measures the ratio of products to reactants, each raised to the power of its coefficient in the chemical equation. Thus,

$$k_{eq} = \frac{[NO_2][NO_3]}{[N_2O_5]} \quad \text{or} \quad [NO_2][NO_3] = k_{eq} [N_2O_5]$$

Recall, $\frac{d[\text{O}_2]}{dt} = k_1 [\text{NO}_2][\text{NO}_3]$. Thus $[\text{NO}_2][\text{NO}_3]$ can be replaced by $k_{\text{eq}} [\text{N}_2\text{O}_5]$, so that

$\frac{d[\text{O}_2]}{dt} = k_1 k_{\text{eq}} [\text{N}_2\text{O}_5]$, which means O_2 formation is directly proportional to $[\text{N}_2\text{O}_5]$.

• PROBLEM 476

You have the reaction $\text{HCrO}_4^- + 3\text{Fe}^{2+} + 7\text{H}^+ \rightarrow \text{Cr}^{3+} + 3\text{Fe}^{3+} + 4\text{H}_2\text{O}$ with the rate law expression: $\text{rate} = k [\text{HCrO}_4^-]$

$[\text{Fe}^{2+}][\text{H}^+]$. Explain why the rate of reaction is not proportional to the number of each species reacting according to the stoichiometry of the balanced equation.

Solution: Stoichiometry predicts only the amount of products expected; it cannot predict when they will be produced. The rate of production is dependent only on the number of molecules in the activated complex and their concentrations in the system. A reaction can occur only when the species come in contact with each other. Thus, the concentration reflects the probability of a collision among the species. Stoichiometry cannot provide any information in these areas, and, as such, the rate of the reaction is not necessarily reflective of the stoichiometry of the balanced equation.


• PROBLEM 477

The rate law for the reaction $\text{H}_2(\text{g}) + \text{Br}_2(\text{g}) \rightarrow 2\text{HBr}(\text{g})$ is

$$\frac{d[\text{HBr}]}{dt} = \frac{k[\text{H}_2][\text{Br}_2]^{1/2}}{1 + k' [\text{HBr}]/[\text{H}_2]} \quad . \quad \text{Derive this law from}$$

the reaction mechanism, which consists of chain-starting, chain-propagating, and chain-terminating steps. Use steady-state approximation.

Solution: A chain-reaction comes about when an intermediate species that is consumed in one step is regenerated in a later step. As a consequence of this, a sequence of steps is set up such that they endlessly repeat themselves until the chain is terminated or the starting material is used up. By various experiments and observations, the mechanism has been found to be the following:



To derive the rate law for this reaction, the formation of HBr, use these individual reactions:

Notice, first, that only those reactions that dealt with HBr in some way were included. In other words, (1) + (5) did not have this molecule, so that they cannot be included. The whole idea of writing the rate in this fashion comes from the fact that when, say $\text{A} + \text{B} \rightarrow \text{products}$,

$\text{rate} = -\frac{d[\text{A}]}{dt} = -\frac{d[\text{B}]}{dt} = k[\text{A}][\text{B}]$. This is called the expression for the rate law of a second order reaction. By order, you mean the number of atoms or molecules whose concentrations directly determine the reaction rate. In other words, for (2), $d[\text{HBr}]/dt = k_2[\text{Br}\cdot][\text{H}_2]$. You can write similar expressions for (3) and (4). The overall rate of HBr formation is the sum of the rates of formation of the HBr in each of these reactions. In (2) and (3), you are making HBr, so that their rate law expressions are added. In (4), however, HBr is being eliminated, so that you subtract. To repeat, you now have

$$\frac{d[\text{HBr}]}{dt} = k_2[\text{Br}\cdot][\text{H}_2] + k_3[\text{H}\cdot][\text{Br}_2] - k_4[\text{H}\cdot][\text{HBr}]$$

To obtain the desired formula, you need to find $[\text{Br}\cdot]$ and $[\text{H}\cdot]$. You can do this by writing overall rate expressions for the formation of $[\text{H}\cdot]$ and $[\text{Br}\cdot]$. To do this, use the same reasoning employed for $[\text{HBr}]$. You find that

$$\frac{d[\text{H}\cdot]}{dt} = k_2[\text{Br}\cdot][\text{H}_2] - k_3[\text{H}\cdot][\text{Br}_2] - k_4[\text{H}\cdot][\text{HBr}]$$

Note: Only (2), (3), and (4) dealt with H, which explains why only they are used. You subtract both (3) and (4) from (2), since they consume H instead of producing it as (2) does. For Br, you find

$$\frac{d[\text{Br}\cdot]}{dt} = 2k_1[\text{Br}_2] - k_2[\text{Br}\cdot][\text{H}_2] + k_3[\text{H}\cdot][\text{Br}_2] -$$

$$k_4[\text{H}\cdot][\text{HBr}] - 2k_5[\text{Br}\cdot]^2.$$

Again, use the same type of reasoning and technique to obtain this expression. You are told steady-state conditions exist. This means that the concentrations of the reactive intermediates that are regenerated do not change with time. This means, therefore that $d[\text{Br}\cdot]/dt = 0$ and $d[\text{H}\cdot]/dt = 0$, since each is being regenerated. Thus, you can equate the above expressions with zero. You obtain

$$\frac{d[\text{Br}]}{dt} = 0 = 2k_1[\text{Br}_2] - k_2[\text{Br}][\text{H}_2] + k_3[\text{H}][\text{Br}] + k_4[\text{H}][\text{HBr}] - 2k_5[\text{Br}]^2.$$

$$\frac{d[\text{H}]}{dt} = 0 = k_2[\text{Br}][\text{H}_2] - k_3[\text{H}][\text{Br}_2] - k_4[\text{H}][\text{HBr}].$$

These two equations can be solved simultaneously to find $[\text{H}]$ and $[\text{Br}]$. If you perform this operation, you find $[\text{Br}] = [k_1/k_5[\text{Br}]]^{1/2}$ and

$$[\text{H}] = k_2 \frac{(k_1/k_5)^{1/2} [\text{H}_2][\text{Br}_2]^{1/2}}{k_3[\text{Br}_2] + k_4[\text{HBr}]}.$$

Recalling, now, that

$$\frac{d[\text{HBr}]}{dt} = k_2[\text{Br}][\text{H}_2] + k_3[\text{H}][\text{Br}_2] - k_4[\text{H}][\text{HBr}],$$

you can

substitute these values for $[\text{H}]$ and $[\text{Br}]$ to obtain

$$\frac{d[\text{HBr}]}{dt} = \frac{2 k_3 k_2 k_4^{-1} k_1^{1/2} k_5^{-1/2} [\text{H}_2][\text{Br}_2]^{1/2}}{k_3 k_4^{-1} + [\text{HBr}][\text{Br}_2]^{-1}}.$$

Now, if you

let $k = 2k_2 k_1^{1/2} k_5^{-1/2}$ and $k' = \frac{1}{k_3 k_4^{-1}}$, you obtain

$$\frac{d[\text{HBr}]}{dt} = \frac{k[\text{H}_2][\text{Br}_2]^{1/2}}{1 + k' [\text{HBr}]/[\text{H}_2]}.$$

CHAPTER 14

THERMODYNAMICS I

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 459 to 502 for step-by-step solutions to problems.

The key to all of the problems in this chapter is an understanding of the First Law of Thermodynamics. The First Law says, simply, that energy can neither be created nor destroyed. Therefore, in any reaction or process, the energy must balance. This is true of bond energies in chemical reactions, of energy changes associated with chemical reactions, and, with allowances, for changes between mass and energy (i.e., $E = mc^2$) in nuclear reactions.

In chemical reactions, there are two energies that are very common and enter into almost all energy calculations associated with reactions. The first is the internal energy, E . This is the bond energy associated with chemical bonds plus the kinetic energy of molecular species. The second is the PV energy. Since so many reactions take place at conditions that cause the system to either expand or contract against the atmospheric surroundings, the effect of PV work on the atmosphere becomes important. The enthalpy is the thermodynamic quantity defined to include both the internal energy and the PV work against the atmosphere.

$$H = E + PV$$

14-1

where H is the enthalpy, E the internal energy, and PV the pressure and volume terms. For condensed phases (i.e., liquids and solids) the volume change is often negligible and $H \approx E$, but for gases the difference is significant. Only changes in enthalpy and internal energy have physical significance; there is no absolute zero point for these quantities. Therefore, tabulation of enthalpy is the difference between the state for which it is tabulated and some reference state either implied or specified.

Important definitions used in the calculations in this chapter follow.

ΔH°_r = enthalpy change in a reaction (typically joules/mole)

ΔE°_r = internal energy change in a reaction (typically joules/mole)

ΔH°_f = enthalpy change for formation of a species from the elements
(joules/mole)

C_p = heat capacity at constant pressure (typically joules/mole/°C)

C_v = heat capacity at constant volume (typically joules/mole/°C)

The superscript $^\circ$ indicates that the value is at some defined standard state. Typically for gases, this is one atmosphere pressure at the specified temperature; for liquids and solids it is the pure species at the specified temperature. If temperature is not specified, it is usually 25°C or 298K.

Since for any balanced reaction, the same number of moles of each element exist in the products and reactants:

$$\Delta H^\circ_r = \sum (\Delta H^\circ_f)_{\text{products}} - \sum (\Delta H^\circ_f)_{\text{reactants}} \quad 14-2$$

where ΔH°_f is the enthalpy associated with forming a species from the elements. It is simply the value of ΔH°_r where all the reactants are elements. Extensive tables of the values of ΔH°_f are found in handbooks. By convention, ΔH is positive for endothermic changes (changes which consume energy) and negative for exothermic changes (changes which produce energy).

Similarly, the First Law of Thermodynamics makes it possible to calculate the energy change associated with a reaction from bond energies. The energy, usually the enthalpy, is associated with a reaction.

$$\Delta H^\circ_r = \sum (\text{bond energies})_{\text{products}} - \sum (\text{bond energies})_{\text{reactants}} \quad 14-3$$

Heat capacities are defined as the energy change required to change the temperature. Again, for liquids and solids, there is little difference in the heat capacity at constant pressure or constant volume and often it is not specified. However, gases expand appreciably when heated and the energy to push back the atmosphere ($P\Delta V$) is significant. For gases, therefore, it is important to specify whether the heat capacity applies for a constant pressure or constant volume process. For ideal gases there is a simple relationship between C_p and C_v .

$$C_p - C_v = nR \text{ (the universal gas constant)} \quad 14-4$$


Heats of fusion and vaporization are heat changes associated with the solid-liquid phase change and the liquid-vapor phase change, respectively. The solid to liquid phase change and the liquid to vapor phase change are always endothermic — i.e., energy must be supplied to cause the phase change to occur. Energy is required to break physical bonds between molecules. If the phase change proceeds in the opposite direction, as in condensation or freezing, the numerical value of the associated energy change is the same, but the sign is opposite.

$$2 \times \Delta H_f^\circ \text{ OH} = 51 \text{ Kcal/mole} + \Delta H_f^\circ \text{ HO-OH}$$

$$\Delta H_f^\circ \text{ OH} = \frac{51 \text{ Kcal/mole} - 32.58 \text{ Kcal/mole}}{2}$$

$$\Delta H_f^\circ \text{ OH} = \frac{18.42 \text{ Kcal/mole}}{2} = 9.21 \text{ Kcal/mole.}$$

(b) CH_3OH dissociates by breaking the C-O bond:



Therefore, the bond energy of the C-O bond equals $\Delta H_{\text{diss}}^\circ$. $\Delta H_{\text{diss}}^\circ$ equals the sum of the ΔH 's of the products minus the ΔH 's of the reactants. Thus,

$$\text{bond energy of C-O} = \Delta H_f^\circ \text{ of } \text{CH}_3^\dagger\text{(g)} + \Delta H_f^\circ \text{ of } \text{OH}^-\text{(g)} - \Delta H_f^\circ \text{ of } \text{CH}_3\text{OH(g)}.$$

$$\begin{aligned} \text{bond energy of C-O} &= 34.0 \text{ Kcal/mole} + 9.21 \text{ Kcal/mole} \\ &\quad + 47.96 \text{ Kcal/mole} \\ &= 91.17 \text{ Kcal/mole.} \end{aligned}$$

• PROBLEM 481

Using the following table of bond energies, calculate the energy change in the following:

- (a) $2\text{H}_2\text{(g)} + \text{O}_2\text{(g)} \rightarrow 2\text{H}_2\text{O(g)}$
- (b) $\text{CH}_4\text{(g)} + 2\text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)} + 2\text{H}_2\text{O(g)}$
- (c) $\text{CH}_4\text{(g)} + \text{Cl}_2\text{(g)} \rightarrow \text{CH}_3\text{Cl(g)} + \text{HCl(g)}$
- (d) $\text{C}_2\text{H}_6\text{(g)} + \text{Cl}_2\text{(g)} \rightarrow \text{C}_2\text{H}_5\text{Cl(g)} + \text{HCl(g)}$.

Bond Energies (in kcal per mole)

H-H	104	C-O	83
H-F	135	C=O	178
H-Cl	103	C-Cl	79
H-Br	88	C-F	105
H-I	71	Si-O	106
Li-H	58	Si-F	136
Cl-Cl	58	C-C	83
C-H	87	C=C	146
O-H	111	C≡C	199
O=O	118	N≡N	225
P-Cl	78		

Solution: When using the bond energies to calculate the net energy change in a reaction, the net energy change is equal to the total bond energy of the bonds formed subtracted from the bond energy of bonds broken. This method is illustrated in the following examples.

• PROBLEM 484

How many calories of heat must be added to 3.0 liters of water to raise the temperature of the water from 20°C to 80°C?

Solution: A calorie is defined as the amount of heat necessary to raise the temperature of one gram of water one degree Centigrade. One can find the mass of 3.0 liters of water by using the density. The density of water is 1 g/ml. 1 liter = 1000 ml. Therefore, 3.0 liters = 3000 ml.

$$\text{Mass} = \text{density} \times \text{volume}$$

$$\text{Mass} = 1 \text{ g/ml} \times 3000 \text{ ml} = 3000 \text{ g}$$

The temperature change is found by subtracting 20 from 80, which is 60°C.

One can find the calories necessary to raise 3000 g of water 60°C by multiplying 1 cal/g°C by 3000 g by 60°C.

$$\begin{aligned} \text{No. of calories} &= 1 \text{ cal/g}^\circ\text{C} \times 3000 \text{ g} \times 60^\circ\text{C} \\ &= 180000 \text{ cal.} = 180 \text{ Kcal.} \end{aligned}$$

• PROBLEM 485

Calculate the quantity of heat required to raise the temperature of one gallon of water (3.78 liters) from 10° to 80°C.

Solution: The unit of heat is called the calorie. It is defined as the quantity of heat necessary to raise the temperature of one gram of water one degree centigrade. In this problem, one is told that 3.78 liters of water are heated from 10°C to 80°C. If one subtracts the final temperature from the original temperature, one can find the number of degrees the temperature of the water is raised.

$$\text{number of } ^\circ\text{C the temperature is raised} =$$

$$\text{final temperature} - \text{original temperature}$$

$$\text{no. of } ^\circ\text{C the temperature is raised} = 80^\circ\text{C} - 10^\circ\text{C} = 70^\circ\text{C.}$$

One must now determine the number of grams in 3.78 liters of water. Because 1 milliliter of water weighs one gram, one must determine the number of milliliters in 3.78 liters of water. There are 1000 milliliters in 1 liter; therefore, liters can be converted to milliliters by multiplying the number of liters present by the factor 1000 ml/1 liter.

$$3.78 \text{ liters} \times 1000 \text{ ml/l liter} = 3780 \text{ ml}$$

Because number of ml = number of grams for water, there are 3780 g of water present in 3.78 liters. One now knows that 3780 g of water have been raised 70°C in temperature. Remembering the definition for calorie, one can find the number of calories absorbed in this process by multiplying the number of grams of water by the number of degrees the temperature was raised by the factor 1 calorie/lg - 1°C, the specific heat of water.

$$\begin{aligned} \text{number of calories absorbed} &= 3780 \text{ g} \times 70^\circ\text{C} \times \\ &\quad 1 \text{ calorie/lg} - 1^\circ\text{C} \\ &= 265,000 \text{ calories} \end{aligned}$$

There are 1000 calories in 1 kilocalorie, so that calories can be converted to kilocalories by multiplying the number of calories by the factor 1 Kcal/1000 cal.

$$265,000 \text{ cal} \times 1 \text{ Kcal/1000 cal} = 265 \text{ Kcal.}$$

265 Kcal are absorbed in this process.

• PROBLEM 486

A piece of iron weighing 20.0 g at a temperature of 95.0°C was placed in 100.0 g of water at 25.0°C. Assuming that no heat is lost to the surroundings, what is the resulting temperature of the iron and water? Specific heats: iron = .108 cal/g-°C; water = 1.0 cal/g-°C.

Solution: The heat lost by the iron must be equal to the heat gained by the water. One solves for the heat lost by the iron by multiplying the number of grams of Fe by the number of degrees the temperature dropped by the specific heat of iron. The specific heat of a substance is defined as the amount of heat energy required to raise the temperature of 1 g of a substance by 1°C. The specific heat for iron is .108 cal/g-°C. Let t = the final temperature of the system.

$$\begin{aligned} \text{amount of heat lost by the iron} &= .108 \text{ cal/g-}^\circ\text{C} \times 20.0 \text{ g} \\ &\quad \times (95.0^\circ\text{C} - t) \end{aligned}$$

The amount of heat gained by the water is the specific heat of water multiplied by the weight of the water multiplied by the rise in the temperature. The specific heat of water is 1.0 cal/g-°C. Let t = final temperature of the system.

$$\begin{aligned} \text{amount of heat gained by water} &= 1.0 \text{ cal/g-}^\circ\text{C} \times 100 \text{ g} \\ &\quad \times (t - 25^\circ\text{C}) \end{aligned}$$

Solving for t:

amount of heat lost by the iron = amount of heat gained by the water. Therefore,

$$(.108 \text{ cal/g}^\circ\text{C})(20.0 \text{ g})(95^\circ\text{C}-t) = (1 \text{ cal/g}^\circ\text{C})(100 \text{ g})(t-25^\circ\text{C})$$

$$205.2 \text{ cal} - (2.16 \text{ cal/}^\circ\text{C})t = (100 \text{ cal/}^\circ\text{C})t - 2500 \text{ cal}$$

$$2705.2 \text{ cal} = (102.16 \text{ cal/}^\circ\text{C})t$$

$$\frac{2705.2 \text{ cal}}{102.16 \text{ cal/}^\circ\text{C}} = t$$

$$26.48^\circ\text{C} = t.$$

• PROBLEM 487

A container has the dimensions 2.5 m \times 50 cm \times 60 mm. If it is filled with water at 32°F, how many kilocalories will be necessary to heat the water to the boiling point? (212°F)

Solution: A kilocalorie is defined as the amount of heat necessary to raise the temperature of 1 kg of water 1°C. Thus, to solve for the number of kilocalories necessary, one must first solve for the number of grams present and the number of °C that the temperature is raised. One cubic cm of water weighs 1 gram. Thus, if one solves for the volume in terms of cubic centimeters, the weight is then quickly determined. One can solve for the volume in cubic centimeters after converting the lengths of dimensions to centimeters. 1 m = 100 cm, therefore 2.5 m = 250 cm.

If 1 cm = 10 mm, then 60 mm = 6 cm.

Solving for the volume:

$$\text{volume} = 250 \text{ cm} \times 50 \text{ cm} \times 6 \text{ cm} = 7.50 \times 10^4 \text{ cc.}$$

Therefore, the weight of the water is $7.5 \times 10^4 \text{ g}$.
1 kg = 1000 g, thus $7.5 \times 10^4 \text{ g} = 75 \text{ kg}$.

To convert from °F to °C one uses the following formula:

$$^\circ\text{C} = 5/9 (^\circ\text{F} - 32)$$

Solving for °C when t = 32°F:

$$^\circ\text{C} = 5/9 (32 - 32) = 0^\circ\text{C}.$$

When t = 212°F.

$$^\circ\text{C} = 5/9 (212-32) = 5/9 (180) = 100^\circ\text{C}.$$


Therefore, the change in temperature is $100^{\circ} - 0^{\circ}$ or 100°C .

Solving for the number of kilocalories needed:

$$\text{no. of Kcal} = \frac{1 \text{ Kcal}}{^{\circ}\text{C kg}} \times 100^{\circ}\text{C} \times 75 \text{ kg} = 7.5 \times 10^3 \text{ Kcal.}$$

• PROBLEM 488

The following reaction using hydrogen and oxygen is carried out in a bomb calorimeter: $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{l})$. The following data are recorded: Weight of water in calorimeter = 2.650 kg., Initial temperature of water = 24.442°C ., Final temperature of water after reaction = 25.635°C ., Specific heat of reaction vessel is $0.200 \text{ Kcal/}^{\circ}\text{C-kg}$, the weight of the calorimeter is 1.060 kg, and the specific heat of water is $1.00 \text{ Kcal/}^{\circ}\text{C-kg}$, calculate the heat of reaction. Assuming the 0.050 mole of water was formed in this experiment, calculate the heat of reaction per mole of liquid water formed. Neglect the specific heat of the thermometer and stirrer.



Solution: The heat emitted during the reaction must be equal to the heat absorbed by the system, including the water and the reaction vessel. Therefore, the heat of reaction can be found by calculating the heat absorbed by the water and the heat absorbed by the reaction vessel.

1) To find the heat absorbed by the water one must be concerned with the specific heat of water, the amount of water present and the number of degrees the temperature is raised. The specific heat is defined as the number of kilocalories absorbed when a kilogram of mass is raised one degree. The specific heat of water is $1 \text{ Kcal/kg-}^{\circ}\text{C}$. To find the total heat absorbed by the water, the weight of the water and the number of degrees that the temperature is raised must be multiplied by the specific heat.

$$\begin{aligned}\text{Number of Kcal absorbed by water} &= 2.650 \text{ kg} \times 1 \text{ kg/}^\circ\text{C} - \text{kg} \\ &\times (25.635^\circ - 24.442^\circ\text{C}) \\ &= 3.161 \text{ Kcal.}\end{aligned}$$

2) To determine the number of Kcal absorbed by the reaction vessel a similar procedure is used. The specific heat of the vessel is multiplied by the weight of the vessel and the number of degrees the temperature was raised.

$$\begin{aligned}\text{Number of Kcal absorbed by the reaction vessel} &= \\ &= 1.060 \text{ kg} \times 0.200 \text{ Kcal/kg-}^\circ \times (25.635^\circ - 24.442^\circ) \\ &= 0.253 \text{ Kcal.}\end{aligned}$$

3) The heat of reaction is the sum of the heat absorbed by the water and the reaction vessel.

$$\text{heat of reaction} = (3.161 + 0.253) \text{ Kcal} = 3.414 \text{ Kcal.}$$

A ratio can be set up to determine the heat of reaction per mole of $\text{H}_2\text{O}(\ell)$ if 3.414 Kcal is the heat of reaction of 0.05 mole of water. The heat of reaction of 1 mole of water will be related to one as 3.415 Kcal is related to 0.05 moles. Let x = the heat of reaction of 1 mole of water.

$$\frac{3.415 \text{ Kcal}}{0.05 \text{ mole}} = \frac{x}{1 \text{ mole}}$$


$$x = 68.3 \text{ Kcal.}$$

Therefore the heat formed when one mole of water is formed is 68.3 Kcal.

ENTHALPY

• PROBLEM 489

Determine ΔH° for the following reaction of burning ethyl alcohol in oxygen:



$$\Delta H_f^\circ \text{ of } \text{C}_2\text{H}_5\text{OH}(\ell) = -65.9 \text{ Kcal/mole}$$


$$\Delta H_f^\circ \text{ of } \text{CO}_2(\text{g}) = -94.1 \text{ Kcal/mole}$$

$$\Delta H_f^\circ \text{ of } \text{H}_2\text{O}(\ell) = -68.3 \text{ Kcal/mole}$$

Solution: The heat reaction (ΔH°) may be found from the heats of formation (ΔH_f°) by subtracting the sum of the heats of formation of all reactants from the sum of the heats of forma-

$$\Delta H^\circ = [2 \times (-267) + (-94.1)] - [3 \times (-197) + (-26.4)]$$


$$= -10.7 \text{ Kcal.}$$



$$\Delta H^\circ = (3 \times \Delta H^\circ \text{ of FeO} + \Delta H^\circ \text{ of CO}_2) - (\Delta H^\circ \text{ of Fe}_3\text{O}_4 + \Delta H^\circ \text{ of CO})$$

$$\Delta H^\circ = (3 \times (-63.7) + (-94.1)) - ((-267) + (-26.4))$$

$$= 8.2 \text{ Kcal.}$$



$$\Delta H^\circ = (\Delta H^\circ \text{ of Fe} + \Delta H^\circ \text{ of CO}_2) - (\Delta H^\circ \text{ of FeO} + \Delta H^\circ \text{ of CO})$$


The ΔH° of any element is 0. Thus, the ΔH° of Fe is 0.

$$\Delta H^\circ = (0 + (-94.1)) - ((-63.7) + (-26.4))$$


$$= -4.0 \text{ Kcal.}$$

• PROBLEM 491

Given the following reactions:




$$\Delta H = -71.0 \text{ Kcal}$$




$$\Delta H = -23.5 \text{ Kcal}$$

calculate ΔH for the reaction:





Solution: The heat of a given chemical change is the same whether the reaction proceeds in one or several steps; in other words, the energy change is independent of the path taken by the reaction. The heat for a given reaction is the algebraic sum of the heats of any sequence of reactions which will yield the reaction in question. For this problem, one can add the 2 reactions together to obtain the third. The ΔH is found for the third reaction by adding the ΔH 's of the other two.

$$\begin{array}{r} \Delta H = -71.0 \text{ Kcal} \\ + \Delta H = -23.5 \text{ Kcal} \\ \hline -94.5 \text{ Kcal} \end{array}$$



This equation can be simplified by subtracting $\text{SO}_2(\text{g})$ from each side and adding O_2 and $\frac{1}{2}\text{O}_2$ together. You have, therefore



The ΔH for this reaction is - 94.5 Kcal, as shown above.

• PROBLEM 492

Given that $\Delta H^\circ_{\text{CO}_2(\text{g})} = - 94.0$, $\Delta H^\circ_{\text{CO}(\text{g})} = - 26.4$, $\Delta H^\circ_{\text{H}_2\text{O}(\text{l})} = - 68.4$ and $\Delta H^\circ_{\text{H}_2\text{O}(\text{g})} = - 57.8$ in Kcal/mole, determine the heats of reaction of (1) $\text{CO}(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g})$, (2) $\text{H}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{H}_2\text{O}(\text{l})$ and (3) $\text{H}_2\text{O}(\text{l}) \rightarrow \text{H}_2\text{O}(\text{g})$.

Solution: To solve this problem, you need to know what heat of reaction means and how to determine it quantitatively. The heat of reaction may be defined as the heat released or absorbed as a reaction proceeds to completion. It is measured quantitatively from the heats of formation of products minus heats of formation of reactants. Namely, heat of reaction, $\Delta H^\circ = \Delta H^\circ_{\text{products}} - \Delta H^\circ_{\text{reactants}}$. (Note: By convention, the ΔH° of any element is defined as being equal to zero.) You proceed as follows:

- $$\begin{aligned} (1) \quad \Delta H^\circ &= \Delta H^\circ_{\text{CO}_2(\text{g})} - \Delta H^\circ_{\text{CO}(\text{g})} - \frac{1}{2}\Delta H^\circ_{\text{O}_2(\text{g})} \\ &= - 94 - (- 26.4) - \frac{1}{2}(0) = - 67.6 \text{ Kcal/mole.} \\ (2) \quad \Delta H^\circ &= \Delta H^\circ_{\text{H}_2\text{O}(\text{l})} - \Delta H^\circ_{\text{H}_2(\text{g})} - \frac{1}{2}\Delta H^\circ_{\text{O}_2(\text{g})} \\ &= - 68.4 - 0 - \frac{1}{2}(0) = - 68.4 \text{ Kcal/mole.} \\ (3) \quad \Delta H^\circ &= \Delta H^\circ_{\text{H}_2\text{O}(\text{g})} - \Delta H^\circ_{\text{H}_2\text{O}(\text{l})} \\ &= - 57.8 - (- 68.4) = + 10.6 \text{ Kcal/mole.} \end{aligned}$$

• PROBLEM 493

You are given the following reactions at 25°C :
 $2\text{NaHCO}_3(\text{s}) \rightarrow \text{Na}_2\text{CO}_3(\text{s}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$, $\Delta H = 30.92 \text{ Kcal/mole}$. $\Delta H^\circ_{\text{Na}_2\text{CO}_3(\text{s})} = - 270.3$, $\Delta H^\circ_{\text{CO}_2(\text{g})} = - 94.0$ and $\Delta H^\circ_{\text{H}_2\text{O}(\text{l})} = - 68.4 \text{ Kcal/mole}$, what is the standard enthalpy of formation for $\text{NaHCO}_3(\text{s})$?

Solution: The ΔH° for a reaction is an indication of the amount of heat released (or absorbed) when the reactants are converted to products. In general, ΔH° for a reaction will be the sum of the heats of formation of products minus the sum of the heats of formation of reactants, each of which is multiplied by its coefficient in the equation. Because you are given this sum and the ΔH° (heats of formation) of all products, the standard enthalpy of formation of NaHCO_3 can be found. Proceed as follows:

$$\Delta H^\circ = \Delta H_f^\circ \text{Na}_2\text{CO}_3(\text{s}) + \Delta H_f^\circ \text{CO}_2(\text{g}) + \Delta H_f^\circ \text{H}_2\text{O}(\text{l}) - 2\Delta H_f^\circ \text{NaHCO}_3(\text{s})$$

Substituting known values,

$$30.92 = (-270.3) + (-94.0) + (-44.8) - 2\Delta H_f^\circ \text{NaHCO}_3(\text{s})$$

Solving for $\Delta H_f^\circ \text{NaHCO}_3(\text{s})$, you obtain

$$\begin{aligned} \Delta H_f^\circ \text{NaHCO}_3(\text{s}) &= (-270.3 - 94.0 - 44.8 - 30.9)/2 \\ &= -440.0/2 = -220.0 \text{ Kcal/mole,} \end{aligned}$$

which is its standard enthalpy of formation.

• PROBLEM 494

Calculate the standard enthalpy change, ΔH° , for the combustion of ammonia, $\text{NH}_3(\text{g})$, to give nitric oxide, $\text{NO}(\text{g})$, and water $\text{H}_2\text{O}(\text{l})$. The enthalpies of formation, ΔH_f° , are -68.32 Kcal/mole for $\text{H}_2\text{O}(\text{l})$, -11.02 Kcal/mole for $\text{NH}_3(\text{g})$, and 21.57 Kcal/mole for $\text{NO}(\text{g})$.

Solution: The enthalpy change, ΔH , refers to the change in heat content between the products and the reactants of a chemical reaction. The symbol ΔH° is the enthalpy change for a reaction in which each reactant and product is in its standard state at a specified reference temperature (the common reference temperature is 25°C).

The standard enthalpy of formation, ΔH_f° , of a substance is defined as the change in enthalpy for the reaction in which one mole of the compound is formed from its elements at standard conditions.

For any chemical reaction, the change in enthalpy, ΔH , may be expressed as

$$\Delta H_{\text{reaction}} = \sum H_{\text{products}} - \sum H_{\text{reactants}}$$

(where the symbol $\sum H$ means the summation of the enthalpies of each substance).

To solve this problem, one must first write and balance the equation for the combustion of $\text{NH}_3(\text{g})$:



From the balanced equation, one knows that 4 moles of $\text{NH}_3(\text{g})$ reacts with 5 moles of $\text{O}_2(\text{g})$ to form 4 moles of $\text{NO}(\text{g})$ and 6 moles of $\text{H}_2\text{O}(\text{l})$. For the combustion of 4 moles of $\text{NH}_3(\text{g})$:

$$4 \cdot \Delta H_f^\circ, \text{NH}_3(\text{g}) = 4 \left(-11.02 \frac{\text{Kcal}}{\text{mole}} \right) = -44.08 \text{ Kcal}$$

Heat of formation may be defined as the heat absorbed or evolved in the synthesis of one mole of a compound from its elements, all components being in their standard states.

The heat of reaction is the sum of the heats of formation of the products minus the sum of the heats of formation of the reactants. In other words,

$$\Delta H_{\text{reaction}} = \sum \Delta H_{\text{formation of products}} - \sum \Delta H_{\text{formation of reactants}}$$

Therefore, for the hydrolysis of urea to CO_2 and NH_3 :

$$\Delta H_{\text{reaction}} = \left[\left(\Delta H_f^\circ, \text{CO}_2 + 2\Delta H_f^\circ, \text{NH}_3 \right) - \left(\Delta H_f^\circ, \text{H}_2\text{N-CO-NH}_2 + \Delta H_f^\circ, \text{H}_2\text{O} \right) \right]$$

Given these values, one can substitute to obtain:

$$\Delta H_{\text{reaction}} = [(-98.69 - 2 \times 19.32) - (-76.30 - 68.32)]$$

From this equation one obtains

$$\Delta H_{\text{reaction}} = 7.29 \text{ Kcal/mole.}$$

• PROBLEM 496

Using the data from the accompanying figure, calculate the heat of reaction for the following process at 25°C :


- a) $\text{CaCl}_2 (\text{s}) + \text{Na}_2\text{CO}_3 (\text{s}) \rightarrow 2\text{NaCl} (\text{s}) + \text{CaCO}_3 (\text{s})$
 b) $\text{H}_2\text{SO}_4 (\text{l}) + 2\text{NaCl} (\text{s}) \rightarrow \text{Na}_2\text{SO}_4 (\text{s}) + 2\text{HCl} (\text{g})$

Standard Heats of Formation, ΔH° ,
in kcal/mole at 25°C .

Substance	ΔH°
$\text{CaCl}_2 (\text{s})$	-190.0
$\text{Na}_2\text{CO}_3 (\text{s})$	-270.3
$\text{NaCl} (\text{s})$	-98.2
$\text{CaCO}_3 (\text{s})$	-288.4
$\text{H}_2\text{SO}_4 (\text{l})$	-193.9
$\text{HCl} (\text{g})$	-22.1
$\text{Na}_2\text{SO}_4 (\text{s})$	-330.9

Solution: Calculate the amount of heat released (or absorbed) from the heats of formation. In general, ΔH for a reaction will be the sum of the heats of formation of products minus the heats of formation of reactants, each of which is multiplied by its coefficient in the equation. Once ΔH is determined, ΔE can be found from $\Delta E = \Delta H - (\Delta n)RT$, where Δn = change of moles, R = universal gas constant and T = temperature in kelvin (Celsius plus 273°)


Proceed as follows:



Thus,

$$\begin{aligned}\Delta H &= 2\Delta H^\circ_{\text{NaCl}(s)} + \Delta H^\circ_{\text{CaCO}_3(s)} - \Delta H^\circ_{\text{CaCl}_2(s)} - \Delta H^\circ_{\text{Na}_2\text{CO}_3(s)} \\ &= 2(-98.2) + (-288.4) - (-190.) - (270.3) \\ &= -24.5 \text{ Kcal/mole.}\end{aligned}$$

$\Delta E = \Delta H - \Delta nRT$. But, in this reaction, no gases appear. This means ΔnRT becomes zero. As such, $\Delta E = \Delta H = -24.5 \text{ Kcal/mole}$.



$$\begin{aligned}\Delta H &= \Delta H^\circ_{\text{Na}_2\text{SO}_4(s)} + 2\Delta H^\circ_{\text{HCl}(g)} - \Delta H^\circ_{\text{H}_2\text{SO}_4(l)} - 2\Delta H^\circ_{\text{NaCl}(s)} \\ &= (-330.9) + 2(-22.1) - (-193.9) - 2(-98.2) = +15.2 \text{ kcal/mole}\end{aligned}$$

$$\Delta E = \text{heat of reaction} = \Delta H - (\Delta n)RT.$$

A gas, HCl , is involved in this reaction.
 Δn = moles products - moles reactants = 2 - 0 = 2

$T = 25 + 273 = 298^\circ\text{K}$; R is in terms of cal. and ΔH in terms of Kcal, so that you must use the conversion factor of 1Kcal/1000 cal.

$$\begin{aligned}\text{Thus, } \Delta E &= -15.2 - (2)(1.987)(298)(1/1000) \\ &= -15.2 - 1.2 \\ &= -16.4 \text{ Kcal/mole.}\end{aligned}$$

• PROBLEM 497

Using the data from the accompanying table find the enthalpy change for the combustion of a mole of $\text{C}_2\text{H}_4(g)$ to form $\text{CO}_2(g)$ and $\text{H}_2\text{O}(g)$ at 298°K and 1 atm standard state conditions.

Solution: When a chemical reaction occurs heat is usually absorbed or given off to the environment. This means that the enthalpy or heat content of the system will also change. If one knows the enthalpy change of the combustion of each reactant and product, one can find the change in enthalpy of a reaction. The reaction in this problem can be written as



For the reaction,

$$\Delta H_{\text{combustion}} = \sum \Delta H_{\text{products}} - \sum \Delta H_{\text{reactants}}.$$

Remembering that the enthalpy change for the formation of an element (i.e. O_2) in its standard state is zero, one then has,

$$\begin{aligned}\Delta H_{\text{combustion}} &= 2\Delta H_{\text{CO}_2} + 2\Delta H_{\text{H}_2\text{O}} - \Delta H_{\text{C}_2\text{H}_4} - 3\Delta H_{\text{O}_2} \\ &= 2(-394) + 2(-242) - (52) - 3(0) \\ &= -1324 \text{ KJ}\end{aligned}$$

Enthalpies of formation in kilojoules per mole from the elements for various compounds at 298°k and 1 atm pressure

Compound	ΔH
$\text{C}_2\text{H}_4(\text{g})$	+ 52
$\text{CO}_2(\text{g})$	-394
$\text{H}_2\text{O}(\text{g})$	-242
$\text{O}_2(\text{g})$	0

• PROBLEM 498

Calculate the quantity of heat required to (a) convert a liter of H_2O at 30°C to a liter of water at 60°C , and (b) heat a 1 kg block of aluminum from 30° to 60°C . Assume the specific heat of water and aluminum is, respectively, 1 cal/g°C and .215 cal/g°C.

Solution: When heat is added to a mass of unspecified substance, the temperature will rise. The quantitative

relationship between the quantity of heat, expressed in kilocalories, and the rise in temperature - provided there is no phase change - is:

$$\text{quantity of heat} = \text{mass} \times \text{specific heat} \times \Delta T,$$

where ΔT is the change in temperature and the specific heat equals the amount of heat required to raise the temperature of 1 g of any substance by 1°C.

(a) To find the mass of H_2O , remember that density = mass/volume, and the density of H_2O is one. Thus, mass = (1 liter H_2O) (1000 ml/liter) (1.00 g H_2O)/ml. H_2O). Therefore,

$$\begin{aligned}\text{quantity of heat} &= (\text{mass})(\text{specific heat})(\Delta T) \\ \text{mass} &= (1\ell \text{H}_2\text{O})(1000 \text{ ml}/\ell)(1 \text{ g}/\text{ml}) \\ &= 1000 \text{ g} \\ \text{specific heat} &= 1 \text{ cal}/\text{g}^\circ\text{C} \\ \Delta T &= (60 - 30)^\circ\text{C} \\ &= 30^\circ\text{C} \\ \text{quantity of heat} &= (1000 \text{ g})(1 \text{ cal}/\text{g}^\circ\text{C})(30^\circ\text{C}) \\ &= 30,000 \text{ cal} = 30 \text{ Kcal}\end{aligned}$$

(b) A similar calculation can be used to determine the amount necessary to raise the temperature of the aluminum from 30 to 60°C.

$$\begin{aligned}\text{mass} &= (1 \text{ kg Al})(1000 \text{ g}/\text{kg}) = 1000 \text{ g} \\ \text{specific heat} &= .215 \text{ cal}/\text{g}^\circ\text{C} \\ \Delta T &= (60 - 30)^\circ\text{C} \\ &= 30^\circ\text{C} \\ \text{quantity of heat} &= (1000 \text{ g})(.215 \text{ cal}/\text{g}^\circ\text{C})(30^\circ\text{C}) \\ &= 6450 \text{ cal} = 6.45 \text{ Kcal}.\end{aligned}$$

● PROBLEM 499

In a pound-for-pound comparison, which would you prefer as a rocket fuel, hydrogen (H_2) or dimethylhydrazine ($(\text{CH}_3)_2\text{NNH}_2$)? $\text{H}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{H}_2\text{O}(\ell)$, $\Delta H = -68.3 \text{ Kcal}/\text{mole}$
 $(\text{CH}_3)_2\text{NNH}_2 + 4\text{O}_2(\text{g}) \rightarrow \text{N}_2(\text{g}) + 4\text{H}_2\text{O}(\text{g}) + 2\text{CO}_2(\text{g})$, $\Delta H = -404.9 \text{ Kcal}/\text{mole}$.

Solution: The material that is the better fuel will release more energy per unit weight in its reaction with oxygen. Calculate the energy released per unit weight for H_2 and $(CH_3)_2NNH_2$. You can do this by using the heats of formation and enthalpy changes given. Calculate the total mass or weight of materials from moles that react and divide the ΔH by it to obtain the amount of energy released per unit weight.

For hydrogen:

1 mole of H_2 reacts with $\frac{1}{2}$ mole of oxygen (1 mole H_2) (1.008 g H_2 /mole H_2) + ($\frac{1}{2}$ mole O_2) (16.00 g O_2 /mole O_2) = 1.008 g + 8.00 g = 9.01 g. $\Delta H = -68.3$ Kcal. Thus,

$$\frac{-68.3}{9.01} = -7.58 \text{ Kcal/g.}$$

For dimethylhydrazine:


[1 mole $(CH_3)_2NNH_2$] [60.10 g $(CH_3)_2NNH_2$ /mole $(CH_3)_2NNH_2$] + (4 moles O_2) (16.00 g. O_2 /mole O_2) = 60.10 g + 64.00 g = 124.10 g. $\Delta H = -404.9$. The amount of heat released per unit weight with $(CH_3)_2NNH_2 + O_2 =$

$$\frac{-404.9 \text{ Kcal}}{124.10 \text{ g}} = -3.26 \text{ Kcal/g.}$$

Hydrogen yields over twice as much energy per gram of fuel mass, and is thus preferable in this respect.

• PROBLEM 500

Calculate ΔH_r° for the combustion of methane, CH_4 . The balanced reaction is



Solution: ΔH_r° is the Standard Enthalpy Change of Reaction. Standard conditions are defined as 25°C and 1 atm.

Enthalpy is the heat content of a system. If the overall change in enthalpy is negative, then heat is given off to the surroundings and the reaction is called exothermic. When the change is positive, heat is absorbed and the reaction is endothermic. Endothermic compounds are often unstable and can sometimes explode. An endothermic compound, however, is a more efficient fuel because, upon combustion, it yields more heat energy.

ΔH_r° is calculated using the enthalpies of formation, ΔH_f° . The sum of enthalpies of formation of products minus the sum of the enthalpies of formation of

The question asks for ΔH by unit weight and volume. Therefore, ΔH by unit weight =

$$\frac{\Delta H^\circ}{\text{mol.wt.CO} + \text{mol.wt.H}_2} = \frac{-526 \text{ KJ}}{30 \text{ g}} = -17.5 \text{ KJ/g.}$$

The sum of the molecular weights of $\text{CO} + \text{H}_2$ is 30. To find ΔH by molar volume, note that two molar volumes are involved, one from H_2 and one from CO , which means

$$\begin{aligned}\Delta H \text{ by volume} &= \frac{\Delta H}{\text{no. of molar volumes}} \\ &= \frac{-526 \text{ KJ}}{2 \text{ molar volumes}} = -263 \text{ KJ/molar volume.}\end{aligned}$$

An analogous method is used for methane. The reaction is $\text{CH}_4 + \text{O}_2 \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g})$.

$$\Delta H^\circ = -394 + 2(-242) - (-75) = -803 \text{ KJ}$$

The molecular weight of $\text{CH}_4 = 16$. Therefore,

$$\Delta H \text{ per unit weight} = \frac{-803 \text{ KJ}}{16 \text{ g}} = -50.2 \text{ KJ/g.}$$

The only molar volume is that of methane. Then


$$\frac{\Delta H}{\text{molar volume}} = \frac{-803 \text{ KJ}}{1}$$

The comparative heats can thus be related as

$$\frac{263}{803} = \frac{.33}{1} \text{ by volume and } \frac{17.5}{50} = \frac{.35}{1} \text{ by weight.}$$

• PROBLEM 502

A common rocket propellant is a mixture of one part liquid hydrazine (N_2H_4) as fuel and two parts liquid hydrogen peroxide (H_2O_2) as oxidizer. These reactants are hypergolic, that is, they ignite on contact. The reaction proceeds according to the equation



If reactants are mixed at 25°C , what is the heat of reaction?

Solution: We will make use of the following expression for the heat of reaction at 25°C (298°K), ΔH_{298}° :

$$\Delta H_{298}^\circ = \text{sum of heats of formation of products} - \text{sum of heats of formation of reactants.}$$

Denoting the heat of formation of species s by $\Delta H_{f,s}^\circ$, this becomes

$$\Delta E^{\circ} = \Delta H^{\circ} - \Delta nRT$$


$$\Delta H^{\circ} = - 279.40 \text{ Kcal}$$

$$\Delta n = 5 \text{ moles}$$

$$R = 1.99 \text{ cal/mole}^{\circ}\text{K}$$

$$T = 298^{\circ}\text{K}$$

$$\begin{aligned}\Delta E^{\circ} &= - 279.4 \text{ Kcal} \times \left(\frac{1000 \text{ cal}}{1 \text{ Kcal}} \right) - (- 5 \text{ mole} \\ &\quad \times 1.99 \text{ cal/mole } ^{\circ}\text{K} \times 298^{\circ}\text{K}) \\ &= - 279,400 \text{ cal} + 2965 \text{ cal} \\ &= - 276,435 \text{ cal} = - 276.4 \text{ Kcal.}\end{aligned}$$



$$\Delta H^{\circ} = - 134.0 \text{ Kcal}$$

ΔH° should be expressed in calories

$$\begin{aligned}\Delta H^{\circ} \text{ in cal} &= - 134.0 \text{ Kcal} \times 1000 \text{ cal/1 Kcal} \\ &= - 134,000 \text{ cal}\end{aligned}$$

$$\Delta n = (\text{no. of moles of SO}_2) - (\text{no. of moles of H}_2\text{S} + \text{no. of moles of O}_2)$$

$$\begin{aligned}\Delta n &= (1 \text{ mole of SO}_2) - (1 \text{ mole of H}_2\text{S} + 1\frac{1}{2} \text{ moles of O}_2) \\ &= (1) - (1 + 1\frac{1}{2}) = - 1.5 \text{ moles}\end{aligned}$$

$$R = 1.99 \text{ cal/mole } ^{\circ}\text{K}$$


$$T = 298^{\circ}\text{K}$$

$$\Delta E^{\circ} = \Delta H^{\circ} - \Delta nRT$$

$$\begin{aligned}&= - 134,000 \text{ cal} - (- 1.5 \text{ moles} \times 1.99 \text{ cal/mole}^{\circ}\text{K} \times 298^{\circ}\text{K}) \\ &= - 134,000 \text{ cal} + 890 \text{ cal} \\ &= - 133,110 \text{ cal} = - 133.1 \text{ Kcal.}\end{aligned}$$

• PROBLEM 508

Calculate ΔH for the reaction at 25°C .



Solution: The change in enthalpy (ΔH) is defined by the first law of thermodynamics as the sum of the change of the internal energy (ΔE) and the work done by a particular system. Work is defined as the product of the pressure (P) and the change in volume (ΔV) of a system, where only pressure-volume work is done. Much of the experimentation done by chemists is pressure-volume work because the work

is usually done in an open container in the laboratory. The pressure, in this case, is constant and the volume of the materials is allowed to change.

$$\Delta H = \Delta E + P\Delta V$$

In this problem, one is given ΔE but not P or ΔV , so that one must calculate $P\Delta V$ in some other way. The Ideal gas Law is defined as:

$$P\Delta V = \Delta nRT,$$

where P is the pressure, ΔV is the change in volume, Δn is the change in the number of moles reacting, R is the gas constant (1.99 cal/mole-°K), and T is the absolute temperature. Here, one knows R and can calculate T and Δn . Therefore, ΔnRT can be substituted for $P\Delta V$ in the equation to find ΔH . Δn is found by subtracting the number of moles reacting from the number of moles formed as products. Here, 1 mole of CO and 0.5 mole of O_2 form 1 mole of CO_2 .

$$\Delta n = 1 \text{ mole } \text{CO}_2 - (1 \text{ mole } \text{CO} + 0.5 \text{ mole } \text{O}_2)$$

$$= - 0.5 \text{ mole.}$$

The absolute temperature is found by adding 273 to the temperature in °C.

$$T = 25 + 273 = 298^\circ\text{K.}$$

$$\begin{aligned} \text{Therefore, } \Delta nRT &= (- 0.5 \text{ mole}) \times 1.99 \text{ cal/mole-}^\circ\text{K} \times 298^\circ\text{K} \\ &= - 0.3 \text{ Kcal.} \end{aligned}$$

One is now ready to find ΔH , via substitution.

$$\Delta H = \Delta E + \Delta nRT$$

$$\Delta E = - 67.4 \text{ Kcal}$$

$$\Delta nRT = - 0.3 \text{ Kcal}$$

$$\Delta H = - 67.4 \text{ Kcal} + (- 0.3 \text{ Kcal}) = - 67.7 \text{ Kcal.}$$

• PROBLEM 509

The equation for the burning of naphthalene is $\text{C}_{10}\text{H}_8(\text{s}) + 12\text{O}_2(\text{g}) \rightarrow 10\text{CO}_2 + 4\text{H}_2\text{O}(\text{l})$. For every mole of C_{10}H_8 burned, - 1226.7 Kcal is evolved at 25° in a fixed-volume combustion chamber. ΔH° for $\text{H}_2\text{O}(\text{l}) = - 64.4 \text{ Kcal/mole}$ and $\text{H}_2\text{O}(\text{g}) = - 57.8 \text{ Kcal/mole}$. Calculate (a) the heat of reaction at constant temperature and (b) ΔH for the case where all of the H_2O is gaseous.

Solution: This problem deals with the heat evolved

when a compound is heated with oxygen to form carbon dioxide and water (the heat of combustion). The heat of reaction, or combustion in this case, ΔH , is given by the formula $\Delta H = \Delta E + \Delta nRT$, where ΔE = amount of heat released per mole, Δn = change in moles, R = universal gas constant and T = temperature in Kelvin (Celsius plus 273°). Thus, to answer (a) substitute these values and solve for ΔE . Δn is moles of gas produced - moles of gas reacted, which is 10 moles - 12 moles = -2 moles based on the coefficients in the equation. $T = 25^\circ + 273 = 298^\circ K$. Use R in terms of kilocalories. As such

$$\Delta H = -1226.7 \text{ kcal} + (-2 \text{ moles})(1.987 \text{ cal/mole } ^\circ K)(298^\circ K)(1 \text{ Kcal}/1000 \text{ cal})$$

$$= -1226.7 - 1.2 = -1227.9 \text{ Kcal/mole.}$$

To find (b), note that the reaction is the same, except that H_2O is gaseous not liquid. You have, therefore, $4H_2O(l) \rightarrow 4H_2O(g)$. The ΔH , change in enthalpy, for this conversion is $4\Delta H^\circ(l) - 4\Delta H^\circ(g)$ of $H_2O = 4(-57.8) - 4(-68.4) = 42.4 \text{ Kcal/mole}$. It follows, then, that the resulting

$$\Delta H = -1227.9 + 42.4 = -1185.5 \text{ Kcal/mole.}$$

• PROBLEM 510

In the reaction, $CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$ at $950^\circ C$ and CO_2 pressure of 1 atm, the ΔH is found to be 176 kJ/mole. Assuming that the volume of the solid phase changes little by comparison with the volume of gas generated, calculate the ΔE for this reaction.

Solution: To solve this problem, you must know that ΔE , the change in energy, can be related quantitatively to the change in enthalpy, ΔH , the pressure (P) and the change in volume (ΔV), by the formula $\Delta E = \Delta H - P\Delta V$. This equation means that if heat is added to a system at constant pressure, part of it goes into increasing the internal energy of the system (ΔE) and the rest is used to do work on its surroundings, $P\Delta V$. Therefore, the total of both ΔE and $P\Delta V$ is the heat content of the system, ΔH . To solve the problem, therefore, you need only substitute in the values for P , ΔV and ΔH .

$$\Delta V = V_{\text{products}} - V_{\text{reactants}}$$

There is little change in the volume of the solid by comparison with the gas. This means, therefore, that ΔV is approximately equal to V_{gas} . V_{gas} can be found from the equation of state; $PV = nRt$ or $V = nRt/P$, where n = number of moles, P = pressure, R = universal gas constant and t = temperature in Kelvin (Celsius temp + 273). From

the reaction, 1 mole of CO_2 gas is produced, thus $n = 1$. R is defined as being equal to $8.31 \text{ J mol}^{-1} \text{ deg}^{-1}$. You are given P and t . Therefore

$$v = \frac{1(8.31)(1223)}{1} = 10.2 \text{ kJ}$$

Recalling, $\Delta E = \Delta H - P\Delta V$, you have

$$\Delta E = 176 - 1(10.2) = 166 \text{ kJ.}$$

• PROBLEM 511

You have the reaction $\text{H}_2\text{O}(l) \rightarrow \text{H}_2\text{O}(g)$. For both states, 1 mole of water is at 100°C and 1 atm pressure. The volume of 1 mole of water = 18 ml, $\Delta H = 9710 \text{ cal/mole}$ and there are 24.2 cal/liter-atm. Calculate the work done in this conversion and the value of ΔE .

Solution: In the conversion from liquid water to steam, the pressure and temperature remain constant, work (w) = P (pressure) times ΔV (change in volume). You are given the pressure.

To find the volume, note ΔV = volume of gas - volume of liquid. Thus, to find the work done in the conversion, find the volume that the steam occupies. This can be done by using the equation of state, which indicates $PV = nRT$, where P = pressure, V = volume, n = moles, R = universal gas constant and T = temperature in Kelvin (Celsius plus 273°). You know all values, except V . Substituting and re-writing,

$$v = \frac{nRT}{P} = \frac{(1)(.0821)(373)}{1} = 30.62 \text{ liters.}$$

This is the volume of the steam. The volume of the liquid = 18 ml or .018 liters. Thus, $\Delta V = 30.62 - .018 = 30.602 \text{ l}$ and $w = \text{work} = (1 \text{ atm})(30.602 \text{ liter}) = 30.602 \text{ liter-atm}$, 24.2 cal. exist per liter-atm. Thus, $w = \text{work} = 24.2 \text{ cal/liter-atm} \times 30.602 \text{ liter-atm} = 740.56 \text{ cal}$. To find ΔE , remember that $\Delta E = \Delta H - P\Delta V$, where ΔE = change of energy, ΔH = change of enthalpy and $P\Delta V$ = work, which was calculated. You are given ΔH , therefore

$$\Delta E = 9710 \text{ cal} - 740.56 \text{ cal} = 8969.44 \text{ cal.}$$

• PROBLEM 512

A chemist expands an ideal gas against a constant external pressure of 700 mmHg, and finds its volume changes from 50 to 150 liters. He finds that 1.55 Kcal of heat have been absorbed in the process. Determine the internal energy change that took place. $24.217 \text{ cal} = 1 \text{ liter-atm}$.


Solution: The internal energy change of a system at constant pressure is given by the formula $\Delta E = \Delta H - P\Delta V$, where ΔE = change in internal energy, ΔH = enthalpy or heat absorbed or released, P = pressure and ΔV = change in volume.

In this problem, $\Delta V = 150 - 50 = 100$ liters. The pressure, in atms, = $(700 \text{ mm})(1 \text{ atm})/(760 \text{ mm}) = .921 \text{ atm}$. To convert, liter-atm to kilocalories, multiply by 24.217 cal./liter-atm and then multiply by Kcal/1000 cal. Therefore,

$$\begin{aligned}\Delta E &= 1.55 \text{ Kcal} - (.921 \text{ atm})(100 \text{ l})(24.217 \text{ cal/liter-atm}) \\ &\hspace{15em} (\text{Kcal/1000 cal}) \\ &= 1.55 - 2.23 = - .68 \text{ Kcal.}\end{aligned}$$

• PROBLEM 513

Exactly one mole of gaseous methane is oxidized at fixed volume and at 25°C according to the reaction



If 212 Kcal is liberated, what is the change in enthalpy, ΔH ?

Solution: We must first find the change in internal energy, ΔE , before applying the equation,

$$\Delta H = \Delta E + \Delta nRT,$$

to find the enthalpy change (Δn = number of moles of gaseous products - number of moles of gaseous reactants, $R = 1.987 \text{ cal/deg-mole}$, T = absolute temperature).

Since heat was evolved (exothermic reaction), ΔE must be negative. Furthermore, since no useful work was done ($\Delta V \neq 0$ if work is done), ΔE is equal in magnitude to the quantity of heat evolved. Thus $\Delta E = - 212 \text{ Kcal}$. Since there is one mole of gaseous product (1 $\text{CO}_2(\text{g})$) and three moles of gaseous reactants ($2\text{O}_2(\text{g}) + 1 \text{CH}_4(\text{g})$), $\Delta n = 1 - 3 = - 2 \text{ mole}$. The absolute temperature is $T = 25 + 273 = 298^\circ\text{K}$. Hence

$$\begin{aligned}\Delta H &= \Delta E + \Delta nRT \\ &= - 212 \text{ Kcal} + (- 2 \text{ mole})(1.987 \text{ cal/deg-mole})(298^\circ\text{K}) \\ &= - 212 \text{ Kcal} - 1180 \text{ cal} \\ &= - 212 \text{ Kcal} - 1.180 \text{ Kcal} \\ &\approx - 213 \text{ Kcal.}\end{aligned}$$

For the following exothermic reaction at 25°C,



the change in energy, ΔE , is - 780 Kcal/mole. Find the change in enthalpy, ΔH .

Solution: The solution is based on the following equation for the enthalpy change

$$\Delta H = \Delta E + \Delta nRT,$$

where Δn is the number of moles of gaseous products minus the number of moles of gaseous reactants, R is the gas constant (1.987 cal/deg-mole), and T is the absolute temperature. Since there are 6 moles of gaseous products ($6\text{CO}_2(\text{g})$) and $7\frac{1}{2}$ moles of gaseous reactants ($7\frac{1}{2}\text{O}_2(\text{g})$), $\Delta n = 6 - 7\frac{1}{2} = -1.5$. The absolute temperature is $T = 25^\circ + 273 = 298^\circ\text{K}$.

Hence, $\Delta H = \Delta E + \Delta nRT$

$$= - 780 \text{ Kcal} + (- 1.5 \text{ mole})(1.987 \text{ cal/deg-mole}) \times (298^\circ\text{K})$$

$$= - 780 \text{ Kcal} - 888 \text{ cal}$$

$$= - 780 \text{ Kcal} - 0.888 \text{ Kcal}$$

$$\approx - 781 \text{ Kcal.}$$

HEATS OF FUSION AND VAPORIZATION

It is known that the heat of vaporization of water is 5 times as great as the heat of fusion. Explain this fact.

Solution: The heat of vaporization is the quantity of heat necessary to vaporize 1 g. of a liquid substance at its boiling point at constant temperature. The heat of fusion is the quantity of heat necessary to liquefy 1 g. of a solid substance at constant temperature at its melting point. With this in mind, let us consider what goes on in each of these phase changes. The volume-increase of the gas in going from a liquid state to a vapor state is much greater than the volume-increase accompanying a solid to liquid transformation. When you increase volume, work is required to overcome the existing external pressure that hinders the

$$\begin{aligned}\text{Total amount of heat absorbed} &= 3200 \text{ cal} + 40 t \text{ cal} \\ &= (3200 + 40 t) \text{ cal}.\end{aligned}$$

To find the amount of heat lost by the 100 g of water at 60°C , when it is cooled to a new temperature, t , the specific heat of water is also used. Here, it means that for every gram of water that is lowered one degree, 1 calorie of heat is released. Therefore, to find the amount of heat released by this water, the specific heat will be multiplied by 100 g, the amount of water present, and by $60-t^{\circ}$, which is the number of degrees that the temperature will drop.

$$(60-t)^{\circ} \times (1 \text{ cal/g-degree}) \times (100 \text{ g}) = (6000 - 100t) \text{ cal}.$$

To calculate t , one must set the amount of heat absorbed by the ice equal to the amount of heat lost by the water at 60°C .

amount of heat absorbed = amount of heat lost

$$\text{amount of heat absorbed} = (3200 + 40 t) \text{ cal}$$

$$\text{amount of heat lost} = (6000 - 100 t) \text{ cal}$$

$$(3200 + 40 t) \text{ cal} = (6000 - 100 t) \text{ cal}$$

$$140 t = 2800$$

$$t = 20^{\circ}$$

Therefore, the final temperature is 20°C .

• PROBLEM 519

Determine the quantity of heat required to convert 10 g of ice at 0°C to vapor at 100°C . For water, heat of fusion = 80 cal/g, heat of vaporization = 540 cal/g, and specific heat = 1 cal/g $^{\circ}\text{C}$.

Solution: Heat is absorbed when ice is transformed from the solid to the liquid state. The heat absorbed when 1 g of solid melts is called the heat of fusion. The heat of fusion for water is 80 cal/g. In calculating the amount of heat needed to raise the temperature of the liquid water, the specific heat must be used. The specific heat is defined as the number of calories of heat necessary to raise the temperature of a gram of a substance 1°C . The specific heat of water is 1° , thus 1 calorie is needed to raise the temperature of 1 gram of water 1°C . The quantity of heat necessary to convert 1 g of a liquid into a vapor is termed the heat of vaporization. For water, 540 calories are necessary to change 1 g of liquid water at 100°C into vapor (steam) at 100°C .

In this problem three steps are involved.

Therefore, 7200 calories are needed to melt 10 g of ice, heat it to 100°C and vaporize it to steam.

• PROBLEM 520

500 g of ice at 0°C is added to 500 g of water at 64°C. When the temperature of the mixture is 0°C, what weight of ice is still present? Heat of fusion of $H_2O = 80 \text{ cal/g}$.

Solution: The amount of heat used to melt the ice is equal to the amount of heat lost by the water. The amount of heat lost by the water is determined by using the specific heat of water. The specific heat of water is 1 cal/g-degree. This means that for each gram of water cooled 1 degree, 1 calorie of heat will be evolved. Here, 500 g of H_2O is lowered 64° - 0° or 64°. The heat evolved by the water can now be found.

$$\begin{aligned}\text{no. of calories evolved} &= 1 \text{ cal/g-degree} \times 500 \text{ g} \times 64^\circ \\ &= 32000 \text{ cal.}\end{aligned}$$

The heat of fusion is the quantity of heat necessary to liquefy 1 g of a solid substance at constant temperature at its melting point. Therefore, if the heat of fusion of water is 80 cal/g, then it takes 80 cal to melt one gram of ice. You found that 32000 calories are absorbed by the ice. The weight of ice melted by 32000 cal is found by dividing 32000 cal by 80 cal/g.

$$\text{no. of grams of ice} = \frac{32000 \text{ cal}}{80 \text{ cal/g}} = 400 \text{ g}$$

Thus 400 g of the ice is melted. Originally, there was 500 g of ice, therefore 100 g of ice is left.

• PROBLEM 521

Determine the heat needed to raise 60 g of Pb from 20°C to 360°C, given its specific heats are 0.0306 (solid) and 0.0375 (liquid). Its heat of fusion is 5.86 cal/g; its melting point is 327°C.

Solution: There are three heats involved in finding the amount of heat absorbed in raising the temperature of Pb from 20°C to 360°C. First, the amount of heat absorbed in raising the temperature of the solid from 20°C to 327°C (its melting point); then, the amount of heat absorbed in melting the compound; last, the amount of heat absorbed in raising the temperature of the liquid from 327°C to 360°C. The heat absorbed in these three processes are added together to find the amount of heat needed to bring Pb from

*Heat capacity in joules per
mole-degree for various
substances at the
temperatures indicated.*

$\text{H}_2\text{O}(s)$ at 239°K	33.30
$\text{H}_2\text{O}(s)$ at 271°K	37.78
$\text{H}_2\text{O}(l)$ at 273°K	75.86
$\text{H}_2\text{O}(l)$ at 298°K	75.23
$\text{H}_2\text{O}(l)$ at 373°K	75.90
$\text{H}_2\text{O}(g)$ at 383°K	36.28

Solution: Enthalpy is an indication of the changes in heat content of a system due to changes in the system. To solve this problem, one must be aware that in going from ice at -10°C to water at 15°C , three things happen: (1) the ice is heated, (2) the ice is melted, (3) the liquid is heated. For the heating processes, (1) + (3), the change in enthalpy is quantitatively related to the heat capacity, C_p , and the change in temperature, ΔT by the relation is $\Delta H = C_p \Delta T$. C_p , the heat capacity, is defined as the amount of heat required to raise one mole of material by one degree. The melting process, (2), is expressed in terms of the molar heat of fusion, which is the amount of heat necessary to melt one mole of solid. The summation of the heat involved in all three processes will give the change in enthalpy. Because each step is defined in terms of moles, one must first calculate the number of moles involved. The molecular weight of H_2O is 18. Therefore, there are $100/18$ or 5.55 moles of H_2O in 100 g. Thus,

(1) To heat ice, one needs,

$$(5.55 \text{ moles})(37.78 \text{ J mol}^{-1} \text{ deg}^{-1})(10 \text{ deg}) = 2,100 \text{ J}$$

(2) To melt ice, one requires

$$(5.55 \text{ moles})(6.02 \text{ kJ/mole})(1000 \text{ J/kJ}) = 33,400 \text{ J}$$

(3) To heat liquid, one needs

$$(5.55 \text{ moles})(75.86 \text{ J mol}^{-1} \text{ deg}^{-1})(15 \text{ deg}) = 6,300 \text{ J.}$$

The total increase in enthalpy is

$$\Delta H = 2,100 + 33,400 + 6,300 = 41,800 \text{ J.}$$

CHAPTER 15

THERMODYNAMICS II

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 503 to 538 for step-by-step solutions to problems.

The Second Law of Thermodynamics is not nearly as easy to state as the First Law, but is nonetheless very important in understanding how spontaneous changes occur. For example, the First Law is satisfied when water falls over a waterfall and the temperature increases to account for the change in potential energy from the top to the bottom of the waterfall. But the First Law would also be satisfied if the water went up the waterfall and was simultaneously cooled. Experience, of course, indicates that this does not happen; the Second Law of Thermodynamics provides a formal explanation.

The Second Law of Thermodynamics states that the change in entropy, defined as

$$\Delta S = \Delta q/T \quad 15-1$$

is positive when contributing to spontaneous change. In Equation 15-1, ΔS is the entropy change, Δq is the change in heat (positive, by convention, for heat entering the system and negative for heat leaving), and T is the absolute temperature. Unlike enthalpy, there is an absolute zero for entropy—the elements at 0K. Therefore, absolute values of entropy are tabulated for various compounds—usually at 298K. But entropy change for a reaction is calculated in the same way as the enthalpy changes.

$$\Delta S^\circ_r = \sum S^\circ_{\text{products}} - \sum S^\circ_{\text{reactants}} \quad 15-2$$

A very important thermodynamic quantity involving both entropy and enthalpy is called the Gibbs Free Energy (G) or, sometimes, simply the free energy. It is defined as

The resulting mole fractions ($N_i = n_i/n_T$) are:

$$N_{N_2} = .236$$

$$N_{H_2} = .709$$

$$N_{NH_3} = .055$$

$$\Sigma = 1.000$$

Step-by-Step Solutions to Problems in this Chapter, "Thermodynamics II"

ENTROPY

• PROBLEM 523

For the following reaction at 25°C



Values of S° , the absolute entropies for the substances are:

CuO(s)	=	10.4 cal/mole
$\text{H}_2\text{(g)}$	=	31.2 cal/mole
Cu(s)	=	8.0 cal/g-atm
$\text{H}_2\text{O(g)}$	=	45.1 cal/mole

Assuming standard conditions find out if the reaction will proceed spontaneously.

Solution: Entropy change is often used to predict the spontaneity of a reaction. A process will occur spontaneously if there is an increase in entropy i.e. ΔS° is positive.

$$\Delta S^\circ = S^\circ_{\text{(products)}} - S^\circ_{\text{(reactants)}}$$

For the above reaction

$$\begin{aligned}\Delta S^\circ &= S^\circ_{\text{Cu(s)}} + S^\circ_{\text{H}_2\text{O(g)}} - S^\circ_{\text{CuO(s)}} - S^\circ_{\text{H}_2\text{(g)}} \\ &= (8 + 45.1 - 10.4 - 31.2) \text{ cal/deg-mole} \\ &= +11.5 \text{ cal/deg-mole}\end{aligned}$$

ΔS° is positive thus this is a spontaneous reaction.

When mercury is vaporized at its boiling point at standard pressure, the entropy change is 20.7 cal/mole-°K. Determine the boiling point of Hg if the heat of vaporization is 65 cal/g.

Solution: When a process occurs at constant temperature, the change in entropy, ΔS , is equal to the heat absorbed divided by the absolute temperature at which the change occurs, i.e.,

$$\Delta S = \frac{\Delta H}{T},$$

where ΔH is the heat of vaporization (in this case) and T is the absolute temperature. T will be equal to the boiling point of mercury in this problem.

Using this equation, one can solve for T after either converting ΔH to cal/moles or converting ΔS to cal/g. Here, one will convert ΔH from cal/g to cal/mole by multiplying ΔH by the molecular weight of Hg. (MW of Hg = 200.6.)

$$\begin{aligned}\Delta H \text{ in cal/moles} &= 65 \text{ cal/g} \times 200.6 \text{ g/mole} \\ &= 13039 \text{ cal/mole}\end{aligned}$$

One can now solve for T .

$$T = \frac{\Delta H}{\Delta S}$$

$$\Delta H = 13039 \text{ cal/mole}$$

$$\Delta S = 20.7 \text{ cal/mole-}^\circ\text{K}$$

$$T = \frac{13039 \text{ cal/mole}}{20.7 \text{ cal/mole-}^\circ\text{K}} = 630^\circ\text{K}$$

$$T \text{ in } ^\circ\text{C} = 630 - 273 = 357^\circ\text{C}.$$

Calculate ΔS for the conversion of one mole of liquid water to vapor at 100°C. Heat of vaporization = 540 cal/g.

Solution: When a process occurs at constant temperature, the change in entropy (ΔS) is equal to the heat absorbed divided by the absolute temperature at which the change occurs.

$$\Delta S = \frac{\Delta H}{T}$$

In this problem, one is given ΔH and one can find T .

The absolute temperature is calculated by adding 273 to the temperature in °C.

$$T = 273 + 100 = 373^{\circ}\text{K}$$

Because one mole of water is reacting here, and the heat of vaporization is given in cal/g, one must multiply the ΔH given by the molecular weight of water to find the ΔH in cal/mole.

The molecular weight of water is 18.

$$\Delta H = 540 \text{ cal/g} \times 18 \text{ g/mole} = 9720 \text{ cal/mole}$$

One can now calculate ΔS .

$$\Delta S = \frac{\Delta H}{T}$$

$$\begin{aligned} \Delta H &= 9720 \text{ cal/mole} \\ T &= 373^{\circ}\text{K} \end{aligned}$$



$$\Delta S = \frac{9720 \text{ cal/mole}}{373^{\circ}\text{K}} = 26.1 \text{ cal/mole-}^{\circ}\text{K}$$

The change in entropy, when one mole of water is vaporized at 100°C, is 26.1 cal/mole-°K.


• PROBLEM 526

A chemist knows that the $\Delta H^{\circ} = 485 \text{ kJ}$ for the reaction $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$ and that $\Delta H^{\circ} = -537 \text{ kJ}$ for $\text{H}_2(\text{g}) + \text{F}_2(\text{g}) \rightarrow 2\text{HF}(\text{g})$. With this information, he calculated the ΔH° for $2\text{H}_2\text{O}(\text{g}) + 2\text{F}_2(\text{g}) \rightarrow 4\text{HF}(\text{g}) + \text{O}_2(\text{g})$ and predicted whether ΔS° was positive or negative. How?

Solution: Hess' Law states that the net heat change resulting from a particular chemical reaction is the same, independent of the steps involved in the transformation. Thus, ΔH° of the sum of two reactions equals the sum of the ΔH° 's of each reaction. The chemist knows that



By doubling (ii) and adding to (i), the chemist obtains the desired equation



The ΔH° for this reaction, according to Hess' Law, equals


The reaction with the more negative free energy change will be the more efficient, since it proceeds spontaneously as written. We will require the following standard free energies of formation:

$$\Delta G^\circ_{\text{H}_2\text{O}_2(\ell)} = -27.2 \text{ Kcal/mole}$$

$$\Delta G^\circ_{\text{H}_2\text{O}(\ell)} = -56.7 \text{ Kcal/mole}$$

$$\Delta G^\circ_{\text{H}_2(\text{g})} = \Delta G^\circ_{\text{O}_2(\text{g})} = 0 \text{ Kcal/mole}$$


For the first process



the standard free energy change is

$$\begin{aligned}\Delta G^\circ &= \Delta G^\circ_{\text{products}} - \Delta G^\circ_{\text{reactants}} \\ &= \Delta G^\circ_{\text{H}_2\text{O}_2(\ell)} - \left[\Delta G^\circ_{\text{H}_2(\text{g})} + \Delta G^\circ_{\text{O}_2(\text{g})} \right] \\ &= -27.2 \text{ Kcal/mole} - (0 \text{ Kcal/mole} + 0 \text{ Kcal/mole}) \\ &= -27.2 \text{ Kcal/mole}\end{aligned}$$

For the second process,




the standard free energy change is

$$\begin{aligned}\Delta G^\circ &= \Delta G^\circ_{\text{products}} - \Delta G^\circ_{\text{reactants}} \\ &= 2\Delta G^\circ_{\text{H}_2\text{O}_2(\ell)} - \left[2\Delta G^\circ_{\text{H}_2\text{O}(\ell)} + \Delta G^\circ_{\text{O}_2(\text{g})} \right] \\ &= 2(-27.2 \text{ kcal/mole}) - (2 \times (-56.7) \\ &\quad \text{kcal/mole} + 0 \text{ kcal/mole}) \\ &= 59.0 \text{ Kcal/2 moles H}_2\text{O}_2 \text{ produced.} \\ &= 29.5 \text{ Kcal/mole.}\end{aligned}$$

Since the first process proceeds spontaneously as written (negative ΔG°) and the second process requires energy to proceed as written (positive ΔG°), the first method for preparing H_2O_2 is more efficient than the second.

Determine ΔG° for the reaction



$$\Delta G_f^\circ \text{ of } \text{NH}_3(\text{g}) = -4.0 \text{ Kcal/mole}$$

$$\Delta G_f^\circ \text{ of } \text{NO}(\text{g}) = 20.7 \text{ Kcal/mole}$$

$$\Delta G_f^\circ \text{ of } \text{H}_2\text{O}(\text{l}) = -56.7 \text{ Kcal/mole}$$


Solution: The change in free energy (ΔG°) may be found by subtracting the sum of free energies (ΔG_f°) of the reactants from the free energies of the products. The free energy of formation of pure elements is always 0. When more than 1 mole of a compound is either reacted or formed, the ΔG_f° of that compound must be multiplied by the stoichiometric coefficient for the specific compound.

$$\Delta G^\circ = (4 \text{ moles} \times \Delta G_f^\circ \text{ of } \text{NO}(\text{g}) + 6 \text{ moles} \times \Delta G_f^\circ \text{ of } \text{H}_2\text{O}(\text{l}))$$

$$- (4 \text{ moles} \times \Delta G_f^\circ \text{ of } \text{NH}_3(\text{g}) + 5 \text{ moles} \times \Delta G_f^\circ \text{ of } \text{O}_2)$$

$$\begin{aligned} G^\circ &= (4 \text{ moles} \times 20.7 \text{ kcal/mole} + 6 \text{ moles} \times (-56.7 \text{ kcal/mole})) \\ &\quad - (4 \text{ moles} \times (-4.0 \text{ kcal/mole}) + 5 \text{ moles} \times 0 \text{ kcal/mole}) \\ &= -241.4 \text{ kcal.} \end{aligned}$$

For sublimation of iodine crystals,



at 25°C and atmospheric pressure, it is found that the change in enthalpy, $\Delta H = 9.41 \text{ Kcal/mole}$ and the change in entropy, $\Delta S = 20.6 \text{ cal/deg-mole}$. At what temperature will solid iodine be in equilibrium with gaseous iodine?

Solution: Use the fact that the system is in a state of equilibrium. The change in Gibb's free energy is related to ΔH and ΔS by the equation

$$\Delta G = \Delta H - T\Delta S,$$

where T is the absolute temperature of the system. At equilibrium, $\Delta G = 0$ and T is the equilibrium temperature, T_{equil} . Hence

$$\Delta G = \Delta H - T\Delta S,$$

$$0 = \Delta H - T_{\text{equil}} \Delta S,$$

$$\text{or, } T_{\text{equil}} = \frac{\Delta H}{\Delta S}.$$

Therefore,

$$T_{\text{equil}} = \frac{\Delta H}{\Delta S} = \frac{9.41 \text{ Kcal/mole}}{20.6 \text{ cal/deg-mole}} = \frac{9410 \text{ cal/mole}}{20.6 \text{ cal/deg-mole}} \\ = 457 \text{ K,}$$

$$\text{or, } T_{\text{equil}} = 457 - 273 = 184^{\circ}\text{C.}$$

• PROBLEM 532

At the melting point of a solid (or the freezing point of a liquid), the free energies of the solid state and the liquid state are equal, $\Delta G = 0$. Likewise, at the boiling point of a liquid, where there is an equilibrium between the liquid and vapor phases, the free energy is equal in the two states. Calculate the change in entropy for the following process at 0°C if the heat of fusion of $\text{H}_2\text{O} = 80 \text{ cal/g}$. $\text{H}_2\text{O(s)} \rightarrow \text{H}_2\text{O(l)}$.

Solution: The free energy, ΔG , can be expressed as $\Delta G = \Delta H - T\Delta S$, where ΔH , in this case, is the heat of fusion, T is the absolute temperature and ΔS is the change in the entropy. One is given that $\Delta G = 0$ for this process and thus, $\Delta H - T\Delta S = 0$. One is told that the process takes place at 0°C . To use the free energy equation, the temperature must be in $^{\circ}\text{K}$. To convert from $^{\circ}\text{C}$ to $^{\circ}\text{K}$, add 273 to the temperature in $^{\circ}\text{C}$.

$$T = 0^{\circ} + 273 = 273^{\circ}\text{K}$$

ΔH is given as 80 cal/g . When solving for the entropy, convert ΔH to cal/mole . There are 18 g of H_2O in one mole, thus the ΔH can be converted by multiplying it by 18 g/mole

$$\Delta H = 80 \text{ cal/g} \times 18 \text{ g/mole} = 1440 \text{ cal/mole}$$

It is now possible to solve for ΔS .

$$0 = \Delta H - T\Delta S \qquad \Delta H = 1440 \text{ cal/mole}$$

$$0 = 1440 \text{ cal/mole} - 273^{\circ}\text{K} \times \Delta S$$

$$- 1440 \text{ cal/mole} = - 273^{\circ}\text{K} \times \Delta S$$

$$\frac{- 1440 \text{ cal/mole}}{- 273^{\circ}\text{K}} = \Delta S = 5.27 \text{ cal/mole-}^{\circ}\text{K}$$

• PROBLEM 533

2 moles of hydrogen chloride are to be made from 1 mole each of hydrogen (H_2) and chlorine (Cl_2) at 25° and 1 atm. Calculate the ΔG for this chemical reaction.

STANDARD ENTHALPIES OF FORMATION,
 ΔH_f° AT $25^\circ C$

Compound	ΔH_f°
$NO(g)$	21.600 kcal/mole
$NO_2(g)$	8.091
$HF(g)$	-64.2
$HCl(g)$	-22.063
$HBr(g)$	-8.66
$HI(g)$	6.20

ENTROPIES OF CERTAIN SUBSTANCES
AT $25^\circ C$ AND 1 ATM

Substance	S (cal deg^{-1} mole $^{-1}$)
H_2	31.21 e.u.
O_2	49.00
Cl_2	53.31
N_2	45.77
HCl	44.61

Solution: Free energy (ΔG) may be expressed quantitatively as $\Delta G = \Delta H - T\Delta S$, where ΔH is the change in enthalpy, ΔS is the change in entropy, and T is temperature in degrees kelvin. Entropy measures the randomness of the system, while enthalpy is the heat content.

The reaction may be written $H_2(g) + Cl_2(g) \rightarrow 2HCl(g)$

Because free energy is defined in terms of ΔH , ΔS , and T , you can see these values to solve for ΔG . The table of standard enthalpies of formation, states that $\Delta H_{f,HCl}^\circ = -22.063$ kcal/mole. Two moles of HCl are formed, ΔH for the reaction is 2 moles $\times (-22,063)$ cal/mole. The ΔS is the final minus the initial entropy;

$$\begin{aligned} S_f - S_i &= S_{2HCl} - S_{H_2} - S_{Cl_2} \\ &= 2(44.61) - 31.21 - 53.31 \\ &= 4.70 \text{ cal } deg^{-1} \end{aligned}$$

Substituting into $\Delta G = \Delta H - T\Delta S$;

$$G = -44,123 \text{ cal} - (298)(4.70 \text{ eu}) = -45,523 \text{ cal.}$$

• PROBLEM 534

Determine the entropy difference between solid and liquid states for a substance melting at $100^\circ C$ and having a heat of fusion of 10,000 J/mol.

Solution: At the melting point, the liquid and solid are in equilibrium, which means the change in free energy

$\Delta H = T_{\text{Mp}} \Delta S$ or $T_{\text{Mp}} = \Delta H / \Delta S$. Because it takes 30.3 KJ/mole to melt NaCl, this must be the heat content or enthalpy. The change of entropy was given as $28.2 \text{ J mol}^{-1} \text{ deg}^{-1}$. You have therefore,

$$T_{\text{Mp}} = \frac{\Delta H}{\Delta S} = \frac{30,300 \text{ J/mole}}{28.2 \text{ J mole}^{-1} \text{ deg}^{-1}} = 1070^{\circ}\text{K}.$$

• PROBLEM 536

The density of ice at 0°C is $.917 \text{ g/cm}^3$ and has an entropy of $37.95 \text{ J mol}^{-1} \text{ deg}^{-1}$. The density of liquid water at this temperature is $.9998 \text{ g/cm}^3$ and has an entropy of $59.94 \text{ J mol}^{-1} \text{ deg}^{-1}$. Given these data, calculate the change of entropy ΔS , change of enthalpy ΔH , and the change of energy ΔE for the conversion of one mole of ice to liquid water at the normal melting point.

Solution: The solution of this problem involves the ability to relate energy, free energy, entropy, and enthalpy.

One must first calculate the change in entropy. You are told the entropies of both ice and water. The change in entropy is their difference or

$$\Delta S = S_{\text{liq}} - S_{\text{solid}} = 59.94 - 37.95 = 21.99 \text{ J mol}^{-1} \text{ deg}^{-1}.$$

To calculate the change in enthalpy at the melting point use the fact that $\Delta G = 0$ at the melting point of a substance. $\Delta G = \Delta H - T_{\text{Mp}} \Delta S$, where ΔH = change in enthalpy, T_{Mp} = temperature of melting point in Kelvin and ΔS = change in entropy. Since $\Delta G = 0$ at the melting point, $\Delta H = T_{\text{Mp}} \Delta S$.

From the previous calculation, you know ΔS . T_{Mp} is given. Thus,

$$\Delta H = T \Delta S$$

$$\Delta H = (273.15\text{K})(21.99 \text{ J mol}^{-1} \text{ deg}^{-1}) = 6010 \text{ J/mole}.$$

To calculate the ΔE for the conversion of one mole of ice to liquid water at the normal melting point, employ the equation $\Delta E = \Delta H - P \Delta V$, where P = pressure, V = volume. $\Delta H = 6010 \text{ J/mole}$. To determine $P \Delta V$, note that ΔV is $V_{\text{liq}} - V_{\text{solid}}$. You are given the densities of ice and water. Recalling that density = mass/volume, and that the molecular weight (mass) of ice or water is 18.015 grams/mole, you have for V_{liq} : $18.015/.9998$. For V_{solid} , you have $18.015/.917$. Therefore

$$\Delta V = V_{\text{liq}} - V_{\text{solid}} = \frac{18.015 \text{ g/mole}}{.9998 \text{ g/cm}^3} - \frac{18.015 \text{ g/mole}}{.917 \text{ g/cm}^3}$$

$$= -1.63 \text{ cm}^3/\text{mole} = -.00163 \text{ liter/mole.}$$

Since one is asked to make these calculations for 1 mole of H_2O , the change in volume, ΔV , equals $-.00163$ liter.

Because you are asked to calculate at the normal melting point, the pressure must be 1 atm (by definition, the "normal" is considered to be under a pressure of one atm.)

Therefore,

$$P\Delta V = (1.00 \text{ atm})(-.00163 \text{ liter}) = -.00163 \text{ liter-atm.}$$

Since 1 liter-atm equals 101.3 J, $P\Delta V$ is equal to


$$(-.00163 \text{ liter-atm})(101.3 \text{ J liter}^{-1} \text{ atm}^{-1}) = -.1651 \text{ J.}$$

Therefore,

$$\Delta E = \Delta H - P\Delta V = 6010 + .1651 = 6010.1651 \text{ J.}$$

• PROBLEM 537

Determine ΔS° , ΔH° , and ΔG° for the following reaction at 25°C .



Use the following table.

Substance	S°	ΔH_f°
CO	47.3	-26.4
Cl_2	53.3	0
COCl_2	69.1	-53.3

S° is expressed in cal/mole $^\circ\text{K}$ and
 ΔH_f° is in kcal/mole.

Solution: The change in entropy (ΔS°) or randomness of a reaction is obtained by subtracting the entropy (S°) of the reactants from the S° of the products. If more than

one mole of a compound reacts or is formed in a reaction, the S° should be multiplied by the appropriate stoichiometric coefficient. Here, one mole of CO reacts with one mole of Cl_2 to form one mole of COCl_2 .

$$\Delta S^\circ = S^\circ \text{ of } \text{COCl}_2 - (S^\circ \text{ of CO} + S^\circ \text{ of } \text{Cl}_2)$$

$$\Delta S^\circ = 69.1 - (47.3 + 53.3) = -31.5 \text{ cal/mole-}^\circ\text{K}$$

The change in enthalpy (ΔH°) of a reaction is determined by subtracting the enthalpy of formation (ΔH_f°) of the reactants from the ΔH_f° of the products. ΔH_f° of any pure element is always 0. When more than one mole of a compound either reacts or is formed in a reaction, the ΔH_f° must be multiplied by the corresponding stoichiometric coefficient when calculating ΔH° :

$$\Delta H^\circ = \Delta H_f^\circ \text{ of } \text{COCl}_2 - [\Delta H_f^\circ \text{ of CO} + \Delta H_f^\circ \text{ of } \text{Cl}_2]$$

$$\Delta H^\circ = -53.3 \text{ kcal} - (-26.4 \text{ kcal} + 0) = -26.9 \text{ kcal}.$$

The change in free energy (ΔG°) can be obtained if one remembers how ΔG° is related to ΔS° , T and ΔH° . Namely, $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, where ΔG° is the change in free energy, ΔH° is the change in enthalpy, ΔS° is the change in entropy and T is the absolute temperature. In this problem, one has already obtained ΔH° and ΔS° , and one can quickly obtain T . The absolute temperature can be found by adding 273 to the temperature in $^\circ\text{C}$.

$$T = 25 + 273 = 298^\circ\text{K}$$

In the previous part of this problem, one calculated ΔS° in cal/mole- $^\circ\text{K}$ and ΔH° in kcal/mole. To use ΔS° and ΔH° in the formula to find ΔG° , one must have them both in the same units. Thus, one should change the ΔS° term from cal to kcal by multiplying it by the conversion factor 1 kcal/1000 cal.

One can now obtain ΔG° .

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$\Delta G^\circ = -26.9 \text{ kcal} - (298\text{K})(-31.5 \text{ cal/}^\circ\text{K})(1 \text{ kcal/1000 cal})$$

$$= -26.9 + 9.4 = -17.5 \text{ kcal}.$$

• PROBLEM 538

Determine the free energy change, ΔG , for transforming liquid water at 100°C and 1 atm. to vapor at the same conditions. $\Delta H = 9720 \text{ cal}$ (molar enthalpy of vaporization).

Solution: This problem deals with the energies of reactions: enthalpy, entropy and free energy. The entropy of a system is a measure of the state of randomness. As

more thermal energy is added to a system, the more random it becomes. Entropy can be measured quantitatively. Entropy or $\Delta S = q_{\text{rev}}/T$, where q_{rev} is the amount of heat added to the system that can be recovered if the reaction were reversed. For example, the energy needed to melt ice can be recovered by freezing the liquid back to ice.

Enthalpy may be defined as the heat content of a system.

The concept of energy and entropy may be combined to define the driving force of a reaction. This driving force is referred to as Gibbs free energy (G) where $\Delta G = \Delta H - T\Delta S$, where ΔH is the enthalpy. This energy must be equal to the heat content of the system (ΔH) minus the energy necessary to increase the randomness, or entropy, of the same system.

Therefore, to solve this problem, you need to substitute into this equation the following values:

$$T = 100^{\circ}\text{C} = 373^{\circ}\text{K}$$

$$q = \Delta H = 9720 \text{ cal}$$

Therefore,

$$\Delta S = \frac{q}{T} = \frac{9720 \text{ cal}}{373^{\circ}\text{K}}$$

Now,

$$\Delta G = 9720 \text{ cal} - 373^{\circ}\text{K} \left(\frac{9720 \text{ cal}}{373^{\circ}\text{K}} \right) = 0.$$


The answer is zero.

The system can derive no usable energy to perform work from this change. For any system when $\Delta G=0$, the system is at equilibrium.

EQUILIBRIUM CALCULATIONS

• PROBLEM 539

Calculate the equilibrium constant for the following reaction at 25°C or 298°K



ΔH° for this reaction is $-17,889 \text{ cal}$.

Solution: At equilibrium the equilibrium constant of any

$$\log \frac{K_2}{K_1} = \frac{\Delta H^\circ}{19.15} \left(\frac{1}{T_1} - \frac{1}{T_2} \right),$$

where K_2 and K_1 are equilibrium constants, ΔH° = the standard enthalpy formation, and T_1 and T_2 are temperatures in Kelvin. Let $K_1 = 1.6 \times 10^{12}$ at $T_1 = 298^\circ\text{K}$; $T_2 = 373^\circ$; and $\Delta H = -113 \text{ J/mole}$. Substitute these values into the equation and solve for K_2 , which will be the solution to this problem. Thus

$$\log \frac{K_2}{1.6 \times 10^{12}} = \frac{-113,000}{19.15} \left(\frac{1}{298} - \frac{1}{373} \right)$$

Solving for K_2 , one obtains $K_2 = 1.7 \times 10^6$.

• PROBLEM 541

Given that $k = 8.85$ at 298°K and $k = .0792$ at 373°K , calculate the ΔH° for the reaction of the dimerization of NO_2 to N_2O_4 . Namely, $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$.

Solution: ΔH° is the standard enthalpy change, a measure of the heat content of the system. It is quantitatively related to the equilibrium constants of a system at different temperatures by the van't Hoff equation:

$$\log \frac{k_2}{k_1} = \frac{\Delta H^\circ}{19.15} \left(\frac{1}{T_1} - \frac{1}{T_2} \right),$$

where k_2 and k_1 are equilibrium constants, and T_1 and T_2 are temperatures in Kelvin. Given that $k_1 = 8.85$ at $T_1 = 298^\circ\text{K}$ and $k_2 = 0.0792$ at 373°K , one can find ΔH° by the substitution of these values in the equation. Thus,

$$\log \frac{.0792}{8.85} = \frac{\Delta H^\circ}{19.15} \left(\frac{1}{298} - \frac{1}{373} \right).$$

Solving for ΔH° , one obtains

$\Delta H^\circ = -58,200 \text{ J/mole}$ for the reaction.

• PROBLEM 542

Ammonia is synthesized commercially according to the equation



under a constant total pressure of 50 atm. At 350°C the equilibrium constant has been determined as 0.0278. The heat of reaction for this process is $\Delta H =$

- 11.04 Kcal/mole. Determine the value of equilibrium constant at 450°C.

Solution: For a reaction with equilibrium constant k_1 at absolute temperature T_1 and equilibrium constant k_2 at temperature T_2 ,

$$\log \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right),$$

where R is the gas constant.

Let $T_1 = 350^\circ\text{C} = 623^\circ\text{K}$, $k_1 = 0.0278$, and $T_2 = 450^\circ\text{C} = 723^\circ\text{K}$. $\Delta H^\circ = -11.04$ Kcal/mole = -11,040 cal/mole. Then

$$\log \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

$$\begin{aligned} \log \left(\frac{k_2}{0.0278} \right) &= \frac{11,040 \text{ cal/mole}}{2.303 \times 1.987 \text{ cal/mole-deg}} \left(\frac{1}{723^\circ\text{K}} - \frac{1}{623^\circ\text{K}} \right) \\ &= -0.535. \end{aligned}$$

Solving for k_2 ,

$$\log \left(\frac{k_2}{0.0278} \right) = -0.535$$


$$\frac{k_2}{0.0278} = 10^{-0.535} = 0.292$$

$$k_2 = 0.0278 \times 0.292 = 0.00812.$$

Thus, the value of the equilibrium constant at 450°C. is 0.00812.

• PROBLEM 543

Calculate the enthalpy change, ΔH° , for the reaction



given the equilibrium constants 4.08×10^{-4} for a temperature of 2000°K and 3.60×10^{-3} for a temperature of 2500°K.

Solution: The effect of temperature on chemical equilibrium is determined by ΔH° (enthalpy or heat content); over moderate ranges in temperature, ΔH° is relatively independent of temperature.

If, as in the case of this problem, ΔH° is independent of temperature then,

$$(1) \Delta G^\circ = -RT \ln k, \text{ where}$$

ΔG° = standard free energy change,

R = gas constant,

T = absolute temperature

and k the equilibrium constant.

$$(2) \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$(3) \Delta H^\circ - T\Delta S^\circ = -RT \ln K \quad \text{or}$$

$-\left(\frac{\Delta H^\circ - T\Delta S^\circ}{RT}\right) = \ln k$, where ΔS° = the change in entropy or randomness of system.

$$(4) \frac{-\Delta H^\circ}{RT} + \frac{\Delta S^\circ}{R} = \ln k$$

For two different temperatures, T_1 and T_2 , equation (4) becomes

$$\ln k_2 - \ln k_1 = \left(\frac{-\Delta H^\circ}{RT_2} + \frac{\Delta S^\circ}{R}\right) - \left(\frac{-\Delta H^\circ}{RT_1} + \frac{\Delta S^\circ}{R}\right) \quad \text{or}$$

$$\ln \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{RT_2} + \frac{\Delta H^\circ}{RT_1}$$

$$\ln \frac{k_2}{k_1} = \frac{\Delta H^\circ (T_2 - T_1)}{RT_2 T_1} \quad \text{or, finally,}$$

$$\log \frac{k_2}{k_1} = \frac{\Delta H^\circ (T_2 - T_1)}{(2.303) R T_2 T_1}$$

$$\text{Thus, } \log \frac{k_{2500^\circ\text{K}}}{k_{2000^\circ\text{K}}} = \log \frac{3.60 \times 10^{-3}}{4.08 \times 10^{-4}}$$

$$= \frac{\Delta H^\circ (2500 - 2000)^\circ\text{K}}{(2.303)(1.987 \text{ cal}^\circ\text{K}^{-1}\text{mole}^{-1})(2500)^\circ\text{K}(2000)^\circ\text{K}}$$

$$\log 8.82 = .945 = \frac{\Delta H^\circ (500^\circ\text{K})}{(2.303)(1.987)(5 \times 10^6) \text{ cal}^\circ\text{K}^{-1}\text{mole}^{-1} \text{ }^\circ\text{K}^2}$$

$$\frac{(0.945)(2.303)(1.987)(5 \times 10^6)}{(500) \text{ mole cal}^{-1}} = \Delta H^\circ$$

$$\Delta H^\circ = 43,240 \text{ cal/mole.}$$

For the reaction $\text{PbSO}_4(\text{s}) \rightarrow \text{Pb}^{2+} + \text{SO}_4^{2-}$, $\Delta H = + 2990 \text{ cal/mole}$. Will the solubility of PbSO_4 increase or decrease with increasing temperature? $K_{\text{sp}} = 1.8 \times 10^{-8}$ at 25°C . Find its k_{sp} at 55°C .

Solution: Whether the solubility of a salt increases or decreases with an increase in temperature can be determined by an investigation of its ΔH , the enthalpy change. The formula for the temperature dependence on k is

$$\log k = \frac{\Delta H^\circ}{2.303 RT} + \frac{\Delta S^\circ}{2.303 R}.$$

A positive ΔH° suggests that solubility increases with increasing temperature. When ΔH° is positive, the first term remains a negative value. When T increases, the first term becomes a smaller negative number. Thus, k increases. Since k measures solubility, the solubility increases. To find the k_{sp} of PbSO_4 at 55°C , use the fact that

$$\log \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right),$$

where k_2 = solubility constant at the temperature in Kelvin (Celsius plus 273°) of T_2 ; k_1 = solubility constant at a temperature of T_1 . Thus, if the k_{sp} at one temperature is known, the k_{sp} at another temperature can be found given ΔH° . Let $T_2 = 55^\circ\text{C}$ or 328°K and $T_1 = 298^\circ\text{K}$. (25°C). Thus

$$\log \frac{k_{328}}{k_{298}} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{328} - \frac{1}{298} \right)$$

$$\log \frac{k_{328}}{1.8 \times 10^{-8}} = \frac{-(2990 \text{ cal/mole})(-3.07 \times 10^{-4}/\text{K})}{(2.303)(1.987 \text{ cal/mole K})}$$


$$= 0.200$$

$$k_{328} = (1.8 \times 10^{-8})(10^{0.200})$$



$$= (1.8 \times 10^{-8})(1.58)$$

$$= 2.8 \times 10^{-8}.$$

The equilibrium constant and standard Gibbs free energy change for ammonia synthesis at 400°C or 673°K are 1.64×10^{-4} and 11,657 cal/mole, respectively. The equation for this reaction is



Calculate the equilibrium constants and standard free energy changes for



Solution: This problem illustrates the necessity of knowing the balanced chemical equation in order to determine the numerical value of the equilibrium constant.

For ideal gases, the equilibrium constant, K_p , is the product of the partial pressure of the product gases, each raised to the power that corresponds with its stoichiometric coefficient in the balanced equation, divided by the product of the partial pressure of the reacting gases, each raised to the power that corresponds with its stoichiometric coefficient in the balanced equation.

The quantity ΔG° is the standard change in Gibbs free energy. It is the difference between the Gibbs free energy of the products and the Gibbs free energy of the reactants. The Gibbs Free Energy and the equilibrium constant are related in the equation

$$\Delta G = -RT \ln K_p$$

where R is the gas constant, T the absolute temperature and K_p the equilibrium constant calculated using partial pressures.

The equilibrium constant for reaction 1 is

$$K_1 = \frac{P_{\text{NH}_3}^2}{P_{\text{N}_2} P_{\text{H}_2}^3} = 1.64 \times 10^{-4}$$

Thus, for (2) $\frac{1}{2}\text{N}_2(\text{g}) + \frac{3}{2}\text{H}_2(\text{g}) = \text{NH}_3(\text{g})$


$$K_2 = \frac{P_{\text{NH}_3}}{P_{\text{N}_2}^{1/2} P_{\text{H}_2}^{3/2}} = K_1^{1/2} = (1.64 \times 10^{-4})^{1/2} = 1.28 \times 10^{-2}$$

By definition, at equilibrium,

$$\Delta G^\circ = -RT \ln K_2$$

$$= - (1.987 \text{ cal}^\circ\text{K}^{-1} \text{ mole}^{-1}) (673^\circ\text{K}) (2.303) \times \log(1.28 \times 10^{-2})$$

$$= 5829 \text{ cal/mole.}$$



$$K_3 = \frac{P_{\text{N}_2} P_{\text{H}_2}^3}{P_{\text{NH}_3}^2} = \frac{1}{K_1} = \frac{1}{1.64 \times 10^{-4}} = 0.6098 \times 10^4$$


$$\Delta G_3^\circ = -RT \ln K_3 = - (1.987 \text{ cal}^\circ\text{K}^{-1} \text{ mole}^{-1}) (673^\circ\text{K}) (2.303) \times \log (0.6098 \times 10^4)$$

$$= - 11,657 \text{ cal/mole.}$$

If a reaction is reversed, the equilibrium constant becomes the reciprocal of that for the first reaction; the standard Gibbs free energy change has the same magnitude but the opposite sign.


• PROBLEM 546

Calculate ΔG° for the reaction



at $400^\circ\text{C} = 673^\circ\text{K}$ using K_c . The value of K_p for this reaction is 1.64×10^{-4} . Interpret the different values of ΔG° calculated from K_p and K_c .

Solution: Given the equilibrium reaction:



there are two constants that measure the equilibrium: K_c relates the concentrations.

$$(ii) \quad K_c = \frac{[C]^c [D]^d \dots}{[A]^a [B]^b \dots}$$

K_p relates the pressures

$$(iii) \quad K_p = \frac{P_C^c P_D^d \dots}{P_A^a P_B^b \dots}$$

To derive K_c from K_p substitute for each pressure $P = cRT$, where R is the gas constant; T = absolute tem-

where R = universal gas constant and T = temperature in Kelvin. Solving for K , you obtain:

$$(ii) \quad \ln K = \frac{\Delta G_{490}^{\circ}}{-RT}$$

In this problem, $R = 8.31 \text{ J mol}^{-1} \text{ deg}^{-1}$, and $T = 490^{\circ}\text{C} = (490^{\circ} + 273^{\circ})\text{K} = 763^{\circ}\text{K}$. ΔG_{490}° is the amount of free energy needed for the reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2 \text{HI}$, i.e., the amount of free energy needed for the production of 2 moles of HI at 490°C . Given that -12.1 kJ will produce 1 mole of HI at 490°C , it follows that

$$\Delta G_{490}^{\circ} = 2(-12.1 \text{ kJ/mole}) = -24.2 \text{ kJ/mole}$$

$$= -24,200 \text{ J/mole.}$$

Substituting these values in equation (ii), you obtain


$$(iii) \quad \ln K = - \frac{\Delta G^{\circ}}{RT} = \frac{-(-24,200 \text{ J/mole})}{(8.31 \text{ J mol}^{-1} \text{ deg}^{-1})(763 \text{ deg})}$$

$$(iv) \quad \ln K = 3.82$$

$$(v) \quad K = 46.$$

• PROBLEM 549

Phosgene is formed at 25°C according to the equation



Calculate the free energy change ΔG° and the equilibrium constant k for this reaction.

Solution: The free energy change for the reaction is the difference between the sum of the free energies of formation of the products and the sum of the free energies of formation of the reactants.

The free energies of formation of CO(g) , $\text{Cl}_2(\text{g})$ and $\text{COCl}_2(\text{g})$ are

$$\Delta G_{\text{CO}}^{\circ} = -32.8 \text{ Kcal/mole}$$

$$\Delta G_{\text{Cl}_2}^{\circ} = 0 \text{ Kcal/mole}$$

$$\Delta G_{\text{COCl}_2}^{\circ} = -50.3 \text{ Kcal/mole.}$$

$$\text{Hence,} \quad \Delta G^{\circ} = \Delta G_{\text{COCl}_2}^{\circ} - \left(\Delta G_{\text{CO}}^{\circ} + \Delta G_{\text{Cl}_2}^{\circ} \right)$$

$$\begin{aligned}
 &= -50.3 \text{ Kcal/mole} - (-32.8 \text{ Kcal/mole} + \\
 &\quad 0 \text{ Kcal/mole}) \\
 &= -17.5 \text{ Kcal/mole} = -17,500 \text{ cal/mole.}
 \end{aligned}$$

The equilibrium constant is calculated from

$$\Delta G^\circ = -2.303 RT \log k,$$

where R is the gas constant and T is the absolute temperature ($T = 25^\circ\text{C} + 273 = 298^\circ\text{K}$). Then,


$$\begin{aligned}
 \Delta G^\circ &= -2.303 RT \log k \\
 -17,500 \text{ cal/mole} &= -2.303 \times 1,987 \text{ cal/mole-deg} \times 298^\circ\text{K} \\
 &\quad \times \log k.
 \end{aligned}$$

Solving for k,

$$\begin{aligned}
 \log k &= \frac{-17,500 \text{ cal/mole}}{-2.303 \times 1,987 \text{ cal/mole-deg} \times 298^\circ\text{K}} = 12.8, \\
 k &= 10^{12.8} = 6 \times 10^{12}.
 \end{aligned}$$

• PROBLEM 550

Sulfur exists as S_2 vapor at temperatures between 700°C and 1500°C . At 1473 K it combines with hydrogen according to the equation



At 750°C the equilibrium constant is 1.07×10^2 and at 1200°C it is 4.39. Determine the heat of reaction in the temperature range 750°C to 1200°C , and the change in free energy at each of these temperatures.

Solution: The change in free energy ΔG is determined from the equation

$$\Delta G^\circ = -2.303 RT \log k,$$

where R is the gas constant, T the absolute temperature, and k the equilibrium constant. From this equation and the expression

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ,$$

where ΔH° is the change in the heat of reaction and ΔS° the change in entropy, a relation between ΔH and T will be found.

$$\text{At } T = 750^\circ\text{C} = 1023^\circ\text{K}, \quad k = 1.07 \times 10^2 \quad \text{and}$$

$$\Delta G^\circ = - 2.303 RT \log k$$

$$= - 2.303 \times 1.987 \frac{\text{cal}}{\text{mole-deg}} \times 1023^\circ\text{K} \times \log (1.07 \times 10^2)$$

$$= - 9500 \text{ cal/mole}$$

$$= - 9.5 \text{ Kcal/mole.}$$

At $T = 1200^\circ\text{C} = 1473^\circ\text{K}$, $k = 4.39$ and

$$\Delta G^\circ = - 2.303 RT \log k$$

$$= - 2.303 \times 1.987 \frac{\text{cal}}{\text{mole-deg}} \times 1473^\circ\text{K} \times \log (4.93)$$

$$= - 4330 \text{ cal/mole}$$

$$= - 4.33 \text{ Kcal/mole.}$$

To find an expression relating ΔH to T , consider the following two equations:

$$\Delta G^\circ = - 2.303 RT \log k, \text{ and } \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

Setting these equal gives

$$- 2.303 RT \log k = \Delta H^\circ - T\Delta S^\circ$$

$$\text{or, } \log k = \frac{-\Delta H^\circ}{2.303 RT} + \frac{\Delta S^\circ}{2.303 R}$$

For two different temperatures, T_1 and T_2 with equilibrium constants k_1 and k_2 , respectively,

$$\log k_1 = \frac{-\Delta H^\circ}{2.303 RT_1} + \frac{\Delta S^\circ}{2.303 R} \quad \text{and}$$

$$\log k_2 = \frac{-\Delta H^\circ}{2.303 RT_2} + \frac{\Delta S^\circ}{2.303 R}$$

Subtracting the first equation from the second,

$$\log k_2 - \log k_1 = \left(\frac{-\Delta H^\circ}{2.303 RT_2} + \frac{\Delta S^\circ}{2.303 R} \right) - \left(\frac{-\Delta H^\circ}{2.303 RT_1} + \frac{\Delta S^\circ}{2.303 R} \right),$$

$$\text{or, } \log \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right).$$

Let $T_1 = 1023^\circ\text{K}$, $k_1 = 1.07 \times 10^2$, $T_2 = 1473^\circ\text{K}$, and $k_2 = 4.39$. Then

$$\log \frac{k_2}{k_1} = \frac{-\Delta H^\circ}{2.303 R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

$$\log \frac{4.39}{1.07 \times 10^2} = \frac{-\Delta H^\circ}{2.303 \times 1.987 \text{ cal/mole-deg}} \left(\frac{1}{1473^\circ\text{K}} - \frac{1}{1023^\circ\text{K}} \right)$$

$$\log \frac{4.39}{1.07 \times 10^2} = \frac{-\Delta H^\circ}{2.303 \times 1.987 \text{ cal/mole-deg}} (0.000679^\circ\text{K}^{-1} - 0.000978^\circ\text{K}^{-1})$$

or,

$$\begin{aligned} \Delta H^\circ &= \frac{\log (4.39/107)}{2.303 \times 1.987 \text{ cal/mole-deg}} (0.000978^\circ\text{K}^{-1} - 0.000679^\circ\text{K}^{-1}) \\ &= -21,300 \text{ cal/mole} \\ &= -21.3 \text{ Kcal/mole.} \end{aligned}$$

Thus, the heat of reaction in the temperature range 1023°K to 1473°K (750°C to 1200°C) is - 21.3 Kcal/mole.

• PROBLEM 551

In the human body, the enzyme phosphoglucomutase catalyzes the conversion of glucose-1-phosphate into glucose-6-phosphate:



At 38°C, the equilibrium constant, k , for this reaction is approximately 20. Calculate the free energy change, ΔG° , for the equilibrium conversion. Calculate the free energy change ΔG for the nonequilibrium situation in which [glucose-1-phosphate] = 0.001 M and [glucose-6-phosphate] = 0.050 M.

Solution: The nonequilibrium free energy change ΔG is related to the standard free energy change ΔG° by

$$\Delta G = \Delta G^\circ + 2.303 RT \log k,$$

where R is the gas constant and T the absolute temperature.

At equilibrium, $\Delta G = 0$, hence

$$\Delta G^\circ = -2.303 RT \log k.$$

Thus, for the equilibrium conversion with $k = 20$ and $T = 38^\circ\text{C} = 311^\circ\text{K}$.

$$\Delta G^\circ = -2.303 RT \log k$$

$$\begin{aligned}
 &= - 2.303 \times 1.987 \text{ cal/mole-deg} \times 311^\circ\text{K} \times \log 20 \\
 &= - 1850 \text{ cal/mole.}
 \end{aligned}$$

The equilibrium constant for the conversion

glucose-1-phosphate \rightleftharpoons glucose-6-phosphate

$$\text{is } k = \frac{[\text{glucose-6-phosphate}]}{[\text{glucose-1-phosphate}]}$$

In the case where $[\text{glucose-6-phosphate}] = 0.050 \text{ M}$ and $[\text{glucose-1-phosphate}] = 0.001 \text{ M}$


$$k_n = \frac{[\text{glucose-6-phosphate}]}{[\text{glucose-1-phosphate}]} = \frac{0.050 \text{ M}}{0.001 \text{ M}} = 50,$$

where the subscript "n" has been added to distinguish this ratio from the equilibrium constant. Then

$$\begin{aligned}
 \Delta G &= \Delta G^\circ + 2.303 RT \log k_n \\
 &= - 1850 \text{ cal/mole} + (2.303)(1.987 \text{ cal/mole-deg}) \\
 &\quad (311^\circ\text{K})(\log 50) \\
 &= - 1850 \text{ cal/mole} + 2420 \text{ cal/mole} \\
 &= + 570 \text{ cal/mole.}
 \end{aligned}$$

• PROBLEM 552

Calculate the equilibrium constant at 25°C for the reaction:



The heat formation of SO_3 at 25°C is $- 94.45 \text{ Kcal/mole}$ and the standard molar entropy changes for S , O_2 , and SO_3 at 25°C are 7.62 , 49.0 and $61.24 \text{ cal/mole}^\circ\text{K}$, respectively.

Solution: One is given the values for ΔH and S which can be related by the equation

$$\Delta G^\circ = \Delta H^\circ - T\Delta S,$$

where ΔG is the change in free energy, ΔH is the change in the heats of formation, T is the absolute temperature, and ΔS is the change in entropy.

After calculating ΔG° , one can solve for the equilibrium constant (K) by using the equation

Determine the equilibrium constant for the following reaction at 25°C:



Solution: The equilibrium constant, k , is related to the cell potential, ΔE° , by means of the following two formulas:



$$\Delta G^\circ = -n F \Delta E^\circ, \quad \text{and} \quad \Delta G^\circ = -2.303 RT \log k,$$

where n is the number of electrons transferred, F the Faraday constant, R the gas constant, ΔG° the free energy, and T the absolute temperature. ΔE° will be determined from the half-cell potentials, E° .

First, the cell potential is determined from the half-cell reactions:



Adding the two half-cell reactions and their respective values for E° ,



The cell potential for the reaction is therefore 2.23 v.

Equating the two expressions

$$\Delta G^\circ = -n F \Delta E^\circ$$

$$\text{and } \Delta G^\circ = -2.303 RT \log k$$

$$\text{gives } -2.303 RT \log k = -n F \Delta E^\circ,$$

$$\text{or, } \log k = \frac{n F \Delta E^\circ}{2.303 RT}.$$

Since two electrons are transferred from Mg(s) to Sn^{2+} , $n = 2$. The absolute temperature is $T = 25^\circ\text{C} + 273 = 298^\circ\text{K}$. Hence

$$\begin{aligned} \log k &= \frac{n F \Delta E^\circ}{2.303 RT} \\ &= \frac{(2)(2.30609 \times 10^5 \text{ cal/mole-v})(2.23 \text{ v})}{(2.303)(1.987 \text{ cal/mole-deg})(298^\circ\text{K})} = 75.6, \end{aligned}$$

$$\begin{aligned}
 \Delta G_{\text{NO}_2}^\circ &= \frac{1}{2} \left(\Delta G^\circ + 2\Delta G_{\text{NO}}^\circ \right) \\
 &= \frac{1}{2} (-69.6 + 2 \times 86.6) \\
 &= 51.8 \text{ kJ/mole.}
 \end{aligned}$$

• PROBLEM 556

Calculate the change in Gibbs free energy (ΔG) for the production of 2 $\text{NO}_2(\text{g})$ at 1 atm from $\text{N}_2\text{O}_4(\text{g})$ at 10 atm at 25°C . The standard Gibbs free energy for the reaction



is + 1161 cal at 25°C .

Solution: If ΔG° has a positive value, the reactants in their standard states will not react spontaneously to give products in their standard states. By increasing the pressures or concentrations of the reactants or decreasing the pressures or concentrations of the products, however, it may be possible to make ΔG negative so that the reaction will proceed spontaneously.

To calculate ΔG under nonstandard conditions, the following formula is used:

$$\Delta G = \Delta G^\circ + RT \ln \pi p_i^{v_i}, \quad \text{where } R = \text{gas constant,}$$

T = absolute temperature (Celsius + 273), and $\pi p_i^{v_i}$ represents the product of the partial pressures of the component gases, each raised to the power that corresponds with its stoichiometric coefficient in the balanced equation. Thus,

$$\begin{aligned}
 \Delta G &= \Delta G^\circ + RT \ln \frac{p_{\text{NO}_2}^2}{p_{\text{N}_2\text{O}_4}} \\
 &= +1161 + (1.987 \text{ cal}^\circ\text{K}^{-1} \text{ mole}^{-1})(298^\circ\text{K}) \\
 &\quad (2.303) \log (1 \text{ atm})^2 / (10 \text{ atm}) \\
 &= 1161 - 1364 = -203 \text{ cal.}
 \end{aligned}$$

The production of $\text{NO}_2(\text{g})$ at 1 atm from $\text{N}_2\text{O}_4(\text{g})$ at 10 atm is a spontaneous process due to the fact that a negative ΔG is obtained. Thus, a continuous process would be thermodynamically feasible if N_2O_4 was maintained at 10 atm and NO_2 was withdrawn by some method so that its partial pressure was maintained at 1 atm.

CHAPTER 16

ELECTROCHEMISTRY


Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 539 to 579 for step-by-step solutions to problems.

Electrochemistry is that branch of chemistry which deals with reactions that involve the production and consumption of electrons. Hence electric currents are consumed or produced in electrochemical reactions. Normally the required electric current is introduced into the reactor, called a cell, through electrodes.

The conduction of electric currents in solutions of electrolytes is effected by the motion of ions within the solution. Positively charged ions migrate toward the anode (the negative electrode) while negatively charged ions migrate toward the cathode (the positive electrode). Another way to identify the difference between the cathode and anode is to remember that reduction reactions always occur at the cathode and oxidation reactions at the anode.

In electrochemical reactions, the charge must be balanced as though it were an elemental species in the reaction, and overall, the electrons produced in one part of the reaction must be consumed in another part. If this were not so, tremendous electric charges would develop for even small amounts of reaction. A typical example is the industrially important electrolysis of molten sodium chloride to produce chlorine gas and sodium metal.

The electrolyte contains sodium (Na^+) and chloride (Cl^-) ions. The chloride ions react at the anode and sodium ions at the cathode according to the equations



Since the same number of electrons enter the electrolyte at the cathode and leave at the anode, the overall reaction is obtained by multiplying the second half-reaction by two and adding the two reactions to obtain



Note that no electrons appear in the overall reaction. It is significant that electrons cannot be stored in sufficient numbers to permit the appreciable independent conduction of electrochemical half-reactions.

The equivalent weight is defined for electrochemical reactions in a manner analogous to that outlined in Chapter 8 for equivalent weights of solution. The equivalent weight of a species is the molecular (or atomic) weight divided by the charge or number of electrons which react. Michael Faraday, in 1832, observed that the amount of a substance undergoing oxidation or reduction at an electrode in an electrochemical cell was directly proportional to the amount of electric current that passed through the cell. This observation is known today as Faraday's Law and the unit of the "faraday" is the charge equal to that in one mole (6.022×10^{23}) of electrons. It is a huge charge, equal to 96,487 coulombs (c), or 96,487 ampere seconds (A-sec).

Electrochemical reactions are often analyzed as oxidation-reduction couples, abbreviated as redox reactions. The key to the balancing of redox reactions is to balance the charge as well as the elements on each side of the reaction. An example of balancing such a reaction is shown later for the displacement of Cu^{+2} ions from solution by iron metal (Equation 16-4).


An important concept in electrochemistry is that of the half-cell standard potential or emf. Table 1 lists a sample of important half-cell reactions and their standard potential. This is the potential produced by the half-cell reaction when balanced against a standard hydrogen electrode with all species present at standard (1 molar) concentrations.

Step-by-Step Solutions to Problems in this Chapter, “Electrochemistry”

CONDUCTION

• PROBLEM 558

Design an experiment which demonstrates that both positive and negative ions move in electrolytic conduction.



Solution: The accompanying figure illustrates the set up of an experiment which will demonstrate the migration of both positive and negative ions in electrolytic conductivity. A U-tube is initially half filled with a deep purple solution of $\text{Cu}(\text{MnO}_4)_2$ in water. The color of the blue hydrated Cu^{2+} ions is masked by the purple of the MnO_4^- ions. A colorless aqueous solution of HNO_3 floats on top of the $\text{Cu}(\text{MnO}_4)_2$ solution in each arm of the U-tube. If an electric field is established for a period of time across the solution by 2 electrodes, the blue color, characteristic of hydrated Cu^{2+} ions, moves into the region marked A, suggesting a migration of positive ions toward the negative electrode. At the same time, the purple color, characteristic of MnO_4^- , moves into the B region, indicating a migration of negative ions toward the positive electrode. Thus, there is empirical evidence of the migration of both positive and negative ions.

Would you expect the mobility of an electrolytic ion to increase or decrease for each of the following? (a) increasing the size of the ion; (b) increasing charge on the ion; (c) increasing temperature and (d) placing it in a solvent of lower viscosity.

Solution: To see how the mobility of the electrolytic ion changes when you alter these parameters, you must know what happens to these ions when an electric field is applied. The ions are free to move randomly around in solution before an electric field is applied. However, when the field is present, the positive ions experience a force in one direction, while the negative ions experience a force in the opposite direction. With this in mind, you proceed as follows:

(a) when you increase the size of the ion, you would anticipate a decrease in mobility. Remember, the ion must move when the electric current is applied; additional mass and volume, therefore, inhibits this movement, since there is greater resistance from the solvent molecules.

(b) If you increase the charge on the ion, you expect mobility to increase due to the fact that the force which results in movement is directly proportional to the attraction of the ions to the poles. Thus, by increasing the charge, you increase this force, which, in turn, increases movement.


(c) When you increase the temperature, you increase the mobility of the ions. By increasing the temperature, the average kinetic energy of the ion increases. Kinetic energy is a measure of movement.

(d) A solution of lower viscosity is expected to have an increase in mobility. This stems from the fact that viscosity measures internal resistance to flow. Thus, if the viscosity decreases, there is less resistance to flow, which means that the ions can move more freely.

EQUIVALENT WEIGHT


Calculate the theoretical quantity of chlorine obtainable by the electrolysis of 2.0 kg of a 20% sodium chloride solution. What other products would be obtained and what would be the weight of each?

Solution: The equation for this reaction is as follows:



One mole of Cl_2 will be formed from every 2 moles of NaCl reacted.

Therefore, to find the quantity of Cl_2 formed one must first know the amount of NaCl reacted. This can be determined by solving for the weight of NaCl in the solution and dividing this number by the mole-



cular weight to find the number of moles present. Since the solution is 20% NaCl, it means that 20% of its weight is made up by NaCl.

$$\text{weight of NaCl} = 0.20 \times 2.0 \text{ kg} = 0.4 \text{ kg}.$$

Therefore, there are 0.4 kg or 400 g of NaCl present.

One can now solve for the number of moles present by dividing 400g by the molecular weight (MW of NaCl = 58.5)

$$\text{no. of moles} = \frac{400\text{g}}{58.5\text{g/mole}} = 6.84 \text{ moles}.$$

From the equation one sees that $\frac{1}{2}$ of this amount is equal to the number of moles of Cl_2 formed.

$$\text{no. of moles of } \text{Cl}_2 = \frac{1}{2} \times 6.84 \text{ moles} = 3.42 \text{ moles}.$$

The weight of this number of moles of Cl_2 will be equal to the molecular weight of Cl_2 times the number of moles.

$$(\text{MW of } \text{Cl}_2 = 71.0)$$

$$\begin{aligned} \text{weight of } \text{Cl}_2 &= 3.42 \text{ moles} \times 71.0\text{g/mole} = 242.8\text{g} \\ &\text{of } \text{Cl}_2 \end{aligned}$$

From the reaction one can also see that if 2 moles of NaCl react, 2 moles of NaOH are formed. Here, 6.84 moles of NaCl is reacted, therefore, 6.84 moles of NaOH are formed. The weight of this quantity is equal to the number of moles \times the molecular weight. (MW of NaOH = 40).

$$\text{weight of NaOH} = 40\text{g/mole} \times 6.84 \text{ moles} = 273.6\text{g}.$$

It is also seen from the reaction, that if 2 moles of NaCl are reacted, 1 mole of H_2 is formed. Thus, if 6.84 moles of NaCl are reacted, 3.42 moles of H_2 are formed. The weight of H_2 can then be found. (MW of H_2 = 2).

$$\text{weight of } \text{H}_2 = 2\text{g/mole} \times 3.42 \text{ moles} = 6.84\text{g}.$$

• PROBLEM 561

.0324 Faradays (F) liberated .651g of Calcium. What is the atomic weight of Calcium?

Solution: A faraday may be defined as the number of coulombs or charges of electricity that liberate one gram-equivalent weight of an element in solution. Note, the equivalent weight of a substance is the amount of substance which 1 mole of electrons can oxidize or reduce into a neutral species.

Calcium has an equivalent weight of $\frac{1}{2}$ its atomic weight, since its oxidation state is +2. In other words, one mole of electrons will neutralize only half of the calcium ions present, since each ion needs two electrons. Thus, if you can find the gram-equivalent weight of calcium, you can find its atomic weight by multiplying it by two. However, you can determine the number of gram-equivalents given the number of faradays of electricity used. You are told that .0324F deposits .651g. By definition, 1 F deposits the gram-equivalent weight. You have, then, the following proportion:

$$\frac{.651 \text{ g Ca}}{.0324 \text{ F}} = \frac{\text{g-eq. wt. of Ca}}{1 \text{ F}} = 20.1$$

Hence, 20.1g is the gram-equivalent of calcium. Thus, the atomic weight is found to be $2 \times 20.1 = 40.2$.

• PROBLEM 562

Two electrolytic cells were placed in series. One was composed of AgNO_3 and the other of CuSO_4 . Electricity was passed through the cells until 1.273g of Ag had been deposited. How much copper was deposited at the same time?

Solution: To find out how much copper was deposited, you need to know how many faradays were contained in the electricity that passed through the solution. One faraday, by definition, deposits one gram-equivalent of material. The equivalent weight of a substance is that amount of the substance which 1 mole of electrons (1 faraday) can oxidize or reduce to a neutral species.

You are told that 1.273g of Ag are deposited. From this, you can compute the number of faradays. The oxidation state of Ag is +1. Thus, the reaction that deposited Ag had to be $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}$. Thus, 1 mole of electrons yields 1 gram-equivalent weight of Ag. In this case, the gram-equivalent weight equals the atomic weight, since 1 mole of Ag is produced with 1 mole of electrons. Recalling the definition of a faraday, you can say that 1 faraday can deposit 107.87g of Ag (the atomic weight). You are told, however, that only 1.273g has been deposited. Thus, the following proportion can be used to find the number of faradays used:

$$\frac{1.273 \text{ g Ag}}{X \text{ Faradays}} = \frac{107.87 \text{ g Ag}}{1 \text{ Faraday}}$$

Solving, $X = .01180$ Faradays. Thus, the amount of faradays passed is .01180F, which means the amount of copper deposited depends on this amount of electricity.

To deposit copper, you must perform the reaction $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$ you see that it takes 2 moles of electrons to deposit one mole of copper. Thus, one faraday can only deposit one half of a mole copper. Thus, 1 faraday reduces $(\frac{1}{2})(63.54) = 31.77$ grams. You have, however, only .01180 faradays. Thus, the amount of Cu deposited can be found using

Two electrochemical cells were fitted with inert platinum electrodes and connected in series. The first cell contained a solution of a metal nitrate; the second cell, metal sulfate. After operation of the cell for a prescribed period of time, it was observed that 675 mg of the metal had been deposited in the first cell and 76.3 ml of hydrogen gas (25° and 1 atm) evolved in the second. Determine the equivalent weight of the metal.

Solution: Equivalent weight may be defined as that amount of substance which one mole of electrons will reduce or oxidize.

Since the electrodes are connected in series, the same current passes through both cells. Thus, equal numbers of equivalents of the metal (m) and hydrogen must be liberated. The cathode reactions for the two cells are: first cell: $M^{n+} + ne^- \rightarrow M(s)$; second cell: $2H^+ + 2e^- \rightarrow H_2(g)$, where n is the number of electrons needed to form the M(s) from the metal ion. One can solve for the number of moles of H_2 formed by use of the Ideal Gas Law.

$$n = \frac{PV}{RT}$$

where n is the number of moles, R is the gas constant, 0.082 liter-atm/mole^oK, T is the absolute temperature, V is the volume and P is the pressure. One is given the temperature in ^oC, to convert to ^oK add 273 to the temperature in ^oC.

$$T = 25^\circ + 273 = 298^\circ K.$$

Now, solving for n:

$$n = \frac{(1 \text{ atm})(.0763 \text{ liters})}{(0.082 \text{ liter-atm/mole}^\circ K)(298^\circ K)} = 3.12 \times 10^{-3} \text{ moles.}$$

From the half equations, one can see that there are two equivalents for each mole of H_2 formed. Thus, there must be $2 \times (3.12 \times 10^{-3})$ moles of equivalents² of the metal present.

Hence, equivalents of M = $2 \times 3.12 \times 10^{-3} = 6.24 \times 10^{-3}$ equiv. One is given that 675mg or .675 g of metal are formed, thus, in .675 g of the metal there are 6.24×10^{-3} equiv. The weight of one equivalent can be found by dividing .675 g by 6.24×10^{-3} equiv.


$$\text{equiv. wt.} = \frac{.675g}{6.24 \times 10^{-3} \text{ equiv.}} = 108.2g/\text{equiv.}$$

In an electrolytic cell, a liter of a 1M aqueous solution of MnO_4^- is reduced at the cathode. Determine the number of faradays required⁴ for each of the following to be made;


- a solution that is .01 M MnO_4^{2-} ;
- 1 gram of MnO_2 ;
- 1 gram-equivalent of Mn metal.

Solution: For each part, you need to write the reaction that occurs. The masses of substances produced during the process are proportional to

Balance the following reaction in acidic aqueous solution:




Solution: Reactions in which electrons are transferred from one atom to another are known as oxidation-reduction reactions or as redox reactions. To balance this type of reaction, you want to conserve charge and matter, i.e., one side of the equation must not have an excess of charge or matter. To perform this balancing, you need to (1) Balance charge by adding electrons. (2) Balance oxygen by adding water. (3) Balance atoms (of hydrogen) by adding H^+ (in acid) or OH^- (in base). These three rules will balance the redox equation. These rules apply to balancing only the half-reactions. The overall reaction, the sum of these, will be balanced by their addition. Proceed as follows; Fe^{2+} goes to Fe^{3+} . It lost an electron thus, it's the oxidation half reaction. To balance charge, add e^- . Thus,





The reduction must be $\text{ClO}_3^- \rightarrow \text{Cl}^-$. Chlorine changes oxidation state from +5 to -1 so 6e^- must be added. To balance the 3 oxygen atoms on left side, add 3 water molecules on right side. You obtain





Since we know the reaction occurs in acidic media, add 6H^+ to the reactants as the source of hydrogen in the water produced.





In summary, you have



To balance the number of electrons appearing in the equations, multiply the oxidation reaction by six. You obtain






Notice; The electrons dropped out.

• PROBLEM 568

Balance the equation for the following reaction taking place in aqueous acid solution:



Solution: The equation in this problem involves both an oxidation and a reduction reaction. It can be balanced by using the following rules: (1) Separate the net reaction into its two major components, the oxidation process (the loss of electrons) and the reduction process (the gain of electrons). For each of these reactions, balance the charges by adding H^+ , if the reaction is occurring in an acidic medium, or OH^- in a basic medium. (2) Balance the oxygens by addition of H_2O . (3) Balance hydrogen atoms by addition of H . (4) Combine the two half reactions, so that all charges from electron transfer cancel out. These rules are applied in the following example.


The net reaction is



The oxidation reaction is




The I atom went from an oxidation number of 0 in I_2 to + 5 in IO_3^- , because O always has a - 2 charge. You begin with I_2 , therefore, 2 moles of IO_3^- must be produced and 10 electrons are lost, 5 from each I atom. Recall, the next step is to balance the charges. The right side has a total of 12 negative charges. Add 12 H^+ 's to obtain



To balance the oxygen atoms, add $6\text{H}_2\text{O}$ to the left side, since there are 6 O's on the right, thus,



Hydrogens are already balanced. There are 12 on each side. Proceed to the reduction reaction:




Cr began with an oxidation state of + 6 and went to + 3. Since 2Cr^{3+} are produced, and you began with $\text{Cr}_2\text{O}_7^{2-}$, a total of 6 electrons are added to the left. Balancing charges; the left side has 8 negative charges and right side has 6


positive charges. If you add 14 H^+ to left, they balance. Both sides now have a net $+3$ charge. The equation can now be written.




To balance oxygen atoms, add $7\text{H}_2\text{O}$'s to right. You obtain




The hydrogens are also balanced, 14 on each side. The oxidation reaction becomes




The reduction reaction is



Combine these two in such a manner that the number of electrons used in the oxidation reaction is equal to the number used in the reduction. To do this, note that the oxidation reaction has 10e^- and the reduction 6e^- . Both are a multiple of 30. Multiply the oxidation reaction by 3, and the reduction reaction by 5, obtaining



Add these two half-reactions together.



Simplifying, you obtain:



This is the balanced equation.

• PROBLEM 569

Balance the equation for the following reaction taking place in aqueous basic solution:




Solution: The equation in this problem involves both oxidation and reduction. When balancing it, you can use


the following rules. Separate the net reaction into its two major components, the oxidation reaction (the loss of electrons) and the reduction reaction (the gain of electrons). For each half-reaction, balance the charges with H^+ , if the medium is acidic, or OH^- , if the medium is basic. Next, balance the oxygens by the addition of H_2O . Balance total atoms by the addition of H atoms. Finally, combine the two half reactions, so that all charges from electron transfer cancel out.

You employ these rules as follows:

The net reaction is




The oxidation process is



The oxygen atoms in H_2O_2 go from -1 to zero in O_2 . Thus, you have a loss of two electrons. To balance charges, add 2 OH^- to the left side, since there exist 2 negative charges on the right side. You obtain



You now have 4 oxygens on the left, but only 2 on the right. Thus, add 2 water molecules to the right, obtaining




There are the same number of H's on each side. Proceed, now, to reduction half-reaction. Here,




Mn begins with a +7 oxidation number and ends up with +4 in MnO_2 . Therefore, 3 electrons must be added to the left side of the equation. To balance the charges, add 4 OH^-


ions to right, since you have a total of 4 negative charges on left. Rewriting the equation



Add 2 water molecules to the left, so that oxygen atoms can be balanced, obtaining




The hydrogens are balanced. Thus,




To combine these two so that electrons cancel out. Select a multiple of 3 and 2, since these are the number of


electrons involved in the half-reactions. This multiple is six. Multiply the oxidation by 3 and the reduction by 2, obtaining



The net reaction is the total. Thus, adding you obtain:



Cancel the electrons, subtract OH^- ions and H_2O 's to obtain:




which is the balanced equation.

• PROBLEM 570

Determine the volume in milliliters of .20 M KMnO_4 required to oxidize 25.0 ml of .40 M FeSO_4 in acidic solution. Assume the reaction which occurs is the oxidation of Fe^{2+} by MnO_4^- to give Fe^{3+} and Mn^{2+} .

Solution: This problem can be solved by two methods: the mole and the equivalent methods. The mole method requires consideration of the balanced equation that illustrates the reaction. From the data provided, this equation becomes



Now 25.0 ml of .40 M FeSO_4 furnishes
 $(.025 \text{ liters})(.40 \text{ mol/liter}) = .010 \text{ moles of Fe}^{2+}$, since the definition of molarity is

$$M = \frac{\text{number of moles of solute}}{\text{number of liters}}.$$

The balanced equation indicates that the number of moles of MnO_4^- will be 1/5 that of Fe^{2+} . As such, the number of moles of $\text{MnO}_4^- = (.010)(1/5) = .002 \text{ moles}$. Since the KMnO_4 solution has a concentration of .2 M, then the number of liters required is

$$\frac{.002 \text{ mol of MnO}_4^-}{.2 \text{ mol/liter}} = .01 \text{ liters, which equals 10 ml.}$$

The equivalent method functions differently. An equivalent is defined as that mass of oxidizing or reducing agent that picks up or releases the Avogadro number of electrons. Normality is defined as the number of equivalents per liter. Since, in going from Fe^{2+} to Fe^{3+} , you lose 1 electron, .40 M FeSO_4 is equal to .40 N FeSO_4 . Recalling the definition of normality, you have


Solution: Whenever a problem deals with weights and electricity, the solution involves an application of Faraday's Law: The passage of 1 faraday of electricity (96,500 coulombs) causes 1 equivalent weight of matter to be oxidized (the loss of 1 electrons) at one electrode and the reduction (the gain of 1 electrons) of 1 equivalent weight at the other electrode. Equivalent weight may be defined as molecular weight divided by number of moles of hydrogen transferred. To solve this problem, therefore, calculate the number of equivalents present in 454g of water. Water has a molecular weight of 18g/mole, but since 2 H's are transferred, water has an equivalent weight of 9g. Therefore, the number of equivalents is

$$\frac{\text{total weight}}{\text{equivalent weight}} = \frac{454}{9} = 50.4 \text{ equiv.}$$

Recalling that 1 faraday of electricity is used per equivalent, 50.4 equivalents times 1 Faraday/equivalent = 50.4 faradays of electricity required to decompose 454g of water by electrolysis.

• PROBLEM 573

The flashlight battery is a zinc and manganese dioxide (MnO_2) dry cell. The anode consists of a zinc can and the cathode of a carbon rod surrounded by a moist mixture of MnO_2 , carbon, ammonium chloride (NH_4Cl_2), and zinc chloride (ZnCl_2). The cathode reaction is written



If the cathode in a typical zinc and manganese dioxide dry cell contains 4.35g of MnO_2 , how long can it deliver a steady current of 2.0 milliamperes (mA) before all its chemicals are exhausted?

Solution: The problem is solved by calculating the amount of charge required to exhaust the supply of MnO_2 and, from this, determining the lifetime of the battery using the relationship charge (coulombs) = current (A) \times time (sec), or,

$$\text{time (sec)} = \frac{\text{charge (coulombs)}}{\text{current (A)}}.$$

The cathode reaction indicates that two moles of MnO_2 , $2F \text{ MnO}_2$, are consumed for every two moles of electrons present ($2F e^-$). The number of moles of MnO_2 present is

$$\frac{\text{mass, MnO}_2}{\text{molecular weight, MnO}_2} = \frac{4.35\text{g}}{87\text{g/mole}} = 0.05 \text{ mole MnO}_2.$$

Hence, it requires 0.05 mole of electrons (or 0.05F) to consume the 0.05 mole of MnO_2 in the cathode. Converting Faraday's to coulombs (there are 96,500 coulombs in 1F), 0.05 F is equivalent to

$$0.05F \times 96,500 \text{ coulombs/F} = 4.8 \times 10^3 \text{ coulombs.}$$

The battery is supposed to deliver 2.0×10^{-3} amp. Therefore, the lifetime of the battery is

$$\text{time} = \frac{\text{charge}}{\text{current}} = \frac{4.8 \times 10^3 \text{ coulombs}}{2.0 \times 10^{-3} \text{ amp}} = 2.4 \times 10^6 \text{ sec.}$$

Therefore the battery lasts 2.4×10^6 sec (about 30 days).

You pass a 1.0 amp current through an electrolytic cell for 1.0 hr. There are 96,500 coul in a Faraday (F). Calculate the number of grams of each of the following that would be deposited at the cathode:

- (1) Ag from an Ag^+ solution, (2) Cu from a Cu^{+2} solution and (3) Al from an Al^{3+} solution.

Solution: To answer this problem, you can use Faraday's Laws of Electrolysis. Electrolysis is the phenomenon that occurs when electricity is passed through a solution, such that ions are generated and move toward an anode or cathode. The laws are as follows: Masses of substances involved are proportional to the quantity of electricity that flows through the electrolytic cell. Masses of different substances produced during the process are proportional to their equivalent weight. The electrical equivalent is defined as a Faraday (F). It is capable of reducing one equivalent of positive charge, i.e., Avogadro's number of individual unit electric charges. With this information, you can calculate how many Faraday's of electricity were passed through the solution. This tells you the equivalents of the substance that are reduced (recall, one Faraday reduces one equivalent of positive charge). From this number, the weight deposited can be determined.

For all 3 parts, the Faradays generated = $(1.00 \text{ amp}) \left(\frac{1 \text{ coul/sec.}}{\text{amp}} \right) (1 \text{ hr})$
 times $\left(\frac{\text{F}}{96,500 \text{ coul}} \right) = .0373 \text{ F.}$

(1) Ag from an Ag^+ solution. 1 electron is transferred in $\text{Ag}^+ + e^- \rightarrow \text{Ag(s)}$. Since one electron is transferred per Ag atom, 1 mole of Ag atoms requires one mole of electrons or 1F. 1 mole of Ag atoms weighs 107.87g (atomic weight - see Periodic Table). You have, though, only .0373 F. Thus, $(.0373 \text{ mole})(107.87\text{g/mole}) = 4.02\text{g}$ of Ag is deposited.

(2) Here, 2 electrons are transferred in $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu(s)}$. Thus, 1F is required for $\frac{1}{2}$ of a mole of Cu(s) to be deposited. 1 mole weighs 63.55, $\frac{1}{2}$ of a mole = $63.55/2 = 31.775\text{g}$. Thus, with .0373 F, you can deposit $.0373(31.775) = 1.19\text{g}$.

(3) Here, 3 electrons are transferred. Thus, 1F can only deposit $1/3$ of a mole.

$$\text{weight deposited} = \left(\frac{\text{mw}}{3} \right) \times (.0373) = \left(\frac{26.98}{3} \right) (.0373) = .335\text{g.}$$

When a current of 5.36 A is passed for 30 min. through an electrolytic cell containing a solution of a zinc salt, it is found that 3.27g of zinc metal is deposited at the cathode. What is the equivalent weight of zinc in the salt?

Solution: To solve this problem one must recall that 1 coulomb = 1A-sec. This means that the number of coulombs passed through the cell is equal to the product of the current (in amperes) and the time (in seconds). In our case, since 5.36 A was passed for 30 min. $(30 \text{ min.} \times 60 \text{ sec/min} = 1800 \text{ sec})$, the number of coulombs is $5.36\text{A} \times 1800 \text{ sec} = 9650 \text{ coulombs}$. One Faraday (19) is equivalent to 96,500

a current of 4.02 amperes flows for 1000 minutes. One ampere is defined as one coulomb per second

ampere = coulomb/second.

To find the total number of coulombs that were transmitted to the Ni^{2+} ions, the amperage of the current must be multiplied by the time the current was flowing. The time must be converted from minutes to seconds by use of the conversion factor, 60 secs/1 minute.

$$\begin{aligned}\text{no. of coulombs} &= 4.02 \text{ coulombs/sec.} \times 1000 \text{ min.} \\ &\times 60 \text{ sec/minute} = 24,120 \text{ coulombs.}\end{aligned}$$

Since one knows that 2 Faradays or 193,000 coulombs ($2 \times 96,500$ coul) plate out 1 mole, or 58.7g, of Ni, one can set up the following ratio to determine the number of grams that 24,120 coulombs plate out. Let x = the number of grams of Ni^{2+} that 24,120 coulombs will convert to Ni.

$$\begin{aligned}\frac{58.7 \text{ grams}}{193,000 \text{ coulombs}} &= \frac{x}{24,120 \text{ coulombs}} \\ x &= \frac{24,120 \times 58.7 \text{ g.}}{193,000} = 7.34 \text{ g.}\end{aligned}$$


7.34g of Ni will be deposited.

• PROBLEM 579

A meter reads that a battery is putting out .450 amp in the external circuit of a cell. The cell is involved in the electrolysis of a copper sulfate solution. During the 30.0 min that the current was allowed to flow, a total of .30g of copper metal was deposited at the cell's cathode. Was the meter an accurate measurement of current? ($F = 96,500$ coul.)

Solution: The key to solving this problem is the determination of the actual current given the amount of copper deposited in 30.0 minutes. This can be done by employing Faraday's Laws of Electrolysis. Electrolysis is the phenomenon that occurs when electricity is passed through a solution, such that ions are generated and moved toward an anode or cathode. The laws that govern this are as follows: Masses of substances involved are proportional to the quantity of electricity that flows through the cell. Masses of different substances produced during the process are proportional to their equivalent weight. The electrical equivalent is defined as a Faraday (F). It is capable of reducing one equivalent of positive charge (Avogadro's number of individual unit electric charges). The problem can now be solved.

You know how much material is deposited in a given time. Thus, a certain amount of electricity based on current had to be used. The reaction that occurred at the cathode was



Thus, each mole of copper ion requires 2 moles of electrons or, in other words, 2 faradays (1 Faraday = 96,500 coul). Thus, a total of 193,000 coul. is required per mole of copper. Current = charge/time. If you had one mole of copper, the charge would be 193,000 coul. You have only

$$\frac{0.30 \text{g(wt. of Cu)}}{63.55 \text{(m.w. of Cu)}} = 4.72 \times 10^{-3} \text{ moles.}$$

Thus, charge = 4.72×10^{-3} (193,000) = 911.075 coul. To determine the current, time must be considered. The time must be in seconds, since

$$\text{current (amps)} = \frac{\text{charge (coults)}}{\text{time (secs)}}$$

You are given the time as 30.0 min, which equals 1800 secs. Thus,

$$\text{current} = \frac{911.075}{1800} = .506 \text{ amp.}$$


The meter said .450, thus, it was inaccurate by

$$\left(\frac{.506 - .450}{.506} \right) \times 100 = 11.1\%$$


• PROBLEM 58C

Discuss the separate half-reactions and net cell reaction that occur in a flashlight battery. If you have a current of .01 amp, how long can the current flow if the battery contains 25.0g of a paste of MnO_2 , NH_4Cl , and ZnCl_2 of which 20% is MnO_2 ?

Solution: A standard flashlight dry cell, or Leclanché cell, is composed of a graphite rod in a moist MnO_2 - ZnCl_2 - NH_4Cl paste, all in a zinc wrapper. The zinc wrapper serves as the electrode for the oxidation half-cell (the anode), while the graphite rod is the electrode for the reduction half-cell (the cathode). Thus, the probable half-reactions are: at the anode:



and at the cathode:



The net reaction for the overall process is $\text{Zn(s)} + 2\text{MnO}_2 \rightarrow \text{ZnO} \cdot \text{Mn}_2\text{O}_3$.

To determine how long the current can flow, calculate the number of moles of MnO_2 present and the number of Faraday's required to reduce it per mole. You are told that 20% of 25 grams of paste is MnO_2 . Thus, you have $(.20)(25\text{g}) = 5.0\text{g}$ of MnO_2 . The molecular weight of $\text{MnO}_2 = 86.94\text{g}$. Thus, you have

$$5\text{g}/86.94\text{g/mole} = .0575 \text{ moles of } \text{MnO}_2$$



From the net equation, you see that 2 electrons are transferred. A Faraday = 96,500 coul and is capable of reducing one equivalent positive charge, which means Avogadro's number of individual unit charges (one mole of electrons). Since 2 electrons were transferred per 2 moles, it takes 2 Faraday's to reduce 2 moles of MnO_2 . According to the net equation 2 moles of MnO_2 are reduced for every mole of $\text{ZnO} \cdot \text{Mn}_2\text{O}_3$ obtained. Thus, you can write that the current will last for

$$(.0575 \text{ moles } \text{MnO}_2) \left(\frac{2\text{F}}{2 \text{ moles } \text{MnO}_2} \right) \left(\frac{96,500 \text{ coul}}{\text{F}} \right) \left(\frac{\text{amp}}{\text{coul/sec}} \right) \left(\frac{1}{.01 \text{ amp}} \right)$$


$$= 154 \text{ hours.}$$

$\text{Cr}_2\text{O}_7^{2-} + 14 \text{H}^+ + 6\text{e}^- \rightleftharpoons 2 \text{Cr}^{3+} + 7 \text{H}_2\text{O}$	1.33
$\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2 \text{Cl}^-$	1.36
$\text{MnO}_4^- + 8 \text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4 \text{H}_2\text{O}$	1.51
$\text{Ce}^{4+} + \text{e}^- \rightleftharpoons \text{Ce}^{3+}$	1.61
$\text{MnO}_4^- + 4 \text{H}^+ + 3\text{e}^- \rightleftharpoons \text{MnO}_2(\text{s}) + 2 \text{H}_2\text{O}$	1.68
$\text{H}_2\text{O}_2 + 2 \text{H}^+ + 2\text{e}^- \rightleftharpoons 2 \text{H}_2\text{O}$	1.77
$\text{O}_3 + 2 \text{H}^+ + 2\text{e}^- \rightleftharpoons \text{O}_2 + \text{H}_2\text{O}$	2.07
$\text{F}_2 + 2\text{e}^- \rightleftharpoons 2 \text{F}^-$	2.87

Solution: The best oxidizing agent will be the one with the greatest ability to gain electrons (be reduced) and will therefore have the most positive standard electrode potential, E° . From the tables,



Thus the substances, in order of decreasing ability as oxidizing agents, are



• PROBLEM 584

Calculate ΔE° for the following cells: (1) Cadmium and Hydrogen, (2) Silver and Hydrogen and (3) Cadmium and silver, using the following data:

Reaction	E° volts
$\text{Cd} - \text{Cd}^{2+} + 2\text{e}^-$	+ .403
$\text{H}_2 - 2\text{H}^+ + 2\text{e}^-$	0.00
$\text{Ag} - \text{Ag}^+ + \text{e}^-$	- .799

Solution: You are asked to calculate the standard cell potential (ΔE°) for each of the given pairs. To do this, you must realize that in such cells, you have 2 half-reactions. Namely, an oxidation reaction (loss of electrons) and a reduction reaction (gain of electrons). The sum of these half-reactions yields the overall reaction and the ΔE° of the whole cell. Thus, to find the ΔE° for each of these pairs, you need to know the E° of the half-reactions. There is one other important fact to be kept in mind.

In cells, if the reaction is to proceed spontaneously, ΔE° must have a positive value. This means, therefore, that you must choose the half-reactions such that their sum always gives a positive ΔE° . You are told the E° of the oxidation half-reaction for each element in each pair. The reduction half-reaction is the reverse of the oxidation reaction for each element, with a change in sign of E° . For example, if you have $\text{A} - \text{A}^{+1} + \text{e}^-$ with an $E^\circ = B$ for oxida-

tion, the reduction is $A^{+1} + e^{-} \rightarrow A$ with a $E^{\circ} = -B$.

With this in mind, the procedure is as follows:

(1) Cadmium (Cd) and hydrogen (H_2).

The reaction for this cell must be the sum of the oxidation and reduction such that the ΔE° is positive. This can only occur if the anode (oxidation) has the higher oxidation potential. Thus, you calculate ΔE° as $+ .403 - (0.000) = +.403v$. Similarly, for (2) and (3), a positive ΔE° can only be obtained with the anode having the higher oxidation potential. Thus, for (2), $E^{\circ} = (0.000) - (-.799) = +.799v$. For (3), $E^{\circ} = (.403) - (-.799) = +1.202v$.

● PROBLEM 585

If I_2 and Br_2 are added to a solution containing I^{-} and Br^{-} , what reaction will occur if the concentration of each species is 1 m?

Some half-reactions and their standard reduction potentials.


	Half-reaction	Standard reduction potential, V
	$2e^{-} + F_2(g) \rightarrow 2F^{-}$	+2.87
	$2e^{-} + Cl_2(g) \rightarrow 2Cl^{-}$	+1.36
	$4e^{-} + 4H_3O^{+} + O_2(g) \rightarrow 6H_2O$	+1.23
	$2e^{-} + Br_2 \rightarrow 2Br^{-}$	+1.09
	$e^{-} + Ag^{+} \rightarrow Ag(s)$	+0.80
	$2e^{-} + I_2 \rightarrow 2I^{-}$	+0.54
	$2e^{-} + Cu^{2+} \rightarrow Cu(s)$	+0.34
	$2e^{-} + 2H_3O^{+} \rightarrow H_2(g) + 2H_2O$	zero
	$2e^{-} + Fe^{2+} \rightarrow Fe(s)$	-0.44
	$2e^{-} + Zn^{2+} \rightarrow Zn(s)$	-0.76
	$3e^{-} + Al^{3+} \rightarrow Al(s)$	-1.66
	$2e^{-} + Mg^{2+} \rightarrow Mg(s)$	-2.37
	$e^{-} + Na^{+} \rightarrow Na(s)$	-2.71
	$e^{-} + Li^{+} \rightarrow Li(s)$	-3.05

Solution: When I_2 and Br_2 are added to the solution, a

dynamic equilibrium is attained between the non-charged element and its ion form. These reactions, redox reactions, are written as:



The voltages listed indicate the electric potential between two electrodes. In other words, it is a measurement of the work done in moving a unit charge (an electron) from one electrode to the other. For any reaction to occur, the



Notice: Since both equations contained 5e^{-} on different sides, they cancelled out. This explains why the oxidation reaction is multiplied by five. Thus you have written the balanced equation.

(b) The ΔE° for the overall reaction is the sum of the E° for the half-reactions, i.e., $\Delta E^{\circ} = E_{\text{red}}^{\circ} + E_{\text{oxid}}^{\circ}$. You know that E_{red}° and E_{oxid}° ; $\Delta E^{\circ} = 1.51 - .77 = 0.74 \text{ eV}$.

(c) A reaction will only proceed spontaneously when $\Delta E^{\circ} =$ a positive value. You calculated a positive ΔE° , which means the reaction proceeds spontaneously.

ELECTROCHEMICAL CELL REACTIONS


• PROBLEM 587

For the following voltaic cell, write the half-reactions, designating which is oxidation and which is reduction. Write the cell reaction and calculate the voltage of the cell made from standard electrodes. The cell is $\text{Co}; \text{Co}^{+2} \parallel \text{Ni}^{+2}; \text{Ni}$.

Solution: The cell reaction is the algebraic sum of the reactions that take place at the electrodes. Every cell has 2 electrodes an anode and a cathode. Oxidation, which is the loss of electrons, occurs at the anode. Reduction, which is the gain of electrons, takes place at the cathode.

The cell is always written as solid; ion in solution \parallel ion in solution; solid (anode) (cathode)

Oxidation and reduction are the half reactions that take place in the cell. For this cell, they are



Sum: $\text{Ni}^{+2} + \text{Co} \rightarrow \text{Ni} + \text{Co}^{+2}$ (Cell reaction).


Since Co is losing electrons, it provides the oxidation reaction and Ni^{+2} , gaining these electrons, takes part in the reduction reaction.

The voltage of a cell is the sum of the oxidation and reduction potentials in units of volts and is designated by E° (under standard conditions).

$$E_{\text{cell}}^{\circ} = E_{\text{oxidation}}^{\circ} + E_{\text{reduction}}^{\circ}$$

The voltages of half cell reactions are usually given as the reduction

Ni, for which the half-reactions are



Since Al is losing electrons (oxidation), and Ni is gaining electrons (reduction), the electron flow is from Al to Ni

E° , the standard electrode potential is the sum of the E° 's for the half-reactions listed above


$$\begin{aligned} E^\circ &= E^\circ_{\text{ox}} + E^\circ_{\text{red}} \\ &= 1.66 + (-.25) \\ &= 1.41 \text{ volts} \end{aligned}$$

• PROBLEM 590


Given the following standard electrode potentials at 25°C: $\text{Sn}^{4+} + 2\text{e}^- \rightarrow \text{Sn}^{2+}$, $E^\circ = 0.15 \text{ eV}$ and $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$, $E^\circ = +0.77 \text{ eV}$ will the reaction $\text{Sn}^{2+} + 2\text{Fe}^{3+} \rightleftharpoons \text{Sn}^{4+} + 2\text{Fe}^{2+}$ proceed spontaneously?

Solution: A reaction will proceed spontaneously only if it has a positive ΔE° as written. Thus, to answer this question, you want to calculate the ΔE° and see whether it is positive. This can be done by considering its half-reactions. The sum of the electrode potentials for these reactions will be the ΔE° .


The overall reaction is




The half-reactions are, oxidation (loss of electrons) and reduction (gain of electrons) reactions. You have then,



This, has the reverse of the standard potential of the reaction given. Thus, its $E^\circ = -(0.15) = -.15 \text{ eV}$.




This has the same standard potential as in the given reaction. Thus, its $E^\circ = 0.77 \text{ eV}$. Adding these two equations together, one obtains the desired net reaction.



The $\Delta E^\circ = E^\circ_{\text{oxid}} + E^\circ_{\text{red}} = -.15 + .77 = +.62 \text{ volt}$. The fact that it is positive means the reaction proceeds spontaneously.

• PROBLEM 591

Construct a galvanic cell based on the reaction




with ΔH negative.

Notice, that both Le Chatelier's principle and the Nernst equation relate what happens to reaction tendencies when concentrations are changed. For example, in the Nernst equation, if the concentrations are such that yield a positive potential, E , then the reaction is favorable in the direction written. The difference between the Nernst equation and Le Chatelier's principle is that the latter relates reaction tendencies with a change of concentration qualitatively, while the former discusses it quantitatively.




• PROBLEM 593

A cell possesses two electrodes. Both half-cells are .01M MnO_4^- ion. One cell is .01M H_3O^+ ion, while the other has a H_3O^+ concentration of .10M. The electrode reaction for the reduction half-cell may be written:



The oxidation half-cell is the reverse of this reaction. 1) Write the net equation for the spontaneous cell process taking place; 2) Find ΔE for the reaction; 3) Find the value of the equilibrium constant.

Solution: 1) The net reaction, in such a situation, is the sum of the balanced half-reactions, i.e., oxidation reaction plus reduction reaction. Since, you are given both reactions, add the equations together to find the overall reaction. You have, with concentrations included,



Notice: all species cancelled out, except H^+ (actually H_3O^+). This is the net equation for the spontaneous reaction taking place.

2) To find ΔE for the reaction, use the Nernst equation, which states

$$\Delta E = \Delta E^\circ - \frac{.059}{n} \log K$$


for a temperature at 25°C , where ΔE = potential for cells under other than standard conditions, ΔE° = standard cell potential, n = number of electrons transferred and K = equilibrium concentration expression. $n = 3$, since from either the oxidation or reduction reaction, 3 electrons are being transferred; $\Delta E^\circ = 0$, since $\Delta E^\circ = E^\circ_{\text{prod}} - E^\circ_{\text{reactants}}$ and both the product and reactant are the same species. K is defined as the ratio of products to reactants, each raised to the power of their coefficients in the net equation. Substituting these values into the Nernst equation, you have

$$\begin{aligned} \Delta E &= 0 - \frac{.059}{3} \log \frac{(.01)^4}{(.10)^4} = 0 - .0197(\log 10^{-4}) \\ &= (-.0197)(-4) = 0.079 \text{ volt.} \end{aligned}$$

3) To find the value of the equilibrium constant, note that there exists a relationship between ΔE° and the constant at 25°C . Namely,



$\Delta E^\circ = \frac{.059}{n} \log K$. From part 2, you found $\Delta E^\circ = 0$. Thus, $K = \text{unity}$ (one), since this is the only value that permits $\log K = 0$, which, then, allows $\Delta E^\circ = 0$, as it must.

You have the following cell process:





$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe(s)}$ with $E^\circ = -.44\text{e}$ and $\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co(s)}$ with $E^\circ = -.28$, find the standard cell potential ΔE° , the cell potential ΔE and the concentration ratio at which the potential generated by the cell is exactly zero.

Solution: Assume that the reaction proceeds spontaneously. This means, therefore, that the reaction must have a positive value for ΔE° . With this in mind, you proceed as follows: You are given two half-reactions:



Both reactions represent reduction (gain of electrons). But, the overall reaction in a cell is a combination of both a reduction and an oxidation reaction. Thus, you must reverse one, keeping in mind that the ΔE° must be a positive value. Recall, also, that $\Delta E^\circ = E^\circ_{\text{oxid}} + E^\circ_{\text{red}}$. You can write



with $\Delta E^\circ = .44 - .28 = .16 \text{ ev}$. Notice: By reversing the Fe reaction and combining it with the other, you obtained the overall reaction with a $\Delta E^\circ = .16 \text{ ev}$, a positive value, which indicates that the reaction proceeds spontaneously. To find ΔE , use the Nernst equation, which states

$$\Delta E = \Delta E^\circ - \frac{.059}{n} \log K,$$

where n = number of electrons transferred and K = equilibrium constant of reaction. In this problem, the number of electrons transferred is 2, so that $n = 2$.

$$K = \frac{[\text{Fe}^{2+}]}{[\text{Co}^{2+}]}, \text{ i.e., the ratio of the concentrations}$$

of products to reactants, each raised to the power of its respective coefficient in the chemical equation. Note, Co(s) and Fe(s) are omitted, because they are solids and, thus, considered constants themselves. You are given $[\text{Fe}^{2+}]$ and $[\text{Co}^{2+}]$ and you have calculated ΔE° . Therefore,


$$\begin{aligned} \Delta E &= \Delta E^\circ - \frac{.059}{n} \log \frac{[\text{Fe}^{2+}]}{[\text{Co}^{2+}]} = .16 - \frac{.059}{2} \log \frac{(1.0)}{(.5)} = .16 - .0295 \log 2 \\ &= .16 - .01 = .15 \text{ ev.} \end{aligned}$$

To find the concentration ratio, K , when $\Delta E = 0$, use the Nernst equation;


$$\Delta E = \Delta E^\circ - \frac{.059}{n} \log K.$$

$$\Delta E = 0, \Delta E^\circ = .16, n = 2 \text{ and } K = \frac{[\text{Fe}^{2+}]}{[\text{Co}^{2+}]}, \text{ so that}$$

and the reduction is



overall reaction



From the overall reaction, recall

$$K = \frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}$$

(The concentrations of solids are not included in a dissociation constant). You are told $\text{Zn}^{2+} = .5\text{m}$ and $\text{Cu}^{2+} = .2\text{m}$, so that

$K = \frac{.50}{.20}$. In the overall reaction, 2 electrons are lost and gained, which means $n = 2$. If you substitute these values in the Nernst equation, you obtain

$$E = 1.1 - \frac{-.592}{2} \log \frac{.50}{.20}.$$

Solving, $E = 1.088\text{v}$ for this Daniell cell.

• PROBLEM 596

Calculate the voltage of the cell $\text{Fe}; \text{Fe}^{+2} \parallel \text{H}^+; \text{H}_2$ if the iron half cell is at standard conditions but the H^+ ion concentrations is .001M.

Solution: The voltage (E) of a cell is found using the Nernst Equation because it involves the use of concentration factors. It is stated

$$E = E^\circ - \frac{RT}{nF} \ln Q,$$

where R is 8.314 joules per degree, F is 96,500 coulombs, n is the number of moles of electrons transferred and Q is the concentration term. $T = 25^\circ \text{C}$, by definition of standard conditions, in this equation (or 298°K).

But to solve the problem we must first obtain E° . This is done by writing down the appropriate half-reactions. Oxidation is the loss of electrons and reduction is the gain of electrons.

Reaction	Type	E°_{red}
$\text{Fe} \rightarrow \text{Fe}^{+2} + 2e^-$	Oxidation	+ .44
$2\text{H}^+ + 2e^- \rightarrow \text{H}_2$	Reduction	0

Next, take the algebraic sum of the E°_{red} and E°_{oxid} , which gives E° .

$$E^\circ = +.44 + 0 = +.44.$$

Now set up the concentration term.

$$\ln Q = \ln \frac{[\text{H}^+]^2}{[\text{Fe}^{+2}]} = \ln \frac{[.001]^2}{[1]}$$

Standard conditions always means a concentration of 1 M. Substituting all these terms into the Nernst equation, one calculates E to be


$$E = +.44 - \frac{.059}{2} \log \frac{10^{-6}}{1} = 0.617\text{v} \quad \text{or} \quad E = +.44 - \frac{0.0257}{2} \ln \frac{10^{-6}}{1} = 0.617\text{v}$$

half reaction is -0.799 volts.

Solution: The silver-silver chloride electrode process is a special case of the (Ag, Ag^+) couple, except that silver ions collect as solid AgCl on the electrode itself. However, solid AgCl has some Ag^+ in equilibrium with it in solution. This $[\text{Ag}^+]$ can be calculated from the K_{sp} equation:

$$[\text{Ag}^+][\text{Cl}^-] = K_{sp} \quad \text{or} \quad [\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]}$$
$$[\text{Ag}^+] = \frac{1.8 \times 10^{-10}}{1} = 1.8 \times 10^{-10}$$

This value for $[\text{Ag}^+]$ can be inserted into the Nernst equation for the (Ag, Ag^+) half-reaction



The Nernst equation is

$$E = E^\circ - \frac{0.05916 \log Q}{n}, \quad \text{where } E^\circ = \text{standard}$$

electrode potential (25°C and 1M), n = number of electrons transferred, and Q = the mass-action expression or equilibrium constant. For this reaction, one sees that one electron is transferred. Hence, $n = 1$.

Also, $Q = \frac{[\text{Ag}^+]}{[\text{Ag}]}$, but, since Ag is solid, its concentration is assumed to be constant; it can be removed from the expression, so that $Q = [\text{Ag}^+]$. Thus,

$$E = E^\circ - \frac{0.05916 \log [\text{Ag}^+]}{1} = -0.799 - 0.05916 \log(1.8 \times 10^{-10})$$
$$= -0.799 + 0.576 = -0.223 \text{ volts}$$

• PROBLEM 601

What will be the relative concentrations of Zn^{+2} and Cu^{+2} when the cell $\text{Zn}; \text{Zn}^{+2} || \text{Cu}^{+2}; \text{Cu}$ has completely run down? ($E^\circ = 1.10$ volts).


Solution: The key phrase to understand before solving this problem is "has completely run down". That means there is no apparent current, leading us to assume a value of zero for the electrode potential, E .

Next, one should realize that whenever concentrations are involved one should apply the Nernst Equation, which states

$$E = E^\circ - \frac{RT}{nF} \ln Q,$$

where Q is the equilibrium constant expression, R is the gas constant (8.314 joules per degree), F is $96,500$ coulombs, n is the number of moles of electrons transferred and T is 25°C (or 298°K).

Since the overall cell reaction is



2 moles of electrons are transferred and $\log Q$ equals $\log \frac{[\text{Zn}^{+2}]}{[\text{Cu}^{+2}]}$.

Remember that solids are omitted from all concentration equations.

This yields (via substitution):

$$E = E^\circ - \frac{RT}{nF} \ln Q$$

and

$$0 = 1.1 \text{ volts} - \frac{.059}{2} \log \frac{[\text{Zn}^{+2}]}{[\text{Cu}^{+2}]}$$

which means $\log \frac{[\text{Zn}^{+2}]}{[\text{Cu}^{+2}]} = 37.3$. Solving,

$$\frac{[\text{Zn}^{+2}]}{[\text{Cu}^{+2}]} = 2 \times 10^{37}$$

This means that the concentration of Zn^{+2} is far greater than Cu^{+2} .

• PROBLEM 602

For a cell reaction of the type $\text{T} + \text{R}^{+2} \rightleftharpoons \text{R} + \text{T}^{+2}$, calculate the value that E° must have so that the equilibrium concentration of T^{+2} is a thousand times the R^{+2} concentration.

Solution: In this question, we are asked to obtain the standard cell potential (E°), not the actual cell potential (E). Since there is no net electron flow at equilibrium, $E = 0$.

There exists an expression which relates E° , E and concentrations. This expression, called the Nernst equation, states $E = E^\circ - .059/n \log Q$, at a temperature of 25°C where n = number of electrons transferred and Q = mass action expression.

From the chemical reaction, you see T goes from an oxidation state of 0 to +2. It lost 2 electrons. Thus, R^{+2} gained 2 electrons. Therefore, the number of electrons transferred, n , equals 2. From the same chemical reaction, Q becomes

$$\frac{[\text{T}^{+2}]}{[\text{R}^{+2}]}$$


You are asked to determine $[\text{T}^{+2}]$ to be 1000 times $[\text{R}^{+2}]$, which means $Q = 1000$. You already determined that $E = 0$. To find E° , substitute these values in the Nernst equation. You obtain

$$0 = E^\circ - \frac{.059}{2} \ln 1000$$

Solving, $E^\circ = .088 \text{ volt}$.

• PROBLEM 603

You are given the reaction:



It has been determined that at 25°C the standard free energy of formation of $\text{AgCl}(\text{s})$ is -109.7 kJ/mole , that of $\text{H}_2\text{O}(\text{l})$ is -237.2 kJ/mole , and that of $(\text{H}_3\text{O}^+ + \text{Cl}^-)(\text{aq})$ is -368.4 kJ/mole . Now assuming the reaction is run at 25°C and 1 atm of pressure in a cell in which the $\text{H}_2(\text{g})$ activity is unity and $\text{H}_3\text{O}^+(\text{aq})$ and $\text{Cl}^-(\text{aq})$ activities are .01, what will be the cell voltage?

Solution: The answer to this question requires the use of the Nernst equation and the relationship between overall free energy change (ΔG) and the cell voltage E .

The Nernst equation implies that the voltage of a cell depends on the concentrations or activity of the species involved. It states

$$E = E^\circ - \frac{RT}{nF} \ln Q,$$

where E is the potential for a reaction at nonstandard conditions, E° = the standard potential, n = number of faradays of electricity, F = the value of a faraday, (for conversion to kilojoules per mole is $96.49 \text{ kJ V}^{-1} \text{ equiv}^{-1}$), T = temperature in kelvin (celsius plus 273°), R = universal gas constant, ($8.31 \text{ J mol}^{-1} \text{ deg}^{-1}$), and Q is the mass-action expression. The relationship between the overall free energy and cell voltage is $\Delta G = -nFE$, where ΔG is the free energy. If all the species are in their standard states, then you can say $\Delta G^\circ = -nFE^\circ$. The sum of ΔG° must be the free energy contribution from each component. Therefore,

$$\begin{aligned}\Delta G^\circ &= 2\Delta G^\circ_{\text{H}_3\text{O}^+\text{Cl}^-} - 2\Delta G^\circ_{\text{AgCl}} - 2\Delta G^\circ_{\text{H}_2\text{O}} \\ &= 2(-368.4) - 2(109.7) - 2(-237.2) \text{ kJ} \\ &= -43.0 \text{ kJ per mole of reaction.}\end{aligned}$$

$$E^\circ = \frac{-\Delta G^\circ}{nF} = - \left[\frac{-43.0 \text{ kJ}}{2(96.49 \text{ kJ/v})} \right] = .223 \text{ v}.$$

Since the activities of solid Ag , of solid AgCl and of H_2O are unity, they can be eliminated from the mass-action expression:

$$Q = \frac{[\text{H}_3\text{O}^+]^2 [\text{Cl}^-]^2}{[\text{H}_2]}.$$

Thus,

$$E = E^\circ - \frac{RT}{nF} \ln \frac{[\text{H}_3\text{O}^+]^2 [\text{Cl}^-]^2}{[\text{H}_2]}.$$

Therefore,

$$\begin{aligned}E &= .223 - \frac{(8.31)(298)(2.303)}{2(96490)} \log \frac{(.0100)^2 (.0100)^2}{1.00} \\ E &= .223 + .236 = .459 \text{ v}.\end{aligned}$$

• PROBLEM 604

You have two half-cells coupled together to form a concentration cell: $\text{H}_2(1 \text{ atm}) + 2\text{H}^+(1\text{M}) + 2\text{e}^-$ and $\text{H}_2(1 \text{ atm}) + 2\text{H}^+(1.0\text{M}) + 2\text{e}^-$. Calculate the free energy and electrical work obtainable. $F = 96,500$ coulombs.


Solution: To find the free energy and electrical work obtainable, calculate the maximum cell voltage in this cell, since $\Delta G = \text{free energy} = -nF\Delta E$, where n = number of electrons transferred, F = Faraday constant, and ΔE = cell voltage.

To find ΔE , use the Nernst equation, which states $\Delta E = \Delta E^\circ - .059/n \log K$, where ΔE° = voltage at standard conditions, and K = equilibrium constant of reaction.

(1) $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag(s)}$ $E^\circ = .80$ volt, and (2) $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al(s)}$ $E^\circ = -1.66$ volt, calculate the standard free energy change, ΔG° , in Kcal/mole. 1 cal = 4.184 joule
1 F = 96,500 coul, 1 Kcal = 1000 cal.

Solution: The standard free energy change, ΔG° , can be related to standard cell potential change, ΔE° , via $\Delta G^\circ = n F \Delta E^\circ$, where n = number of electrons transferred per mole. To find ΔG° , calculate (1) ΔE° , (2) n .

(1) You are told a voltaic cell exists. This means that electricity is generated. You are given the E° values for the half-reactions. For electricity to be generated, however, the ΔE° for the overall reaction must be positive. Both half-reactions represent reduction processes (gain of electrons). But $\Delta E^\circ = E^\circ_{\text{oxid}} + E^\circ_{\text{red}}$. Thus, to find ΔE° , you must reverse one of these reactions to oxidation (loss of electrons) so that ΔE° for the overall reaction, is positive. This condition can only be met if (2) is reversed to $\text{Al(s)} \rightarrow \text{Al}^{3+} + 3\text{e}^-$, $E^\circ = -(-1.66) = 1.66$ and (1) remains as $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag(s)}$. Before adding these two to obtain the overall reaction, multiply (1) by 3, so that all electrons in the overall equation can be cancelled. Thus



(2) The number of electrons being transferred = 3, so that $n = 3$.

$$\Delta G^\circ = n F \Delta E^\circ.$$

Substitute in values to obtain

$$\begin{aligned} \Delta G^\circ &= (3)(96500)(2.46)(\text{cal}/4.184 \text{ joules}) \times (\text{Kcal}/1000 \text{ cal}) \\ &= +170 \text{ Kcal/mole.} \end{aligned}$$

• PROBLEM 606

Calculate the standard E° , electrode potential, of the hydrogen-oxygen fuel cell. The standard Gibbs free energy of formation (ΔG_f°) of $\text{H}_2\text{O(l)}$ = -56,690 cal/mole.

Solution: The standard Gibbs free energy of formation (ΔG_f°) is a measure of the energy available to do work when compounds are formed from their elements in the standard state. The electrode potential,

i.e., the voltage across the two electrodes in a galvanic cell, such as the hydrogen-oxygen cell, can be computed from ΔG_f^0 , by using the equation $\Delta G^0 = -nF E^0$, where n = number of electrons transferred and F = a faraday of electricity, which is $23,060 \text{ cal v}^{-1} \text{ mole}^{-1}$. In a hydrogen-oxygen fuel cell, water is produced from the transfer of 2 electrons. Thus, $n = 2$. Solving for E^0 :

$$-56,690 \frac{\text{cal}}{\text{mole}} = -2(23,060 \text{ cal v}^{-1} \text{ mole}^{-1}) E^0 .$$

$$E^0 = 1.229 \text{ v for this fuel cell.}$$

CHAPTER 17



ATOMIC THEORY

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 581 to 611 for step-by-step solutions to problems.

The atomic weight is defined as the weight of an atom compared to ^{12}C being exactly 12 atomic mass units (amu). This translates into a mole of ^{12}C having a mass of exactly 12 grams. Tabulated values of atomic weights are the weighted averages of the naturally occurring isotopes. (Isotopes have the same number of protons but different numbers of neutrons.) For example, the atomic weight of chlorine is tabulated as 35.45, natural chlorine is 75.5% ^{35}Cl (atomic weight 34.97), and 24.5% ^{37}Cl (atomic weight 36.97).

$$\text{Avg. At. Wt.} = .755(34.97) + .245(36.97) = 35.45 \qquad 17-1$$

Many periodic tables show the electronic configuration for each atomic species. This will be discussed in more detail in the following paragraphs and in Chapter 18, but since chemistry is the study, principally, of the interaction of electrons in the outer or valence orbitals, it is often simplified by considering only the numbers of valence or reactive electrons. The key is to note that in the first orbital there are two valence electrons. The neutral atom with one electron is hydrogen; the one with two is helium. No more electrons can enter the first orbital for reasons to be outlined in Chapter 18. For all other orbitals, up to eight electrons can exist. The tendency or driving force for reaction is to fill the outermost orbital by sharing or acquiring electrons from other atomic species. Therefore, atoms react in a way to get eight electrons in the outermost or valence orbital (except for hydrogen which reacts to get two).

If the valence orbital is filled by sharing electrons the bond is called covalent; if it is filled by acquiring an electron from another atom, the bond is called ionic. For large differences in electronegativity, ionic bonds are

formed; for moderate differences, covalent bonds are formed. The relative electronegativity of the elements is tabulated. An arbitrary scale, called the Pauling scale, is used, and the value of electronegativity for each of the elements is shown in a table in Problem 642. If the difference in electronegativity between two elements is greater than 1.7, the bond between them will be ionic; if less, the bond will be covalent. However, if the difference in electronegativity is near 1.7, bonds may have both covalent and ionic character. These bonds are polar in that the electrons are preferentially attracted to the more electronegative atom. The greater the difference in electronegativity between bonded atoms, the more polar the bond will be until eventually, when the electron is completely attracted to one atom, it becomes ionic.

There are several examples in the problem list showing electron dot diagrams and putting atoms together to obtain the desired number of electrons in the valence orbit. When atoms in Groups I and II of the Periodic Table donate their one or two valence electrons, the electronic configuration that results is that of the filled stable orbit under the valence electrons.

Electrons are numbered for each element by the following convention which is explained by quantum theory and discussed in Chapter 18.

- a) the first number is the principal orbit (e.g., 1, 2, 3, etc.).
- b) the second digit is designated by a letter (e.g., *s*, *p*, *d*, *f*).
- c) the superscript following the letter designates the number of electrons of that type.

The maximum number of each type of electrons in any orbital is 2 for *s* electrons, 6 for *p* electrons, 10 for *d* electrons, and 14 for *f* electrons. The *s* and *p* electrons are the principal reactive electrons for small atoms. The *d* electrons are somewhat less reactive and account for the multiple valence states in many transition metals.


In the first row of the Periodic Table, only *s* electrons exist. In the second and third rows, *s* plus *p* electrons exist, and in subsequent rows the *d* and *f* electrons may also exist. Table 1 shows the electron configuration of a sample of elements from the first three rows of the Periodic Table.

Table 1: Sample of the Electronic Configurations of Elements

Element	At. No.	1s	2s	2p	3s	3p	Electronic Configuration*
H	1	1					1s ¹
He	2	2					1s ²
C	6	2	2	2			1s ² , 2s ² , 2p ²
Na	11	2	2	6	1		1s ² , 2s ² , 2p ⁶ , 3s ¹
Cl	17	2	2	6	2	5	1s ² , 2s ² , 2p ⁶ , 3s ² , 3p ⁵
Ar	18	2	2	6	2	6	1s ² , 2s ² , 2p ⁶ , 3s ² , 3p ⁶

* The superscripts of the electronic configuration add up to the atomic number.

To calculate bond distances and angles, the key is to rely on the geometry of the bond and calculate the relationship between the distances and angles from standard trigonometric principles. For example, for a molecule with a tetrahedral conformation, such as methane, CH₄, or carbon tetrachloride, CCl₄, the H-C-H angle or Cl-C-Cl angle is the tetrahedral angle of 109.5°. The bond distance is, to a first approximation, the sum of the atomic radii of the two atoms between which the bond is formed.



Solution: To find the formula of this compound, it is necessary to determine the valence of the elements from which it is composed. The valence of an element is the number of electrons that are involved in chemical bonding.

To find the valence of phosphorus and hydrogen, consider their atomic number and electronic configuration.

Hydrogen: Atomic number = 1.

Electronic configuration = $1s^1$

Phosphorus: Atomic number = 15.

Electronic configuration = $1s^2 2s^2 2p^6 3s^2 3p^3$.

The outer electrons are $1s^1$ for hydrogen, and $3p^3$ for phosphorus. It takes one additional electron to fill hydrogen's s orbital; its valence is one. It takes 3 more electrons to fill phosphorus' p orbital; its valence is three. Elements react with the purpose of filling all their orbitals with the maximum number of electrons by either a transfer of electrons or by sharing electrons.

It would take three hydrogen atoms to complete phosphorus' outer orbital. In turn, each electron of phosphorus would serve to complete the outer orbital of each hydrogen atom. This can be pictured in an electron-dot formula as shown above.

In this figure, the X's represent the outer electrons of hydrogen and dots represent the electrons in the outer shell of phosphorus. The formula of this compound is, thus, PH_3 .

• PROBLEM 614

H_2O_3 , hydrogen trioxide, a close relative of hydrogen peroxide, has recently been synthesized. It is extremely unstable and can be isolated only in very small quantities. Write a Lewis electron dot structure for H_2O_3 .


Solution: A Lewis dot structure of a compound shows the arrangement of valence electrons. Valence electrons are defined as an element's outer electrons which participate in chemical bonding. Thus, to write an electron dot struc-

ture for H_2O_3 , calculate the total number of valence electrons.

This can be done by considering the electronic configurations of H (hydrogen) and O (oxygen).

Hydrogen possesses one valence electron while oxygen has 6 valence electrons. In H_2O_3 , a total of $(2)1 + (3)6 = 20$ valence electrons are involved.

With this in mind, the Lewis electron dot structure becomes



Notice, you have represented the required 20 valence electrons.

• PROBLEM 615


Using electron-dot notation, show for each of the following the outer shell electrons for the uncombined atoms and for the molecules or ions that result:





- (a) $\text{H} + \text{H} \rightarrow$ hydrogen molecule
- (b) $\text{Br} + \text{Br} \rightarrow$ bromine molecule
- (c) $\text{Br} + \text{Cl} \rightarrow$ bromine chloride
- (d) $\text{Si} + \text{F} \rightarrow$ silicon fluoride
- (e) $\text{Se} + \text{H} \rightarrow$ hydrogen selenide
- (f) $\text{Ca} + \text{O} \rightarrow$ calcium oxide

Solution: When electrons are transferred from one atom to another, ions are formed, which gives rise to ionic bonding. Two atoms, both of which tend to gain electrons, may combine with each other by sharing one or more pairs of electrons. These two atoms form a covalent bond.

To solve this problem, one must know the number of valence electrons, in each of the atoms in the equations. The valence number reflects the combining capacity of an atom. Next, one must know which atoms combine to form ionic bonds and which form covalent bonds. The only ionic bond formed in these equations is for $\text{Ca} + \text{O}$; the other bonds are covalent, and electrons are shared to form an isoelectronic electron cloud such as a noble gas.

Thus,





• PROBLEM 616

The sulfate ion consists of a central sulfur atom with four equivalent oxygen atoms in a tetrahedral arrangement. Keeping in mind the octet rule, draw the electronic structure for the ion. What should the internal O-S-O bond angle be?



Figure a



Figure b



Figure c




Figure d

Solution: The formula for the sulfate ion is SO_4^{2-} . To write the electronic structure of the ion in this problem, one must consider the definition of valence and the octet rule. Valence electrons refer to those outer electrons that participate in chemical bonding. The octet rule states that for stability, there can be no more than eight electrons in the outer orbit of an atom, either as a re-

sult of transfer or sharing. Thus, after determining the number of valence electrons present, arrange them so that they obey the octet rule. The electronic configuration of sulfur is $1s^2 2s^2 2p^6 3s^2 3p^4$. It has 6 valence electrons; they are found in the 3s and 3p orbitals. The electronic configuration of oxygen is $1s^2 2s^2 2p^4$, which means it also has six valence electrons; the first orbital contains only 2 electrons. The electronic structure of each of these atoms can be represented as shown in figure a and figure b, where x's and dots indicate valence electrons. The electronic structure of the ion can be pictured as figure c. Here, one can see that the sulfur and two of the oxygen atoms are surrounded by eight electrons. Because two of the oxygen atoms are only surrounded by seven electrons, each of these atoms possesses a negative charge. This structure may also be drawn as shown in figure d.

For greatest stability the oxygen atoms must take positions as far apart as possible. It has been shown that when four atoms surround a fifth the most stable arrangement is tetrahedral. In this arrangement, the O-S-O bonds form a 109.5° angle.


IONIC AND COVALENT BONDING

• PROBLEM 617


Distinguish a metallic bond from an ionic bond and from a covalent bond.

Solution: The best way to distinguish between these bonds is to define each and provide an illustrative example of each.

When an actual transfer of electrons results in the formation of a bond, it can be said that an ionic bond is present. For example,



When a chemical bond is the result of the sharing of electrons, a covalent bond is present. For example:



ELECTRONEGATIVITY

• PROBLEM 622

What is the meaning and significance of the Pauling electronegativity scale?

Solution: In any chemical bond, electrons are shared between the bonding atoms. In covalent bonds, the valence electrons are shared almost equally - the shared electrons spend about the same amount of time with each atom. In ionic bonds, the valence electrons are monopolized by one atom. The degree to which a bond will be ionic or covalent is dependent upon the relative electron-attracting ability of the bonding atoms.

The Pauling electronegativity scale provides a measure for the relative electron-attracting abilities or electronegativity of each element. The most electronegative element, fluorine, is assigned the highest number, 4.0. The least electron attracting elements (and consequently, those most willing to lose electrons), cesium and francium, are assigned the lowest number, 0.7. Numbers are assigned to the remaining elements so that no element has a higher number than an element more electronegative than it. Furthermore, the numbers are assigned so that the differences between the electronegativities of two elements is indicative of the ionic quality of the bond that forms between them. In particular, if the difference is 1.7 or greater, then the shared electrons are monopolized by the more electronegative atom to such a great extent that the bond is said to be ionic. If the difference is less than 1.7, the bond is covalent.

Thus, when given a compound, the Pauling electronegativity scale can be used to determine the species that acts as the electron donor, the species that acts as the electron acceptor, and the degree of electron polarization of the bond.

• PROBLEM 623

Classify the bonds in the following as ionic, ionic-covalent, covalent-ionic and covalent: I_2 , $LiCl$, $MgTe$, Cl_2O , H_2S .

Solution: The bonds in these molecules can be classified by considering the electronegativities of the atoms involved.

The electronegativity of any element is the tendency of that element to attract electrons. The greater the

• PROBLEM 625

What is the difference between electronegativity and electron affinity? Use specific examples.

Solution: Electron affinity represents the energy released when an electron is added to an isolated neutral gaseous atom. Electronegativity is the tendency of an element to attract electrons, not in the gaseous state, but from a nucleus joined to it by a covalent bond. The electron affinity is a precise quantitative term while the electronegativity is a more qualitative concept.

• PROBLEM 626

Assuming the ionization potential of sodium is 5.1 eV and the electron affinity of chlorine is 3.6 eV, calculate the amount of energy required to transfer one electron from an isolated sodium (Na) atom to an isolated chlorine (Cl) atom.

Solution: Ionization potential is the amount of energy required to pull an electron off an isolated atom. Electron affinity is the amount of energy released when an electron is added to an isolated neutral atom. In this problem, one must add energy to remove an electron from Na and energy will be released upon the addition of an electron to Cl. 5.1 eV are needed to expel an electron from Na and 3.6 eV are released when Cl accepts an electron. Thus, the amount of energy required for the overall process to occur is the difference between the ionization potential of Na and the electron affinity of Cl.

The energy necessary to be added to the system for this reaction to occur is $5.1 \text{ eV} - 3.6 \text{ eV}$ or 1.5 eV.

• PROBLEM 627

It is a fact that the second ionization potential of alkali atoms falls off more rapidly with increasing atomic number than does the first ionization potential. Why?

Solution: The first ionization potential is the energy necessary to remove one electron from a neutral atom in the gaseous state. After this first electron is removed, the atom becomes an ion with a +1 ionization state. Second ionization potential refers to the energy necessary to remove an electron from the +1 ion in the gaseous state.

As the atomic numbers of elements increase the atomic radii of the atoms increase. The atomic radius determines the distance from the nucleus to the outermost electron of the atom.

The first and second ionization potentials should fall off as the atomic radius increases. As the radius increases, the outer electron, which is the one that will be removed, is at a greater distance from the nucleus. Thus, the "pull" on it by the nucleus, is less than that of an electron closer to the nucleus. Hence, less energy should be required to remove an electron that is far away from the nucleus than one that is close.

In considering the second ionization potential, one has an added factor involved; the element is charged. The concentration of charge will be decreased as the atomic radius increases. This increased diffusion of charge with the increased atomic number is the reason for the added fall off of the second ionization potential as compared to first ionization potential.

• PROBLEM 628

40.0 kJ of energy is added to 1.00 gram of magnesium atoms in the vapor state. What is the composition of the final mixture? The first ionization potential of Mg is 735 kJ/mole and the second ionization potential is 1447 kJ/mole

Solution: Ionization potential may be defined as the energy required to pull an electron away from an isolated atom. The second ionization potential is the amount of energy required to pull off a second electron after the first has been removed.

The composition of the final mixture is determined by calculating the number of electrons that will be removed from the magnesium ions. To do this one must determine the number of moles of Mg present in 1 g. From this one can determine the number of electrons that will be liberated by using the values for the first and second ionization potentials of Mg.

The atomic weight of Mg is 24.3. Since moles = grams/atomic weight, there are in 1 gram of Mg, $1/24.30$ or 4.11×10^{-2} moles present. The first ionization potential of Mg is 735 kJ/mole. Therefore, 4.11×10^{-2} moles of Mg requires 4.11×10^{-2} moles \times 735 kJ/mole or 30.2 kJ to ionize all of the atoms once. 40 kJ was added to the system leaving 40 kJ - 30.2 kJ or 9.8 kJ to remove the second electron. If one has 9.8 kJ and 1447 kJ/mole is required to remove the second electrons, then

$$\frac{9.8 \text{ kJ}}{1447 \text{ kJ/mole}} = 6.77 \times 10^{-3} \text{ moles of atoms can}$$

When the following reaction



occurs, 26 Kcal/mole are released. The bond energies of F_2 and Cl_2 are 37 and 58 Kcal/mole, respectively. Calculate the dissociation energy of the ClF bond.

Solution: The dissociation energy is the amount of energy needed to break the chemical bonds of a molecule. The reaction for the dissociation of 2ClF bonds is



This is the opposite of the reaction described in the problem, therefore, 26 Kcal/mole are consumed. The amount of energy consumed is equal to twice the dissociation energy of ClF (because 2 moles of ClF are present) subtracted from the sum of the dissociation energies of F_2 and Cl_2

$$\Delta H = \left(\Delta H_{\text{F}_2} + \Delta H_{\text{Cl}_2} \right) - 2\Delta H_{\text{Cl-F}} = -26 \text{ Kcal/mole}$$


$$26 \text{ Kcal/mole} = 2\Delta H_{\text{Cl-F}} - 37 \text{ Kcal/mole} - 58 \text{ Kcal/mole}$$

$$2\Delta H_{\text{Cl-F}} = 26 \text{ Kcal/mole} + 37 \text{ Kcal/mole} + 58 \text{ Kcal/mole}$$

$$= \frac{121}{2} \text{ Kcal/mole} = 60.5 \text{ Kcal/mole.}$$

Thus, 60.5 Kcal are needed to break one mole of Cl-F bonds.

Based upon the following thermochemical data, show that ozone, O_3 , is considerably more stable than a cyclic structure would suggest. The enthalpy for the O-O bond is approximately 33 Kcal/mole.




Solution: The bond energy is the amount of energy needed to separate atoms joined in a chemical bond. The greater the bond energy the more stable the bond. The stability of a molecule is proportional to the sum of the bond energies of the bonds in the molecule.


If O_3 assumes a cyclic conformation, there are 3 O-O bonds present. Thus, the stability of this structure is proportional to 3 times the bond energy of an O-O bond, 33 Kcal/mole.

$$\text{stability of the cyclic structure} = 3 \times 33 \text{ Kcal/mole or } 99 \text{ Kcal/mole}$$

One can calculate the actual stability of O_3 by using the data supplied in the problem. The overall reaction for the dissociation of O_3 is





One can derive this equation from the two given in the problem. The reverse of the equation and the Δ dissociation for the formation of O_3 is



If one multiplies the second equation by 3/2 it and its ΔH becomes:



One obtains the overall equation (i) by adding together equations (ii) and (iii). One finds the overall ΔH by adding together the ΔH 's of these reactions.



The actual stability of O_3 is, therefore, proportional to 144.5 Kcal/mole. This is greater than that which would be expected if O_3 had a cyclic structure.

• PROBLEM 633

What is the explanation for the following trends in lattice energies?

NaF	- 260 Kcal/mole	NaCl	- 186 Kcal/mole
NaCl	- 186 Kcal/mole	KCl	- 169 Kcal/mole
NaBr	- 177 Kcal/mole	CsCl	- 156 Kcal/mole

Solution: Lattice energy is an indication of the stability of the species.

In the first column, since the sodium (Na) ion is common to all species, one should concentrate on F^- , Cl^- ,

this problem (1) to (5), the reaction becomes (1) + $\frac{1}{2}$ (2) + (3) + (4) + (5). The $\frac{1}{2}$ value for (2) is derived from the fact that one mole of NaF is required, but, in (2), 2 moles of F(g) atoms are generated. Thus, take $\frac{1}{2}$ this number.

Therefore, heat of reaction is:

$$\Delta H = \Delta H(1) + \frac{1}{2}\Delta H(2) + \Delta H(3) + \Delta H(4) + \Delta H(5)$$

$$= 26 + 18.3 + 120 - 83.5 - 216.7 = -135.9 \text{ Kcal/mole.}$$

BOND LENGTH AND ANGLES

• PROBLEM 635

Calculate the interatomic nonbonded distance between 2 bromine nuclei in a molecule of carbontetrabromide (CBr_4). The C-Br distance is 1.94 Å.



Diagram A




Diagram B

Solution: From diagram A, one can see that CBr_4 is in a tetrahedral conformation. Thus, one knows that the Br-C-Br angle is equal to 109.5° . As such, diagram B is then true.

One is solving for the length of $\overline{\text{Br}_1\text{Br}_2}$.

Angle $\text{Br}_1\text{-C-A}$ is $109.5^\circ/2$ or 54.75° , since one can let line $\overline{\text{CA}}$ be an angle bisector. To find $\overline{\text{Br}_1\text{A}}$, note that $\sin 54.75 = \overline{\text{Br}_1\text{A}}/1.94 \text{ Å}$. Therefore,

$$\overline{\text{Br}_1\text{A}} = 1.94 \text{ Å} \times \sin 54.75$$

$$\overline{\text{Br}_1\text{A}} = 1.94 \text{ Å} \times .8166 = 1.58 \text{ Å}$$

structure of the trans form as described in the problem is:

One is asked to find the distance from O_2 to H. This can be done by solving for the length of O_2C and CH. The sum of the squares of these two lengths is equal to the square of the length of O_2H . This can be done through the use of trigonometry. To find O_2C , one must find NA and AC. O_2C is equal to the sum of O_2N , NA, and AC. The measure of angle ONA is equal to $180^\circ - 118^\circ$ or 62° , since they are supplementary.

$$\cos 62^\circ = 0.469 = \frac{NA}{NO_1} = \frac{NA}{0.146 \text{ nm}} \left(\frac{\text{adjacent}}{\text{hypotenuse}} \right)$$

$$0.469 = \frac{NA}{0.146 \text{ nm}}$$

$$NA = 0.069 \text{ nm}$$

$$O_2A = NA + O_2N = 0.069 \text{ nm} + 0.120 \text{ nm} = 0.189 \text{ nm}$$

$\overline{AC} = \overline{O_1B}$, since they are the opposite sides of a rectangle.

$$\angle BO_1H = 90^\circ - \angle HO_1A \quad (\angle = \text{angle})$$

$$\angle NO_1A = 90^\circ - \angle O_1NA = 90^\circ - 62^\circ = 28^\circ$$

$$\angle HO_1A = \angle HO_1N - \angle NO_1A = 105^\circ - 28^\circ = 77^\circ$$

$$\angle BO_1H = 90^\circ - 77^\circ = 13^\circ \text{ from above.}$$

$$\cos 13^\circ = \frac{\overline{OB}}{\overline{OH}} = 0.974$$

$$\frac{\overline{O_1B}}{\overline{O_1H}} = \frac{\overline{O_1B}}{0.098 \text{ nm}} = 0.974$$

$$\overline{O_1B} = (0.098 \text{ nm}) \times (0.974) = 0.095 \text{ nm}$$

$$\text{Thus, } \overline{AC} = 0.095 \text{ nm.}$$

Solving for $\overline{O_2C}$:

$$\overline{O_2C} = \overline{O_2N} + \overline{NA} + \overline{AC}$$

$$= 0.120 \text{ nm} + 0.069 \text{ nm} + 0.095 \text{ nm} = 0.284 \text{ nm}$$

$$\overline{HC} = \overline{OA} - \overline{BH}$$

$$\sin \angle ONA = \frac{\overline{O_1A}}{\overline{NO_1}}$$

$$\angle ONA = 62^\circ$$

charge distribution in the molecule where one atom is partially positive and the other is partially negative. The degree of polarity is measured by finding the difference in the abilities of the two atoms to attract electrons. This tendency to accept electrons is called the electronegativity. The greater the electronegativity difference, the greater the degree of polarity.

From the table of electronegativity values, the following electronegativities can be obtained:

	<u>Compounds</u>	<u>Electronegativity Difference</u>
(1)	HCl	$3.0 - 2.1 = .9$
	HBr	$2.8 - 2.1 = .7$
(2)	H ₂ O	$3.5 - 2.1 = 1.4$
	H ₂ S	$2.5 - 2.1 = .4$
(3)	BrCl	$3.0 - 2.8 = .2$
	IF	$4.0 - 2.5 = 1.5$

In pair (1) (HCl and HBr), HCl has the larger electronegativity. Hence, HCl has a greater degree of polarity than HBr. For the same reasons, H₂O in pair (2) and IF in pair (3) have the greater degrees of polarity.

● **PROBLEM 642**

Of the following pairs, which member should exhibit the largest dipole moment. Use the data from the accompanying table. (a) H-O and H-N; (b) H-F and H-Br; (c) C-O and C-S.

Pauling electronegativities
(H = 2.1)

[illegible]

The values given in the table refer to the common oxidation states of the elements. For some elements variation of the electronegativity with oxidation number is observed, for example, Fe(II) 1.8, Fe(III) 1.9; Cu(I) 1.9, Cu(II) 2.0; Sn(II) 1.8, Sn(IV) 1.9.

(a) Net dipole moment

$$= (4.80 \times 10^{-10} \text{ esu}) (2 \times 10^{-8} \text{ cm})$$

$$= 9.6 \times 10^{-18} \text{ esu-cm.}$$

$$1 \text{ Debye} = 1 \times 10^{-18} \text{ esu-cm.}$$

$$\frac{9.6 \times 10^{-18} \text{ esu-cm}}{1 \times 10^{-18} \text{ esu-cm/Debye}} = 9.6 \text{ Debyes}$$

(b) Net dipole moment

$$= (2) (4.80 \times 10^{-10}) (2 \times 10^{-8})$$

$$= 19.2 \times 10^{-18} \text{ esu-cm}$$

Converting to Debyes


$$\frac{19.2 \times 10^{-18} \text{ esu-cm}}{1 \times 10^{-18} \text{ esu-cm/Debye}} = 19.2 \text{ Debyes.}$$

• PROBLEM 644

The net dipole moment of water is 1.84 debyes and the bond angle is 104.45° . What moment can be assigned to each O-H bond?




(a)




(b)



(c)



(d)



(e)

Solution: A dipole is defined as a molecule which is electrically unsymmetrical - that is, the centers of positive and negative charges are not located at the same point within the molecule.

Quantitatively, a dipole is described by its dipole moment, which is equal to the charge times the distance between the positive and negative centers. The decrease in dipole moment will reflect a decrease in either the value of the charges or the distance between them. It is given that the bond length increases in this sequence, which means that distance increases. This suggests an increase in dipole moment within the sequence. But the dipole moments decrease, this means that the charge must be decreasing. This can be explained by noting that the electronegativity values are decreasing within the sequence. Electronegativity measures the tendency of an atom to attract shared electrons in a molecule. By attracting these electrons, it develops a negative charge. If the electronegativity decreases, so does the charge, which would then account for the decrease in dipole moment.

CHAPTER 18

QUANTUM CHEMISTRY

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 613 to 669 for step-by-step solutions to problems.

The name "quantum" comes from the concept that an electron (or any other particle) can only have discrete energies. Most values of the energy are forbidden so that if an electron changes energy it must do so in a quantum jump.

All electrons in the atoms can be described by four quantum numbers and no two electrons can have the same set of quantum numbers. The quantum numbers are, for historical reasons, called respectively n , l , M_l , and M_s . n is the principal quantum number, l is the orbital quantum number, M_l is the magnetic quantum number, and M_s is the spin. The allowed quantum numbers are as follows:

Quantum No.	Permitted Values
n	1, 2, 3,... etc.
l	0, 1, 2, ... $n - 1$
M_l	$+l$ to $-l$
M_s	$+1/2$, $-1/2$ only

In Chapter 17, the electronic configuration designations, for example $1s^2$, $2s^2$, $2p^2$ for carbon, are, in fact, a tabulation of the quantum numbers of each of the electrons in the atom. Tables 1 and 2 show the relationship between the designation of Chapter 17 and the quantum numbers associated with each electron. In Table 1 the following electronic configuration for neon is tabulated by the rules discussed in Chapter 17.

Table 1: Electronic Configuration of Neon

Element	At. No.	1s	2s	2p	3s	3p	Electronic Configuration
Ne	10	2	2	6			$1s^2, 2s^2, 2p^6$

Table 2: Quantum Numbers for Electrons in Neon Atom

n	l	M_l	M_s	Designation of Chapter 17
1	0	0	+1/2	1s
1	0	0	-1/2	1s (two total 1s electrons)
2	0	0	+1/2	2s
2	0	0	-1/2	2s (two total 2s electrons)
2	1	+1	+1/2	2p
2	1	+1	-1/2	2p
2	1	0	+1/2	2p
2	1	0	-1/2	2p
2	1	-1	+1/2	2p
2	1	-1	-1/2	2p (six total 2p electrons)

The "exclusion" of two electrons having the same set of quantum numbers is called the Pauli exclusion principle. The order in which the quantum numbers occur in the ground, or lowest, energy state is governed by Hund's rule. This rule is outlined in detail in Problem 650.

Two principal types of bonds, sigma (σ) and pi (π) bonds, can be formed in a molecular orbital. The sigma bond is formed when the molecular orbital (of shared electrons) is symmetrical around the nucleus while the pi bond is formed from parallel p orbitals. A succinct description of σ and π bonds is given in Problem 657, and sketches of these molecular orbitals are shown in the problems which follow.

An early attempt to describe the quantum nature of electrons was undertaken by Niels Bohr, and his model is called the Bohr atom. It is simplistic by modern understanding but, for his time, represented a giant step forward in understanding atomic structure. Bohr postulated that the hydrogen atom consisted of a nucleus and a single electron in orbit around the nucleus much as a planet of the solar system orbits the sun. The Coulombic attractive forces are balanced by the centrifugal force of the electron in orbit.

$$e^2/r^2 = mv^2/r \quad 18-1$$

Where e = the charge on one electron, r = distance between the nucleus and the electron, and mv^2/r = centrifugal force. The quantum nature of the electronic energy was introduced by postulating that the angular momentum could equal only quantized values.

$$mvr = nh/2\pi \quad 18-2$$

where h is Planck's constant (6.626×10^{-27} erg sec) and n is the principal quantum number. From Bohr's theory it is possible to calculate the orbital radius from each quantum number n .

The quantum theory indicates matter has both wave and particle characteristics. The deBroglie equation relates the momentum of a particle to its wavelength.

$$\lambda = h/p \quad 18-3$$

where λ is the wavelength and p the momentum ($m \cdot v$) of a particle. An electron, because of its wave character, is not a particle as hypothesized by Bohr. Its position and energy are uncertain as specified by the Heisenberg uncertainty principle. The Heisenberg uncertainty principle states that it is impossible to simultaneously determine the exact position and momentum of a particle and is given by:

$$(\Delta x)(m\Delta V_x) \geq h/4\pi \quad 18-4$$

where $m\Delta V_x$ is equal to the momentum in the x direction and x is equal to the position of the particle.

The energy of a quantum of light or other electromagnetic radiation is given by the equation

$$E = h\nu = \frac{hc}{\lambda} \quad 18-5$$

where ν is the frequency, c is the speed of light, and λ is the wavelength. Because of the quantum nature of energies which an electron can possess, when it moves from one energy state to another, it emits or absorbs energy of a wavelength given by Equation 18-5. This principle forms the basis of the science of atomic spectroscopy, which is the study of atomic structure from light emissions. A diagram of the energy levels showing relative energies for an atom and one electron is shown in Problem 692.

Step-by-Step Solutions to Problems in this Chapter, "Quantum Chemistry"

PAULI EXCLUSION PRINCIPLE, HUND'S RULE AND ELECTRONIC CONFIGURATION

• PROBLEM 646

Explain the following: Pauli exclusion principle and Hund's rule.

Solution: An atom is described by four quantum numbers: the principal quantum number N , the angular momentum quantum number l , the magnetic quantum number m , and the spin quantum number s . According to the Pauli exclusion principle, no two electrons in an atom can have the same four quantum numbers. If two electrons did, they would have identical fingerprints, a situation forbidden by nature.

Each orbital can accommodate a maximum of two electrons. With this in mind, Hund's rule states that once an electron is in an orbital, a second electron will not enter into that same orbital if there exist other orbitals in that subshell that contain zero electrons. In other words, all orbitals in a subshell must contain one electron, before a second one can enter. Hund's rule also states that single electrons in their separate orbitals of a given subshell will have the same spin quantum number.

• PROBLEM 647

Discuss the following statement: The Pauli exclusion principle is the main reason why atoms do not collapse to a point.

Solution: The Pauli exclusion principle states that no two electrons in the same atom can be completely identical;

that is, have the same values for all four quantum numbers. The quantum numbers measure the energy level of the electron, the order of increasing distance of the average electron distribution from the nucleus, the angular shape of electron distribution, (the electronic magnetism), and the two possible orientations of electron spin. An electron in an atom is completely described by its four quantum numbers. Inherent in this description is the point of location of an electron at a specific time. If two electrons have the same four quantum numbers, they would have the same point of location at a specific time. This means, therefore, that all the electrons would collapse to a point at a specific time. The Pauli exclusion principle states that no two electrons can have the same four quantum numbers and therefore the atom cannot collapse to a point.

• PROBLEM 648

Given the ground-state oxygen atom, tabulate each of the electrons by its quantum number.

Solution: Begin by writing the electronic configuration of the ground-state oxygen atom. An electron is in its ground state when it is in its lowest energy level, this is the normal state of the electrons. The atomic number of oxygen is 8. Thus, the oxygen atom has 8 electrons. The electronic configuration of the oxygen atom is $1s^2 2s^2 2p^4$. The quantum numbers are N , l , M , and s . N is the principal quantum number which, in the electronic configuration, is the integer in front of the subshells (s and p) and corresponds to the energy level occupied by the electron. The other quantum numbers can be found once N is determined, l is equal to $0, 1, 2, \dots, N-1$, depending upon N . M is equal to $+l$ to $-l$ and S is equal to $+\frac{1}{2}$ or $-\frac{1}{2}$. l is the orbital quantum number which denotes the subshell and angular shape of electron distribution. M is the magnetic quantum number and s denotes the spin of electron. All four are needed to describe each electron. To tabulate the electrons according to their quantum numbers proceed as follows: the electron configuration is $1s^2 2s^2 2p^4$. Take the first shell, or $N = 1$, $1s^2$. The superscript indicates that 2 electrons are present in this shell. Thus, 2 sets of quantum numbers are needed. For both electrons, $N = 1$. Because $l = 0 \dots n - 1$, $l = 0$ for both electrons in the s subshell. $M + l$ to $-l$, thus, $M = 0$ for both also. The four quantum numbers for each electron must be different. Therefore, one electron has a spin of $+\frac{1}{2}$ and one has a spin of $-\frac{1}{2}$. The sets of quantum numbers for these two electrons are $1, 0, 0, +\frac{1}{2}$ and $1, 0, 0, -\frac{1}{2}$. For the second shell, $N = 2$. There are two subshells, s and p . Consider the s subshell first. In the $2s^2$ orbital, $N = 2$ and there are two electrons to be described. For both electrons $l = 0$, because both are in the s subshell. $M = 0$, because $M = +l$ to $-l$. For one electron $s = +\frac{1}{2}$, for the other $s = -\frac{1}{2}$. The sets of quantum numbers for these two electrons

Since the atomic number of nitrogen is 7 and it is a neutral atom, there are 7 electrons to place in atomic orbitals. The two orbitals of lowest energy are the 1s and the 2s. These are filled according to the Pauli Exclusion Principle - 2 electrons of unpaired spin in each orbital. The orbitals with the lowest energy of those remaining are the three 2p orbitals, $2p_x$, $2p_y$, $2p_z$. Hund's rule dictates that the remaining three electrons enter these orbitals singly and with parallel spin.

The ground state electronic distribution for nitrogen can therefore be pictured as shown in the accompanying Figure.

• PROBLEM 652

Predict the total spin for each of the following electronic configurations: (a) $1s^2 2s^1$; (b) $1s^2 2s^2 2p^1$; and (c) $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^2$.




Figure A

electrons	↑↓	↑↓
orbitals	1s	2s

Figure B

electrons	↑↓	↑↓	↑	↑	↑
orbital	1s	2s	2p		

Figure C

sum of the electrons) and the net positive charge (the atomic number, which is equal to the number of protons) For hydrogen, ($Z = 1$) $1s^1$, the net negative charge is -1 and the net positive charge is $+1$, thus, the atom is neutral and no ionic symbol is used. One uses this method to determine the ionic symbols for the atoms described in the problem.

(a) ($Z = 11$) $1s^2 2s^2 2p^6$. The net negative charge is equal to $(-2) + (-2) + (-6)$ or -10 . The net positive charge is 11 . The net charge on the atom is $11 - 10 = +1$. From the periodic table one sees that this atom is Na^+ , because the atomic number of sodium is 11 .

(b) ($Z = 25$) $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5$. Total number of electrons in configuration is $2 + 2 + 6 + 2 + 6 + 5 = 23$. The net charge on the atom is, then $25 - 23$ or $+2$. From the periodic table one can determine that this atom is Mn^{+2} .

(c) ($Z = 29$) $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10}$. Total number of electrons in configuration = $2 + 2 + 6 + 2 + 6 + 10 = 28$. $Z = 29$ = number of electrons in a neutral atom. Difference: $29 - 28 = +1$. The atom is Cu^{+1} .

• PROBLEM 655

Assume that 90% of the electron density is representative of the volume of the atom. The following atomic radii have been obtained for 7 elements across the second period of the periodic table and for six elements down through the first family:

H						
0.37						
Li	Be	B	C	N	O	F
1.23	.89	.80	.77	.74	.74	.72
Na						
1.57						
K						
2.03						
Rb						
2.16						
Cs						
2.35						

Explain the observed trends of these atomic radii.

Solution: Atomic radii increase going down a family, but decrease going across a period. To explain this, one must consider the change in the elements in going across a period or down a family and see how it could affect atomic radii. In moving across a period, the atomic number increases, which means the nuclear positive charge increases. But the principal quantum number of the outside electrons remains constant. This means that the outer electrons, assuming all other factors, are constant, will remain at

Ne ($Z = 10$): $1s^2 2s^2 2p^6$ Here, all orbitals contain two electrons each. No electron is unpaired. Thus, Ne is not paramagnetic.

Ca ($Z = 20$): $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2$. Again, all orbitals have two electrons. Therefore, calcium is not paramagnetic.

Al ($Z = 13$): $1s^2 2s^2 2p^6 3s^2 3p^1$ The 3p subshell has only 1 electron for three orbitals. It must be unpaired, as such. It is paramagnetic.

Zn ($Z = 30$): $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2$. Each orbital has two electrons. Thus, no paramagnetism exists.

The order of paramagnetism (increasing) is proportional to the number of unpaired electrons. Thus, H, Al, O and N is the order of increasing paramagnetism.

MOLECULAR ORBITAL THEORY

• PROBLEM 657

What is a sigma bond? What is a pi bond? What are their basic differences?

Solution: A molecular orbital that is symmetrical around the line passing through two nuclei is called a sigma (σ) orbital. When the electron density in this orbital is concentrated in the bonding region between the two nuclei, the bond is called a sigma bond. The covalent bonds in H_2 and HF are sigma bonds.

In the formation of the bonding orbital between two fluorine atoms, the 2p orbitals overlap in a head-to-head fashion to form a sigma bond. However, there is a second way in which half-filled p orbitals of two different atoms may overlap to form a bonding orbital.

If the two p orbitals are situated perpendicular to the line passing through the two nuclei, then the lobes of p orbitals will overlap intensively sideways to form an electron cloud that lies above and below the two nuclei. The bond resulting from this sideways or lateral overlap is called a pi (π) bond; the bonding orbital is called a pi orbital. It differs from a sigma orbital in that it is not symmetrical about a line joining the two nuclei. Pi bonds are present in molecules having two atoms connected by a double or triple bond. The sigma bond has greater orbital overlap and is usually the stronger bond; a pi bond, with less overlap, is generally weaker.

Compare the bond order of He_2 and He_2^+ .

Solution: The bond order, or number of bonds in a molecule, is equal to the difference in the sum of the number of bonding electrons and the number of antibonding electrons divided by two.

$$\text{Bond order} = \frac{\text{no. of bonding electrons} - \text{no. of antibonding electrons}}{2}$$

This means that the number of bonding and antibonding electrons must be determined. There are 2 electrons in He, thus in He_2 there are 4. These electrons are all in the 1s level. For each level, there exists bonding and antibonding orbitals, each of which holds 2 electrons. Thus, in He_2 , 2 electrons are bonding and 2 are antibonding. From this, the

$$\text{Bond order} = \frac{2 - 2}{2} = 0.$$

Thus, there are no bonds in He_2 ; and two He atoms will not bond together to form a molecule of He_2 .

In He_2^+ , one electron is removed from He_2 , which means that there are now three electrons present. They are all in the 1s level. This is the lowest energy level that an electron can assume. The three electrons are distributed so that two are in bonding orbitals and one is in an antibonding orbital. Thus,

$$\text{no. of bonds} = \frac{2 - 1}{2} = 0.5 = \text{bond order}.$$

Because the bond order is not zero, this molecule can form.

Consider an octahedral complex having a univalent negative ion at each vertex of the octagon. Explain why the $d_{x^2-y^2}$ orbital of the central atom is less stable relative to the d_{xy} orbital.

Solution: Both the $d_{x^2-y^2}$ and the d_{xy} orbitals are centered about the x-y plane; hence, the difference in the stability of the two orbitals must be related to their different orientations along the x- and y-axes.




Figure A





Figure B

In the $d_{x^2-y^2}$ orbital, the lobes are directed along the axes; whereas in the d_{xy} orbital, the lobes are directed along the bisectors of the angles between the axes (see Figure A). In the octahedral configuration the ligands (the negatively charged ions electrostatically bound to the central atom) lie on the axes (see Figure B).

Hence, the $d_{x^2-y^2}$ orbital brings the negatively charged electrons in the lobes closer to the negatively charged ligands than does the d_{xy} orbital. Since the negative charge of the electrons repels that of the ligands, and this electrostatic repulsion falls off as the reciprocal of the square of the distance between the charges, the greater degree of repulsion between the $d_{x^2-y^2}$ orbital and the ligands, relative to the repulsion between the d_{xy} orbital and the ligands, decreases the stability of the $d_{x^2-y^2}$ orbital.

How would one expect the bond strength of NO to compare with that of O₂?



Solution: The molecular orbital is the sum of the atomic orbitals. For example, when two atomic orbitals are combined, two molecular orbitals are formed. One orbital is called a bonding orbital and the other is an anti-bonding orbital. The bonding orbital is at a lower energy level than the antibonding orbital. If possible, electrons seek out the bonding orbital rather than the antibonding orbital. The orbitals in the production of H₂ from H atoms can be visualized as shown in the accompanying figure.

The greater the number of antibonding orbitals in a molecule, the weaker the bond. Bond order is a quantity that indicates the strength of bonding orbitals. It is defined as half the number of bonding electrons minus half the number of antibonding electrons. Thus, the higher the bond order, the stronger the chemical bond.

To compare the bond strength of NO with O₂, compare their bond orders. To do this, consider the total number of valence electrons in each element. Valence electrons are the outer electrons, which participate in bonding. For NO, the total number of valence electrons is $3 + 4 = 7$. There exist 3 bonding p orbitals, which accommodate 6 electrons. The 1 unpaired electron must be in an antibonding orbital. This means that the bond order of NO is $\frac{1}{2}(6) - \frac{1}{2}(1) = 3 - 0.5 = 2.5$.

In O₂ there is a total of 8 valence electrons. This means 2 electrons must be in antibonding orbitals, since the 3 bonding orbitals can accommodate only 6 electrons. The bond order of O₂ is $\frac{1}{2}(6) - \frac{1}{2}(2) = 2$. The bond order of NO is higher, which means, its chemical bond is stronger than that of O₂.

the geometry of the molecule. A diagram of the bonding resembles Fig. A.

Given that BCl_3 is planar, and since sp^2 hybridization yields a planar structure, B has sp^2 hybridized bonding with angles of 120° (Fig. B).

EARLY QUANTUM CHEMISTRY

• PROBLEM 663

The accompanying figures show a cathode-ray tube (figure A) and the deflection of an electron in a region of magnetic field (figure B). If deflection in the tube is being produced by the magnetic field alone, predict the effect on the observed deflection by increasing (a) mass of particles, (b) velocity of the particles, (c) magnetic field, and (d) charge on the particles.




Figure A

Solution: The magnetic field, charge, velocity, mass and the deflection of a particle are related by the following equation,

$$Hev = \frac{mv^2}{r}$$

where H is the strength of the magnetic field, e the

region. During a particular run of this experiment, an electron enters the magnetic field at a distance of 30 cm away from the detecting screen and travels 25 cm after leaving the field. If the strength of the magnetic field is 0.18 g/coulomb-sec and the beam deflection on the screen is 4.0 mm, what is the velocity of the electrons in the beam?




Figure A




Figure B

Solution: We are looking for a relationship between the magnetic field strength H , the deflection of the electron and the electron velocity v . Such a relationship is provided by the equation of motion of an electron in a magnetic field, $v = He r / m$, where e is the electronic charge ($e = 1.602 \times 10^{-19}$ coulomb), r is the radius of curvature, H is the strength of the magnetic field, and m is the electron mass ($m = 9.11 \times 10^{-28}$ g). In order to apply this equation to solve for v , we must first determine r .

In the following discussion to determine r , we will make reference to figure B (which is not drawn to scale for reasons of clarity). In the problem, we are given that the distance from the point at which the electron beam leaves the field to the deflecting screen is 25 cm. The point at which the electron leaves the field is D. H and D lie on the same vertical line, hence the distance

from D to the screen is the same as the distance from H to the screen, or $HJ = 25$ cm. Since the electron deflection is small, we can assume that the path of the electron is not far from the path the electron would have in the absence of a field. Hence, we can assume that HI is much smaller than CI and HJ , and that FG is much smaller than FB and BG . If we make this assumption that HI is negligible, then $IJ = HJ - HI \approx HJ = 25$ cm. The distance from the point at which the electron enters the field (point C) to the screen is 30 cm, or $CJ = 30$ cm. Then $CI = CJ - IJ = 30$ cm - 25 cm = 5 cm. Denoting the center of the circle containing the magnetic field by G, we have $CG = GI = \frac{1}{2}CI = 5$ cm/2 = 2.5 cm. Using our assumption on the smallness of HI , we obtain $GH = GI - HI \approx GI = 2.5$ cm. In summary, so far we have obtained the following distances:

$$CG = GI \approx GH = 2.5 \text{ cm}, \quad IJ \approx HJ = 25 \text{ cm}.$$

The strategy from this point on will be as follows. We will use the similarity of triangle GJE and triangle GHD to determine HD. Then, using the similarity of triangle CHD and triangle CFB, we will obtain the ratio BF/CF , and from this, the radius of curvature, r .

There is a theorem from elementary plane geometry which states that when a triangle is cut by a line parallel to one of its sides, the two resulting triangles are similar. Since HD and JE are both drawn vertically, and are therefore parallel, triangle GJE is similar to triangle GHD. Hence,

$$\frac{GH}{GJ} = \frac{HD}{JE} \quad \text{or}$$

$$\begin{aligned} HD &= JE \times \frac{GH}{GJ} = 4 \text{ mm} \times \frac{2.5 \text{ cm}}{27.5 \text{ cm}} = 0.4 \text{ cm} \times \frac{2.5 \text{ cm}}{27.5 \text{ cm}} \\ &= 0.036 \text{ cm}. \end{aligned}$$

Next we note that, using the theorems of plane geometry, it can be shown that triangle CHD is similar to triangle CGB. Then

$$\frac{HD}{CG} = \frac{CD}{BG}$$

Because $r = BG$, we solve this ratio for BG. We have already determined HD and CG and can find CD by using the Pythagorean Theorem, thus, we can find BG. Solving for CD:

$$\begin{aligned} CD^2 &= CH^2 + HD^2 = (5.0 \text{ cm})^2 + (0.036 \text{ cm})^2 = 25.001 \text{ cm}^2; \\ CD &= 5.00 \text{ cm}. \end{aligned}$$

If one knows the thickness of the foil and the atomic radius of the particles that compose the foil, one can calculate the number of particles which make up the width of the foil. If one assumes cubic packing in the foil, the sum of the lengths of the diameters of the gold atoms will be equal to the thickness of the foil. Proceed as follows:

The radius, and therefore the diameter, of the gold atom is given in terms of angstroms. The thickness of the foil must be converted to the same units.

$$\begin{aligned}\text{The thickness} &= (10^{-4} \text{ in}) (2.54 \text{ cm/in}) (1 \times 10^8 \text{ \AA/cm}) \\ &= 2.54 \times 10^4 \text{ \AA}.\end{aligned}$$

$$\text{Diameter of gold nucleus} = 2 \times \text{Radius} = 2 \times 1.5 \text{ \AA} = 3 \text{ \AA}.$$

The minimum number of gold atoms passed by alpha particles is equal to

$$\frac{2.54 \times 10^4 \text{ \AA}}{3 \text{ \AA}} = 8.466 \times 10^3$$

• PROBLEM 666

A chemist has a piece of foil that is approximately 5×10^4 atoms thick. If an alpha particle must come within 10^{-12} cm of a nucleus for deflection to occur, what is the probability that an alpha particle will be deflected, assuming the nuclei are not directly behind one another? Assume that the area of one atom is 4×10^{-16} cm².

Solution: To calculate the chance of deflection occurring in any one layer of the foil, assume that each layer is one atom thick. The foil is, therefore, 5×10^4 layers thick. If the number of deflections in any one layer is known, it can be multiplied by 5×10^4 to find the total number of possible deflections as an alpha particle moves through the foil. To find the chance of a deflection occurring in one layer, assume that the atoms are spherical. At 10^{-12} cm from the center of the circle, a deflection can occur. The radius of the circle for which a deflection will occur is 10^{-12} cm. The area of a circle is equal to πr^2 where $\pi = 3.14$ and $r = \text{radius}$.

The area of one atom in which deflection can occur is equal to $3.14 \times (10^{-12})^2$ or 3.14×10^{-24} cm². The area of each atom is 4×10^{-16} cm². The thickness of each layer is one atom. Therefore, the chance of getting a deflection in any given layer of atoms is

$$\frac{3.14 \times 10^{-24} \text{ cm}^2}{4 \times 10^{-16} \text{ cm}^2} = 7.85 \times 10^{-9}.$$

What is the de Broglie wavelength of a radium alpha particle (a helium nucleus, He^{2+}) having an energy of 4.8 MeV (million electron volts)?

Solution: The de Broglie wavelength, λ , of a particle of mass, m , moving at speed, v , is given by

$$\lambda = \frac{h}{mv},$$

where h is Planck's constant ($h = 6.6 \times 10^{-27}$ erg-sec). The mass of an alpha particle is 6.6×10^{-24} g.

The energy of an alpha particle is given by $E = \frac{1}{2} mv^2 = 4.8$ MeV. Since 1 MeV is equal to 1.6×10^{-6} erg, we have:

$$\frac{1}{2} mv^2 = 4.8 \times 1.6 \times 10^{-6} \text{ erg.}$$

We will employ this value of the energy in order to obtain the value of mv without having to find the velocity of the particle.

$$\begin{aligned} (mv)^2 &= 2(\frac{1}{2} mv^2) m \\ &= 2(4.8 \times 1.6 \times 10^{-6} \text{ erg})(6.6 \times 10^{-24} \text{ g}) \\ &= 1.0 \times 10^{-28} \text{ erg-g} = 1.0 \times 10^{-28} \text{ g}^2 \text{ cm}^2/\text{sec}^2, \end{aligned}$$


where we have used 1 erg = 1 g - cm²/sec². Taking the square root of both sides gives:

$$\begin{aligned} (mv)^2 &= 1.0 \times 10^{-28} \text{ g}^2 \text{ cm}^2/\text{sec}^2 \\ mv &= 1.0 \times 10^{-14} \text{ g cm/sec.} \end{aligned}$$

Substituting this value into the expression for the de Broglie wavelength gives:

$$\begin{aligned} \lambda &= \frac{h}{mv} = \frac{6.6 \times 10^{-27} \text{ erg-sec}}{1.0 \times 10^{-14} \text{ g-cm/sec}} = \frac{6.6 \times 10^{-27} \text{ g-cm}^2/\text{sec}}{1.0 \times 10^{-14} \text{ g-cm/sec}} \\ &= 6.6 \times 10^{-13} \text{ cm.} \end{aligned}$$

A chemistry student observed each of the following objects: (a) a 10,000 kg truck moving at 27.8 m/sec, (b) a 50 mg flea flying at 1 m/sec, and (c) a water molecule moving at 500 m/sec. The student proceeded to calculate the wavelength in centimeters of each object. What were these wavelengths? $h = 6.626 \times 10^{-34}$ J sec.



Graphs of ψ , ψ^2 , and $4\pi r^2 \psi^2 dr$ for a 3s electron.

Using $0 = 27 - \frac{18Zr}{a_0} + \frac{2Z^2r^2}{a_0^2}$ solve for r .

$$0 = 27 - \frac{18(11)(r)}{(.0529)} + \frac{2(11)^2(r)^2}{(.0529)^2}$$

$$= 27 - 3.74 \times 10^3 r + 8.67 \times 10^4 r^2$$

Using the quadratic equation one finds that r is equal to 0.0092 or 0.034 nm.

• PROBLEM 680

The mass of an electron is 9.11×10^{-31} and Planck's constant = 6.63×10^{-34} J sec. How much energy does an electron possess in its five lowest states, if it is placed in a cube of dimension 50 nm?

Solution: Wave function theory states that the permitted values of the energy of a particle in a box is given as

$$E = \frac{N^2 h^2}{8 m L^2},$$

where E = energy, N = principal quantum number, h = Planck's constant, m = mass of particle, and L = length of box. Substitute in the known values and solve for E . The principal quantum number N corresponds to the energy levels of the electrons. The five lowest energy states are $N = 1$, $N = 2$, $N = 3$, $N = 4$, and $N = 5$, where $N = 1$ is the lowest and $N = 5$ the highest energy level.

Substituting:

$$E = \frac{N^2 h^2}{8m L^2} = \frac{N^2 (6.63 \times 10^{-34} \text{ J sec})^2}{8(9.11 \times 10^{-31} \text{ kg})(5 \times 10^{-8} \text{ m})^2}$$

$$= N^2 (2.41 \times 10^{-23})$$

(Note: $50 \text{ nm} = 5.0 \times 10^{-8} \text{ m}$)

1st energy level (lowest): $N^2 = 1$, $E = 2.41 \times 10^{-23} \text{ J}$

2nd energy level : $N^2 = 4$, $E = 9.64 \times 10^{-23} \text{ J}$

3rd energy level : $N^2 = 9$, $E = 2.16 \times 10^{-22} \text{ J}$

4th energy level : $N^2 = 16$, $E = 3.85 \times 10^{-22} \text{ J}$

5th energy level : $N^2 = 25$, $E = 6.025 \times 10^{-22} \text{ J}$.

• PROBLEM 681

If one increases the angle away from the Z axis, how does the wave function for a $2P_z$ electron change?

Solution: The wave function of a $2P_z$ electron is

$$\Psi = \frac{1}{4\sqrt{2\pi}} \left(\frac{z}{a_0}\right)^{3/2} e^{-Zr/2a_0} r \cos \theta,$$

where Ψ is the wave function and θ = the polar angle. The polar angle measures the angular deviation away from the vertical, or Z , axis. Ψ^2 is proportional to the electron density at any given radius.

One needs to consider only Ψ and θ to answer the question. To see how the wave function, Ψ , changes when the angle from the Z axis is increased, take two values for θ , and see what happens to Ψ . For example, when $\theta = 0$ (that is, looking along the Z axis), $\cos \theta$ is one, the maximum value a cosine can assume. The wave function is dependent upon θ . The cosine θ is at a maximum, then Ψ is at a maximum. Thus, Ψ^2 is also at a maximum and the electron density is greatest when looking along the Z axis. When $\theta = 90^\circ$ (that is, when one is looking 90° away from the Z


axis, i.e. along the X or Y axis), $\cos \theta$ is zero, the minimum value a cosine can assume. This means, therefore, that ψ^2 is not at a minimum, which means that the probability of finding the electron at this location is zero. In summary, when the angle was increased from 0° to 90° , the wave function decreased. Thus, when one increases the angle away from the Z axis, the wave function and the electron density for a $2p_z$ electron decreases.

• PROBLEM 682

Does the electron probability distribution for 1s and 2s support or refute the Bohr picture of the shell?

90%

contour



1s orbital

Schematic representations of the 1s orbital of hydrogen: a cross section of the probable density of the electron cloud; this cross section represents a slice through the middle of the cloud.

spherical node



90% contour

2s orbital

Solution: This question involves the comparison of the Bohr concepts and the electron probability distribution for the 1s and 2s orbitals. Bohr believed that the shells have spherical symmetry with the average distance from the nucleus of the 2s electron being greater than that for the 1s. The electron probability distribution (orbitals) for 1s and 2s shells are, spherical in symmetry with 2s having a greater radius than the 1s. The Bohr picture does agree very well with the electron distribution for 1s. However, the Bohr picture does not describe as accurately the spread-out nature of the true distribution or the node separating the two regions which the 2s electron occupies. This is

• PROBLEM 686

How much energy is emitted by Avogadro number of atoms, if they each emit a light wave of 400 nm wavelength?

Solution: Energy is related to wavelength in the following equation, $E = hc/\lambda$, where E is energy, h Planck's constant (6.626×10^{-34} Js), c the speed of light (3.0×10^8 m/sec) and λ the wavelength. Using this equation, one calculates the energy emitted by each atom. To find the total amount of energy produced by Avogadro's number of atoms, multiply the amount of energy one atom emits by 6.02×10^{23} .

Solving for E : $1 \text{ nm} = 10^{-9} \text{ m}$.

$$E = \frac{hc}{\lambda} = \frac{(6.626 \times 10^{-34} \text{ Js})(3.0 \times 10^8 \text{ m/s})}{(400 \times 10^{-9} \text{ m})}$$

$$= 4.97 \times 10^{-19} \text{ J/atom}$$

Thus, the total amount of energy emitted is

$$E = (4.97 \times 10^{-19} \text{ J/atom})(6.02 \times 10^{23} \text{ atoms/mole})$$

$$= 2.98 \times 10^5 \text{ J}.$$

• PROBLEM 687

The flame test for barium involves placing a barium-containing compound in a Bunsen burner flame. The barium compound then decomposes to give barium atoms which subsequently undergo an electronic transition of energy 3.62×10^{-12} erg. What color flame would this give rise to?

Solution: To solve this problem we use the relationship $E = hc/\lambda$, where E is the energy of the transition, h is Planck's constant, c is the speed of light, and λ is the wavelength of the emitted light. Solving for λ ,

$$\lambda = \frac{hc}{E} = \frac{6.62 \times 10^{-27} \text{ erg-sec} \times 3.0 \times 10^{10} \text{ cm/sec}}{3.62 \times 10^{-12} \text{ erg}}$$

$$= 5.490 \times 10^{-5} \text{ cm}.$$

Converting to \AA ,

$$\lambda = 5.490 \times 10^{-5} \text{ cm} = 5.490 \times 10^{-5} \text{ cm} \times 10^8 \frac{\text{\AA}}{\text{cm}} = 5490 \text{ \AA},$$

which corresponds to a green flame.

Solution: This problem entails the understanding of the photoelectric effect. The photoelectric effect states that when one shines light of a given frequency on a metallic material, electrons will be emitted. The equation: $E = h\nu$ (where E = energy, ν = frequency, and h = Planck's constant) determines the amount of energy produced by a given frequency of light. The ionization potential is the amount of energy needed to cause the electrons to be ejected from the metal. To solve this problem, therefore, one must calculate how much energy is present in light with a wavelength of 700 nm and then compare this to the ionization potential of a Cs atom.

It is given that the wavelength of the light is 700 nm. The frequency of this light is equal to speed of light/wavelength = $(3.00 \times 10^8 \text{ m/s}) / 700 \times 10^{-9} \text{ m}$. Thus,

$$E = (6.63 \times 10^{-34} \text{ J sec}) \frac{(3.00 \times 10^8 \text{ m/s})}{700 \times 10^{-9} \text{ m}}$$

$$= 2.84 \times 10^{-19} \text{ J/photon}$$

1 eV = $1.60 \times 10^{-19} \text{ J}$. The ionization energy is 3.89 eV or (3.89 eV) $(1.60 \times 10^{-19} \text{ J/eV}) = 6.22 \times 10^{-19} \text{ J}$.

Because Cs is sensitive to light with a wavelength of 700 nm, one photon of this light flashed on the metal Cs will cause 1 electron to be emitted from the Cs. Therefore, the number of electrons emitted when $6.22 \times 10^{-19} \text{ J}$, (the ionization potential of an isolated Cs atom), is added to the metallic system is found by dividing $6.22 \times 10^{-19} \text{ J}$ by $2.84 \times 10^{-19} \text{ J}$

$$\text{no. of electrons emitted} = \frac{6.22 \times 10^{-19} \text{ J}}{2.84 \times 10^{-19} \text{ J}} = 2.2 \text{ electrons}$$


Therefore, the maximum number of electrons that can be emitted is 2.

• PROBLEM 692

The accompanying figure shows an energy-level diagram. Make a comparison of the energy of the $n = 5$ and $n = 4$ transition for an electron with a nucleus of $Z = 3$ and the energy of the transition from $n = 2$ to $n = 1$ for an electron with a nucleus of $Z = 2$.

Solution: When an atom is in the ground state, it is at its lowest energy level, $n = 1$. If energy is added, the electrons become "excited" and move to a higher energy level; $n = 2, 3, 4 \dots$. After a time, the electron falls back to lower energy states and releases the energy needed to move it to the higher energy levels. The equation

$$E_b - E_a = Z^2 \frac{e^2}{2a_0} \left(\frac{1}{n_a^2} - \frac{1}{n_b^2} \right),$$



Energy-level diagram showing relative energies for an atom consisting of a nucleus with charge Z^+ plus one electron with various values of n .

where n = the (quantum) energy level, E = energy, e = charge on electron, a_0 = Bohr radius, and Z = atomic number, measures this energy release between energy levels. You are asked to compare the energy releases between two different atoms and energy levels. You do not need to know the actual energy released. Because $e^2/2a_0$ is a constant, it need not be evaluated. (It will cancel out when you compare).

Therefore, you have for $n = 5$ to $n = 4$

$$E_b - E_a = \frac{e^2}{2a_0} \left(\frac{1}{n_a^2} - \frac{1}{n_b^2} \right) Z^2 = \frac{e^2}{2a_0} \left(\frac{1}{16} - \frac{1}{25} \right) \times 9$$

For $n = 2$ to $n = 1$, you have

$$E_b - E_a = \frac{e^2}{2a_0} \left(\frac{1}{n_a^2} - \frac{1}{n_b^2} \right) Z^2 = \frac{e^2}{2a_0} \left(\frac{1}{1} - \frac{1}{4} \right) \times 4$$

By comparison, in the form of a ratio, one has

$$\frac{\frac{e^2}{2a_0} \left(\frac{1}{16} - \frac{1}{25} \right) \times 9}{\frac{e^2}{2a_0} \left(\frac{1}{1} - \frac{1}{4} \right) \times 4} = \frac{\left(\frac{1}{16} - \frac{1}{25} \right) \times 9}{\left(\frac{1}{1} - \frac{1}{4} \right) \times 4} = \frac{.2025}{3}$$

$\frac{.2025}{3} \times 100\% = 6.8\%$. Thus, the energy involved in

the $n = 5$ to $n = 4$ transition is 6.8% as large as compared to the energy emitted in the $n = 2$ to $n = 1$ transition.

• PROBLEM 693

What wavelength of light is needed to excite an electron in a 0.2 nanometer ($1 \text{ nm} = 10^{-9} \text{ m}$) box from the ground state to the second excited state? What wavelength of light is emitted when the same electron falls from the second excited state to the first excited state?

Solution: The solution to this problem requires the application of two equations, one to determine the energies of the various states, from which may be obtained the energy of a transition between states, and one to relate energy to wavelength. The first equation is the energy of a particle-in-a-box,

$$E_n = \frac{n^2 h^2}{8mL^2},$$

where n is the principal quantum number ($n = 1, 2, 3, \dots$), h is Planck's constant ($6.626 \times 10^{-27} \text{ erg-sec} = 6.626 \times 10^{-34} \text{ J-sec}$), m is the mass of the particle, and L is the length of the box. For an electron in a 0.2 nm

$$= 0.2 \times 10^{-9} \text{ m} = 2 \times 10^{-10} \text{ m box,}$$

$$E_n = \frac{n^2 h^2}{8mL^2} = \frac{n^2 (6.626 \times 10^{-34} \text{ J-sec})^2}{8 \times 9.11 \times 10^{-31} \text{ kg} \times (2 \times 10^{-10} \text{ m})^2}$$

$$= 1.506 \times 10^{-18} \text{ J} \times n^2.$$

The second equation relates the energy difference, ΔE , between two states and the wavelength λ : $\Delta E = hc/\lambda$, or $\lambda = hc/\Delta E$, where c is the speed of light.

In the following table, we calculate the energies corresponding to the ground state ($n = 1$), the first excited state ($n = 2$), and the second excited state ($n = 3$).

n	$E_n = n^2 \times 1.506 \times 10^{-18} \text{ J}$
-----	--

1	$E_1 = (1)^2 \times 1.506 \times 10^{-18} \text{ J} = 1.506 \times 10^{-18} \text{ J}$
---	--

2	$E_2 = (2)^2 \times 1.506 \times 10^{-18} \text{ J} = 6.024 \times 10^{-18} \text{ J}$
---	--

3	$E_3 = (3)^2 \times 1.506 \times 10^{-18} \text{ J} = 1.355 \times 10^{-17} \text{ J}$
---	--

In going from the ground state to the second excited state, the electron must absorb a quantum of light of energy

$\Delta E = E_3 - E_1 = 1.355 \times 10^{-17} \text{ J} - 1.506 \times 10^{-18} \text{ J} = 1.204 \times 10^{-17} \text{ J}$. The corresponding wavelength is

$$\lambda = \frac{hc}{\Delta E} = \frac{6.626 \times 10^{-34} \text{ J} \times 3 \times 10^8 \text{ m/sec}}{1.204 \times 10^{-17} \text{ J}}$$

$$= 16.510 \times 10^{-9} \text{ m} = 16.510 \text{ nm}.$$

In falling from the second excited state to the first excited state the electron will emit a quantum of light of energy $\Delta E = E_3 - E_2 = 1.355 \times 10^{-17} \text{ J} - 6.024 \times 10^{-18} \text{ J} = 7.526 \times 10^{-18} \text{ J}$. This corresponds to a wavelength of

$$\lambda = \frac{hc}{\Delta E} = \frac{6.626 \times 10^{-34} \text{ J} \times 3 \times 10^8 \text{ m/sec}}{7.526 \times 10^{-18} \text{ J}}$$

$$= 26.412 \times 10^{-9} \text{ m} = 26.412 \text{ nm}.$$

• PROBLEM 694

Consider an atom containing one electron and having an atomic number Z . It is observed that light of a wavelength of 0.1 nm will excite this atom from the ground state to infinite ionization. Determine the value of Z .

Solution: We know the wavelength λ of the exciting light and the quantum numbers of the ground state ($n_g = 1$) and the ionized state ($n_i = \infty$). We must determine Z , the number of protons in the nucleus of the atom.

There is a relationship to find Z using n_g and n_i . It is provided by the Bohr theory of the atom. It is:

$$\frac{c}{\lambda} = \frac{Z^2 e^2}{2ha_0} \left(\frac{1}{n_g^2} - \frac{1}{n_i^2} \right),$$

where c is the speed of light ($c = 2.9969 \times 10^{10} \text{ cm/sec}$), e is the electronic charge ($e = 4.80 \times 10^{-10} \text{ esu}$), h is Planck's constant ($h = 6.626 \times 10^{-27} \text{ erg-sec}$), and a_0 is the Bohr radius ($a_0 = 0.05292 \times 10^{-7} \text{ cm}$). If we combine some of these constants, we arrive at the value $e^2/2ha_0 = 3.290 \times 10^{15} \text{ sec}^{-1}$. Hence, our expression becomes

$$\frac{c}{\lambda} = Z^2 \left(\frac{e^2}{2ha_0} \right) \left(\frac{1}{n_g^2} - \frac{1}{n_i^2} \right)$$

$$= Z^2 \times 3.290 \times 10^{15} \text{ sec}^{-1} \left(\frac{1}{n_g^2} - \frac{1}{n_i^2} \right).$$

To solve this problem, we must solve this equation for Z and substitute c , n_g , n_i , and λ ($\lambda = 0.1 \text{ nm} = 0.1 \times 10^{-9} \text{ m} = 10^{-10} \text{ m} = 10^{-8} \text{ cm}$) into the resulting equation.

Solving for Z , we obtain

$$Z^2 = \frac{c}{\lambda} \left[3.290 \times 10^{15} \text{ sec}^{-1} \left(\frac{1}{n_g^2} - \frac{1}{n_i^2} \right) \right]^{-1}, \quad \text{or}$$

$$Z = \left(\frac{c}{\lambda \times 3.290 \times 10^{15} \text{ sec}^{-1} \left(\frac{1}{n_g^2} - \frac{1}{n_i^2} \right)} \right)^{\frac{1}{2}}$$

Substituting the values of c , λ , n_g , and n_i into this expression yields

$$Z = \left(\frac{2.9969 \times 10^{10} \text{ cm/sec}}{10^{-8} \text{ cm} \times 3.290 \times 10^{15} \text{ sec}^{-1} \left(\frac{1}{1} - \frac{1}{\infty} \right)} \right)^{\frac{1}{2}} \approx 30,$$

where we used the equality $1/\infty = 0$. Hence, this atom must have 30 protons.

• PROBLEM 695

The Rydberg-Ritz equation governing the spectral lines of hydrogen is $\frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$, where R is the Rydberg constant, n_1 indexes the series under consideration ($n_1 = 1$ for the Lyman series, $n_1 = 2$ for the Balmer series, $n_1 = 3$ for the Paschen series), $n_2 = n_1 + 1, n_1 + 2, n_1 + 3, \dots$ indexes the successive lines in a series, and λ is the wavelength of the line corresponding to index n_2 . Thus, for the Lyman series, $n_1 = 1$ and the first two lines are 1215.56 \AA ($n_2 = n_1 + 1 = 2$) and 1025.83 \AA ($n_2 = n_1 + 2 = 3$). Using these two lines, calculate two separate values of the Rydberg constant. The actual value of this constant is $R = 109678 \text{ cm}^{-1}$.

Solution: The first thing to do is to convert the wavelengths from \AA to more manageable units, i.e. centimeters. Using the relationship $1 \text{ \AA} = 10^{-8} \text{ cm}$, the first two Lyman lines are $1215.56 \text{ \AA} = 1215.56 \times 10^{-8} \text{ cm}$ for $n_2 = 2$, and $1025.83 \text{ \AA} = 1025.83 \times 10^{-8} \text{ cm}$ for $n_2 = 3$. Solving the Rydberg-Ritz equation for R , one obtains

$$R = \left[\lambda \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \right]^{-1},$$

For the first line,

$$R = \left[\lambda \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \right]^{-1} = \left[1215.56 \times 10^{-8} \text{ cm} \left(\frac{1}{1^2} - \frac{1}{2^2} \right) \right]^{-1} \\ = 109689 \text{ cm}^{-1}$$

and for the second line,

$$R = \left[\lambda \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \right]^{-1} = \left[1025.83 \times 10^{-8} \text{ cm} \left(\frac{1}{1^2} - \frac{1}{3^2} \right) \right]^{-1} \\ = 109667 \text{ cm}^{-1}.$$

The first of these is 0.0100% greater than the true value, and the second is 0.0100% less than the true value.

• PROBLEM 696

Wave number ($\bar{\nu}$) are the reciprocals of wavelengths, λ , and are given by the expression $\bar{\nu} = 1/\lambda$. For the hydrogen atom, the Bohr theory predicts that the wave number for the emission line associated with an electronic transition from the energy level having principal quantum number n_2 to that with principal quantum number n_1 is

$$\bar{\nu} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right),$$

where R_H is the Rydberg constant. In what region of the electromagnetic spectrum would there appear a spectral line resulting from the transition from the tenth to the fifth electronic level in hydrogen?

Solution: This problem can be solved by applying the Rydberg formula used for determining the wavenumbers of the spectral lines.

The Rydberg constant is $1.10 \times 10^5 \text{ cm}^{-1}$. $n_2 = 10$ and $n_1 = 5$. Substituting:

$$\bar{\nu} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) = 1.10 \times 10^5 \text{ cm}^{-1} \left(\frac{1}{5^2} - \frac{1}{10^2} \right) \\ = 1.10 \times 10^5 \text{ cm}^{-1} \left(\frac{1}{25} - \frac{1}{100} \right) = 1.10 \times 10^5 \text{ cm}^{-1} (0.04 - 0.01) \\ = 1.10 \times 10^5 \text{ cm}^{-1} \times 0.03 \\ = 3.3 \times 10^3 \text{ cm}^{-1}.$$

This line appears in the infrared region of the spectrum (approximately 20 cm^{-1} to 10^4 cm^{-1}).

• PROBLEM 697

What is the maximum number of emission lines one would expect to see in a molecule containing only the six electronic energy levels depicted in diagram A?




Diagram A




Diagram B

Solution: Emission is observed when an electron falls from an electronic level of higher energy to one of lower energy.

$$\frac{1}{\lambda} = R \left[\frac{1}{(n_1)^2} - \frac{1}{(n_2)^2} \right],$$

where λ is wavelength. The electron falls from an energy level of n_2 to n_1 . The speed of light is related to the frequency in the equation, $v = c/\lambda$.

These equations are combined.

$$(c) \left(\frac{1}{\lambda} \right) = cR \left[\frac{1}{(n_1)^2} - \frac{1}{(n_2)^2} \right] = \frac{c}{\lambda} = v$$

$$\text{Thus, } v = cR \left[\frac{1}{(n_1)^2} - \frac{1}{(n_2)^2} \right]$$

In the Balmer series n_1 is defined as being equal to 2. For successive lines, $n_2 \rightarrow \infty$ and therefore $1/(n_2)^2$ approaches 0. The above equation becomes

$$v = cR \left[\frac{1}{(2)^2} - \frac{1}{(\infty)^2} \right] = \frac{cR}{4}.$$

• PROBLEM 701

Find the difference in characteristic vibrational frequency of a diatomic molecule of mass 20 amu if (a) the mass is equally distributed between the two ends, and (b) one atom of 1 amu is at one end and an atom of 19 amu is at the other.

Solution: The vibrational frequency is frequency of the oscillation between the two atoms of the molecule. It is expressed quantitatively by

$$v = \frac{1}{2} \pi \sqrt{K/M_{\text{eff}}}$$

where v is the vibrational frequency, K a force constant, and M_{eff} the effective mass of the molecule.

$$M_{\text{eff}} = \frac{M_A M_B}{M_A + M_B}$$

where M_A is the mass of one atom and M_B is the other.

(a) If the mass is equally distributed between both atoms, each weighs 10 amu.

$$M_{\text{eff}} = \frac{10 \times 10}{10 + 10} = 5$$

Assume that $K = 1$.

$$\nu = \frac{1}{2\pi} \sqrt{\frac{1}{5}} = 0.0711$$

$$(b) M_{\text{eff}} = \frac{1 \times 19}{20} = \frac{19}{20}$$

$$\nu = \frac{1}{2\pi} \sqrt{\frac{19}{20}} = 0.1632$$

Therefore, the atoms in the molecule described in
b are vibrating more quickly than those in molecule a.

• PROBLEM 702

Calculate the characteristic vibrational frequency of the lowest vibration state for H_2 . Assume $k = 5.1$ newtons/cm and $h = 6.63 \times 10^{-34}$ J-sec.

Solution: Picture a diatomic molecule, such as H_2 , as two balls connected by a spring. The spring represents the chemical bond. When the two balls (atoms) are pulled apart the string acts as a restoring force to bring them back together to the equilibrium separation. When they are compressed, the string tends to force them apart. Such a system, in which the restoring force is proportional to the amount of distortion, is called a harmonic oscillator. The vibrational frequency of the spring going back and forth is given by the formula

$$\nu = \frac{1}{2\pi} \sqrt{k/M_{\text{eff}}}$$

where ν = characteristic vibrational frequency
of the particular bond, k = force constant
and M_{eff} = effective mass for the vibrating atoms:

$$M_{\text{eff}} = \frac{M_A M_B}{M_A + M_B}$$

If the masses of the two atoms are equal, M_{eff} becomes

$$\frac{M^2}{2M} = \frac{1}{2} M.$$

To find the characteristic vibration frequency, use this formula.

To find the M_{eff} , find the mass of one hydrogen atom. The number of grams in one mole (molecular weight) is one. There are 6.02×10^{23} atoms of H in one mole. Thus, the mass of one atom is

$$\frac{1 \text{ g/mole}}{6.02 \times 10^{23} \text{ /mole}} = 1.66 \times 10^{-24} \text{ g.} \quad \text{Thus,}$$

$$M_{\text{eff}} = \frac{(1.66 \times 10^{-24} \text{ g})}{2} = 8.31 \times 10^{-25} \text{ g.}$$

Solving for v :

$$k = 5.1 \frac{\text{newtons}}{\text{cm}} = 5.1 \frac{\text{kg M}}{\text{cm} \cdot \text{sec}^2}$$

$$v = \frac{1}{2\pi} \sqrt{\frac{5.1 \frac{\text{kg M}}{\text{cm sec}^2} \times 1000 \frac{\text{g}}{\text{kg}} \times 100 \frac{\text{cm}}{\text{m}}}{8.31 \times 10^{-25} \text{ g}}}$$

$$= 1.25 \times 10^{14} \text{ sec}^{-1}.$$

Note: 1000 g/kg and 100 cm/m are conversion factors used to obtain the correct units.

• PROBLEM 703

Find the wavelength of light required for the ionization of sodium atoms to occur. The ionization potential of sodium is $8.17 \times 10^{-19} \text{ J}$.

Solution: The energy necessary to ionize a sodium atom is equal to the ionization potential. Here, one is looking for the wavelength of light, which, when flashed upon sodium, will possess an energy equal to the ionization potential. Wavelength is related to energy by the equation $E = hc/\lambda$, where E is the energy, h Planck's constant ($6.626 \times 10^{-34} \text{ J sec}$), c the speed of light ($3.0 \times 10^8 \text{ m/sec}$) and λ the wavelength. One can rewrite this equation as $\lambda = hc/E$ and then substitute the given to solve for λ .

$$\lambda = \frac{(6.626 \times 10^{-34} \text{ J sec})(3.0 \times 10^8 \text{ m/s})}{(8.17 \times 10^{-19} \text{ J})}$$

$$= 2.43 \times 10^{-7} \text{ m} = 2.43 \times 10^{-7} \text{ m} \times \frac{1 \text{ \AA}}{10^{-10} \text{ m}} = 2430 \text{ \AA}.$$

• PROBLEM 704

Find the wavelength (λ) of the first transition, from the first excited state to the ground state in the Lyman and Paschen series. These series show the lines of the emission spectra of H_2 gas. The Lyman series defines its lowest energy level n_1 as being equal to 1, the Paschen series defines n_1 as 3.

vibrational, or rotational transitions of individual atoms or molecules, and not from any collective behavior. But since these transitions are well defined energetically, the spectrum exhibits sharp lines.

In a liquid or solid, the atoms or molecules are sufficiently close together to affect each other's energy levels. That is, in addition to the individual energy levels of each atom or molecule, there exist energy levels for the collective excitation of groups of atoms or molecules. In general, the interaction of one atom or molecule with another decreases with internuclear distance or intermolecular distance in a continuous fashion, so that a continuous spectrum is observed.

CHAPTER 19


NUCLEAR CHEMISTRY

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 671 to 685 for step-by-step solutions to problems.

Nuclear chemistry is the chemistry of reactions that involve the nuclei of atoms rather than the electrons in the outer orbitals. They are, in general, much more energetic than ordinary (non-nuclear) reactions and often result in conversion of appreciable mass to energy according to Einstein's well known equation

$$E = mc^2 \qquad 19-1$$

The key to solving problems involving nuclear reactions lies in understanding that, instead of balancing equations to obtain the same number of elements on both sides of the equation, you must balance the sum of the atomic numbers and the sum of atomic weights, usually in amu (atomic mass units), on each side of the equation. Elements are, in general, not conserved. An illustration involving a typical nuclear reaction follows.



The subscripted leading number is the atomic number and is redundant since the chemical symbol specifies the atomic number. Nitrogen **always** has an atomic number of 7; H **always** has an atomic number of 1; etc. However, it is convenient to carry the atomic number specifically in nuclear reactions since they must balance. The superscripted number preceding the chemical symbol is the atomic weight in amu and differs for different isotopes of the same element. The atomic weight may be thought of as the sum of neutrons and protons in the nucleus. Note that the precise atomic weight will, in

general, differ from the approximate integral value used to balance equations. This difference is, in fact, a reflection of mass converted to energy in nuclear reactions.

In Equation 19-2, the sum of atomic numbers (9) is equal on both sides of the equation as is the sum of atomic weights (18). This is a requirement for balancing nuclear reactions.

There are several subatomic species that commonly enter into nuclear reactions. The most common ones are defined below with their corresponding atomic number and approximate (integral) atomic weight. These atomic numbers and atomic weights of subatomic species can be used to balance nuclear reactions in the manner identical to that used with atomic nuclei.

alpha particle (${}^4\text{He}$ nucleus)	${}^4_2\alpha$	(also ${}^4_2\text{He}$)
beta particle	${}^0_{-1}\beta$	(an electron)
positron	${}^0_1\beta$	
gamma ray	${}^0_0\gamma$	(a gamma ray has no mass, only energy)
proton (H^1 nucleus)	1_1p	(also ${}^1_1\text{H}$)
neutron	1_0n	

For balancing reactions, it is often convenient, though not precisely correct, to consider a neutron as a proton plus an electron and a proton as a neutron plus a positron.

Many elements have isotopes that are unstable and spontaneously decompose. Such isotopes are called radioactive isotopes or radioisotopes because their spontaneous decomposition is typically accompanied by the emission of high energy radiation. Radioactive decay is a first-order reaction. As discussed in Chapter 13, the half-life in a first-order reaction is independent of the amount of reactant initially present, so the rate constant for radioactive decay is often expressed as a half-life for a particular species. Equation 13-10 gives the rate constant in terms of the half-life for first-order reactions. It applies for radioactive decay reactions.

$$k = \ln(2)/t_{1/2} \text{ (for first-order reactions)} \quad 13-10$$

$$A/[A_0] = e^{(-kt)} = e^{(-t(\ln 2)/t_{1/2})} \quad 19-3$$

The energy associated with nuclear reactions is often sufficiently large that the accompanying change in mass is significant. When precise weights for each species are known, the energy of a nuclear reaction can be calculated as follows.

$$\Delta E = (\Delta m)c^2 \quad 19-4$$

where

$$\Delta m = \sum (m)_{\text{products}} - \sum (m)_{\text{reactants}} \quad 19-5$$

and is the mass change associated with the reaction.

$$A = p_1 M_1 + p_2 M_2$$

$$35.453 \text{ amu} = p_1 \times 35 \text{ amu} + p_2 \times 37 \text{ amu}$$

where $1 = p_1 + p_2$.

One must determine p_1 and p_2 (the probabilities of occurrence of isotope 35 and 37, respectively).

From the second of these relationships, $1 = p_1 + p_2$ one obtains $p_2 = 1 - p_1$. Then

$$A = p_1 M_1 + p_2 M_2 = p_1 M_1 + (1 - p_1) M_2$$

$$= p_1 M_1 + M_2 - p_1 M_2$$

$$= M_2 + (M_1 - M_2) p_1$$

$$\text{or, } p_1 = \frac{A - M_2}{M_1 - M_2} = \frac{35.453 \text{ amu} - 37 \text{ amu}}{35 \text{ amu} - 37 \text{ amu}} = 0.7735.$$

$$\text{Then, } p_2 = 1 - p_1 = 1 - 0.7735 = 0.2265.$$

Thus, the isotope of mass 35 amu occurs with probability of 0.7735 or $0.7735 \times 100 \% = 77.35 \%$ and the isotope of mass 37 amu occurs with a probability of 0.2265 or $0.2265 \times 100 \% = 22.65 \%$.

In reality, the two isotopes do not have integral atomic masses, and the percent occurrences calculated above are not exactly correct.

• PROBLEM 710

Chromium exists in four isotopic forms. The atomic masses and percent occurrences of these isotopes are listed in the following table:

<u>Isotopic mass (amu)</u>	<u>Percent occurrence</u>
50	4.31%
52	83.76%
53	9.55%
54	2.38%

Calculate the average atomic mass of chromium.

Solution: We will make use of the definition of average:

$$A = p_1 M_1 + p_2 M_2 + \dots + p_N M_N$$

where A is the average value, M_i is the atomic mass of isotope "i" and p_i is the corresponding probability of occurrence. For the four isotopes of chromium, we have:

$$M_1 = 50 \text{ amu} \quad p_1 = 4.31\% = 0.0431$$

$$M_2 = 52 \text{ amu} \quad p_2 = 83.76\% = 0.8376$$

$$M_3 = 53 \text{ amu} \quad p_3 = 9.55\% = 0.0955$$

$$M_4 = 54 \text{ amu} \quad p_4 = 2.38\% = 0.0238$$

Hence, the average atomic mass of chromium is

$$A = p_1 M_1 + p_2 M_2 + p_3 M_3 + p_4 M_4$$

$$= 0.0431 \times 50 \text{ amu} + 0.8376 \times 52 \text{ amu} + 0.0955 \times 53 \text{ amu} \\ + 0.0238 \times 54 \text{ amu}$$

$$= 2.155 \text{ amu} + 43.555 \text{ amu} + 5.062 \text{ amu} + 1.285 \text{ amu}$$

$$= 52.057 \text{ amu.}$$

• PROBLEM 711

Given that the masses of a proton, neutron, and electron are 1.00728, 1.00867, and .000549 amu, respectively, how much missing mass is there in $^{19}_9\text{F}$ (atomic weight = 18.9984)?

Solution: The total number of particles in $^{19}_9\text{F}$ and their total weight can be calculated. The amount of missing mass in $^{19}_9\text{F}$ will be the difference of this calculated weight and the given atomic weight of $^{19}_9\text{F}$. The subscript number 9, in $^{19}_9\text{F}$, indicates the atomic number of fluorine (F). Because the atomic number equals the number of protons, there are 9 protons in F. The superscript, 19, indicates the total number of particles in the nucleus. Since the nucleus is composed of protons and neutrons, and there are 9 protons, there are 10 neutrons present. In a neutral atom, the number of electrons equals the number of protons. Thus, there are 9 electrons. The total number of particles is, thus, 28. The mass and quantity of each particle is now known. Calculating the total weight contribution of each type of particle:

$$\text{Protons: } 9 \times 1.00728 = 9.06552$$


$$\text{Neutrons: } 10 \times 1.00867 = 10.0867$$

$$\text{Electrons: } 9 \times .000549 = \underline{\underline{.004941}}$$

$$\text{Total mass} = 19.1572.$$


It is given that the mass of the fluorine atom is 18.9984. Therefore, the missing mass is $19.1572 - 18.9984 = 0.1588 \text{ amu}$.

Calculate ΔE for the proposed basis of an absolutely clean source of nuclear energy



Atomic masses: ${}^1_1\text{B} = 11.00931$, ${}^4_2\text{He} = 4.00260$, ${}^1_1\text{H} = 1.00783$.

Solution: The energy of nuclear reactions, ΔE , is calculated from the difference between the masses of products and reactants in accordance with the Einstein Law. Einstein's Law can be stated $\Delta E = \Delta mc^2$, where Δm is the difference in the masses of the products and the reactants, and c is the speed of light (3×10^{10} cm/sec). The total reaction here can be written



Thus, Δm is equal to the mass of ${}^1_1\text{B}$ and ${}^1_1\text{H}$ subtracted from the mass of $3 {}^4_2\text{He}$. The mass of $3 {}^4_2\text{He}$ is equal to 3 times the mass of ${}^4_2\text{He}$.

$$\begin{aligned}\Delta m &= (3 \times m \text{ of } {}^4_2\text{He}) - (m \text{ of } {}^1_1\text{B} + m \text{ of } {}^1_1\text{H}) \\ &= (3 \times 4.00260) - (11.00931 + 1.00783) \\ &= 12.0078 - 12.01714 = -9.34 \times 10^{-3} \text{ g/mole.}\end{aligned}$$

One can now solve for ΔE by using Δm .

$$\begin{aligned}\Delta E &= \Delta mc^2 = -9.34 \times 10^{-3} \text{ g/mole} \times (3.0 \times 10^{10} \text{ cm/sec})^2 \\ &= -9.34 \times 10^{-3} \text{ g/mole} \times 9.0 \times 10^{20} \text{ cm}^2/\text{sec}^2 \\ &= -8.406 \times 10^{18} \text{ g cm}^2/\text{mole sec}^2 \\ &= -8.406 \times 10^{18} \text{ ergs/mole.}\end{aligned}$$

There are 4.18×10^{10} ergs in 1 Kcal, thus ergs can be converted to Kcal by dividing the number of ergs by the conversion factor 4.18×10^{10} ergs/Kcal.

$$\begin{aligned}\text{no. of Kcal} &= \frac{-8.406 \times 10^{18} \text{ ergs/mole}}{4.18 \times 10^{10} \text{ ergs/Kcal}} \\ &= -2.01 \times 10^8 \text{ Kcal/mole.}\end{aligned}$$

ΔE for this reaction is -2.01×10^8 Kcal per mole.

Calculate Δm , the difference in mass between the final and

and mass are related in the following equation,

$$\Delta E = \Delta mc^2$$

where, ΔE is the change in energy, Δm the change in mass and c the speed of light (3.0×10^{10} cm/sec). Δm is found by subtracting the mass of ${}^{238}_{90}\text{Th}$ from the sum of the masses of ${}^{234}_{88}\text{Th}$ and ${}^4_2\text{He}$.

$$\begin{aligned}\Delta m &= (234.0437 \text{ amu} + 4.0026 \text{ amu}) - 238.0508 \text{ amu} \\ &= - .0045 \text{ amu.}\end{aligned}$$

Energy is expressed in ergs ($\text{g}\cdot\text{cm}^2/\text{sec}^2$), therefore, Δm must be converted to grams before solving for ΔE .

$$\begin{aligned}\Delta m &= (- .0045 \text{ amu}) \frac{(1.673 \times 10^{-24} \text{ g})}{(1.0073 \text{ amu})} \\ &= - 7.47 \times 10^{-27} \text{ g.}\end{aligned}$$

Solving for ΔE :

$$\begin{aligned}\Delta E &= (- 7.47 \times 10^{-27} \text{ g}) (3.0 \times 10^{10} \text{ cm/sec})^2 \\ &= - 6.72 \times 10^{-6} \text{ g cm}^2/\text{sec}^2 = - 6.72 \times 10^{-6} \text{ erg}\end{aligned}$$

Therefore, 6.72×10^{-6} ergs are released.

• PROBLEM 715

Complete the following nuclear equations.

- (a) ${}_{7}\text{N}^{14} + {}_2\text{He}^4 \rightarrow {}_8\text{O}^{17} + \dots$
- (b) ${}_4\text{Be}^9 + {}_2\text{He}^4 \rightarrow {}_6\text{C}^{12} + \dots$
- (c) ${}_{15}\text{P}^{30} + {}_{14}\text{Si}^{30} + \dots$
- (d) ${}_1\text{H}^3 \rightarrow {}_2\text{He}^3 + \dots$

Solution: The rules for balancing nuclear equations are: (1) the superscript assigned to each particle is equal to its mass number and the subscript is equal to its atomic number or nuclear charge; (2) a free proton is the nucleus of a hydrogen atom, and is therefore written as ${}_1\text{H}^1$; (3) a free neutron has no charge and is therefore assigned zero atomic number. Its mass number is one and its notation is ${}_0\text{n}^1$; (4) an electron, β^- , has zero mass and its atomic number is -1 , hence the notation ${}_{-1}\text{e}^0$; (5) a positron has zero mass and its atomic number is $+1$, hence the notation ${}_{+1}\text{e}^0$; (6) an alpha particle (α -particle) is a helium nucleus, and is represented by ${}_2\text{He}^4$ or α ; (7) Gamma radiation (γ) is a form of light, and has no mass and no charge; (8) in a balanced equation, the sum of the subscripts must be the same on both sides of the equation; the sum of the superscripts must also be the same on both sides of the equation.

Solution: The half-life is defined as the time it takes for $\frac{1}{2}$ of the amount of a certain compound present to decompose. For example, if a substance has a half-life of 1 day, after one day there will only be $\frac{1}{2}$ of the original amount left. When given the original and final amount of a substance, after a given time has elapsed, the number of half-lives that have passed can be found. This is done by dividing the original amount by the final amount and determining how many factors of 2 are present in the quotient. The half-life is then found by dividing the time elapsed by the number of half-lives.

Solving for the half-life of ^{33}P :

$$\frac{\text{original amount}}{\text{final amount}} = \frac{2.000 \text{ pg}}{0.250 \text{ pg}} = 8$$

$8 = 2 \times 2 \times 2$, therefore 3 half-lives have passed. One is given that these half-lives elapse in 75.9 days.

$$\text{half-life} = \frac{75.9 \text{ days}}{3} = 25.3 \text{ days.}$$

• PROBLEM 720

The radioactive decay constant for radium is 1.36×10^{-11} . How many disintegrations per second occur in 100 g of radium?

Solution: The number of disintegrations per second of a given amount of a particular element can be determined by using the following equation.

$$D = \lambda N$$

where D is the number of disintegrations per second, λ the decay constant and N the number of atoms present.

In this problem one is given λ and must determine N before solving for D . One is told that 100 g of radium is present, the number of moles present is determined by dividing 100 g by the molecular weight of radium, 226 g/mole.

$$\text{no. of moles} = \frac{100 \text{ g}}{226 \text{ g/mole}} = 0.442 \text{ moles}$$

There are 6.02×10^{23} particles per mole, thus, one can calculate the number of atoms (N) in 0.442 moles.

$$\begin{aligned} N &= (0.442 \text{ moles})(6.02 \times 10^{23} \text{ atoms/mole}) \\ &= 2.66 \times 10^{23} \text{ atoms} \end{aligned}$$

Solving for D :

$$D = \lambda N = (1.36 \times 10^{-11})(2.66 \times 10^{23}) = 3.62 \times 10^{12} \text{ dis/sec.}$$

$$t = \frac{.327 \times 5730 \text{ yrs}}{.301} = 6225.0 \text{ yrs.}$$

Therefore, the tree is 6,225 years old.

CHAPTER 20


ORGANIC CHEMISTRY I: NOMENCLATURE AND STRUCTURE

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 687 to 716 for step-by-step solutions to problems.

Organic chemistry is the chemistry of substances that contain carbon. The number of catalogued organic compounds is in the millions and is expanding rapidly. In this chapter, the rules for the structure of some simple organic compounds and the associated rules for naming them are outlined.

Hydrocarbons contain only carbon and hydrogen, and are the simplest organic compounds. Their classical names form the basis of naming other organic compounds. Those compounds for which all possible carbon bonds contain hydrogen are called alkanes or saturated hydrocarbons. Unsaturated hydrocarbons have carbon-carbon double bonds, triple bonds, or aromatic (benzene-like) bonds.

The structure of a sample of normal or straight-chain alkanes is shown below. The chemical formula for the alkanes is $C_nH_{(2n+2)}$ and a sample of the names follow in Table 1.











Table 1: Names of Some Common Alkanes

n	Formula $[\text{C}_n\text{H}_{(2n+2)}]$	Name
1	CH_4	methane
2	C_2H_6	ethane
3	C_3H_8	propane
4	C_4H_{10}	butane
5	C_5H_{12}	pentane
8	C_8H_{18}	octane
10	$\text{C}_{10}\text{H}_{22}$	decane
12	$\text{C}_{12}\text{H}_{26}$	dodecane
16	$\text{C}_{16}\text{H}_{34}$	hexadecane
20	$\text{C}_{20}\text{H}_{42}$	eicosane

The structure of alkanes need not be in a straight chain. Branched chains can form numerous isomers with the same chemical formula, but with different structure and different physical and chemical properties. The C_{20} alkane, for example, has over 366,000 isomers and the C_{25} alkane has over 37,000,000. The straight-chain isomers are called normal hydrocarbons and their name is preceded by a small n as in n -butane, n -octane, etc.

The analogous series of hydrocarbons with one double bond (C_nH_{2n}) are called the alkenes; and with one triple bond ($\text{C}_n\text{H}_{(2n-2)}$) are called the alkynes.



Other, nonhydrocarbon, functional groups can also replace hydrogen on any of the hydrocarbons molecules discussed above. Table 2 shows the names used when other functional groups are substituted.

Table 2: IUPAC Names for Substituents on Organic Compounds

FAMILY	FUNCTIONAL GROUP	SUBSTITUENT NAME
Alcohols	—OH	add suffix <i>-ol</i> (e.g., 1-propanol)
Halogens	Fl, Cl, Br, I,	add prefix: <i>fluoro</i> , <i>chloro</i> , <i>bromo</i> , <i>iodo</i>
Ethers	C—O—C	add internal <i>-oxy</i> (e.g., methoxy, ethoxy)
Carboxylic acids	$ \begin{array}{c} \text{O} \\ \\ \text{C} - \text{OH} \end{array} $	add suffix <i>-oic acid</i> (e.g., ethanoic acid) also carboxy
Aldehydes	$ \begin{array}{c} \text{O} \\ \\ \text{C} - \text{H} \end{array} $	add suffix <i>-al</i> (e.g., pentanal)
Ketones	$ \begin{array}{c} \text{O} \\ \\ - \text{C} - \end{array} $	add suffix <i>-one</i> (e.g., pentanone)
Amides	$ \begin{array}{c} \text{O} \\ \\ - \text{C} - \text{NH}_2 \end{array} $	add suffix <i>-amide</i> (e.g., ethanamide)

In all cases the position of the substituent is numbered in the usual fashion.

It must be emphasized that there are many common names of organic compounds that were used before the IUPAC formal naming rules were established. The common names remain in use — particularly in commerce — resulting in many compounds being known by more than one name.

Step-by-Step Solutions to Problems in this Chapter, "Organic Chemistry I: Nomenclature and Structure"


ALKANES

• PROBLEM 724

Explain the terms primary, secondary, and tertiary in regards to covalent bonding in organic compounds.







Solution: If a carbon atom is bound to only one other carbon atom, then the former carbon atom is called primary. If a carbon atom is bonded to two other carbon atoms, then that carbon atom is called secondary. If a carbon atom is bonded to three other carbon atoms, then that carbon atom is called tertiary.

Any group that is attached to a primary, secondary, or tertiary carbon is called a primary, secondary, or tertiary group. For example,



Carbons 1, 2, and 5 are primary, carbon 4 is secondary, and carbon 3 is tertiary. The hydrogen atoms attached to carbons 1, 2, and 5 are called primary hydrogens, those attached to carbon 4 are called secondary hydrogens, and those attached to carbon 3 are called tertiary hydrogens. This same principle applies to alcohols, and depending upon where the hydroxyl group (- OH) is attached (that is, primary, secondary, or tertiary carbon), the alcohol is called primary, secondary, or tertiary, respectively.

Name each of the following alkanes. Indicate which, if any, are isomers.



Solution: Isomers are related compounds that have the same molecular formula but different structural formulas.

Isomerism is not possible among the alkanes until there are enough carbon atoms to permit more than one arrangement of the carbon chain.

To name the above compounds, one uses a set of rules to provide each compound with a clear name. These rules for nomenclature are the IUPAC rules (International Union of Pure and Applied Chemistry), and are referred to as systematic nomenclature.

One of the rules of the IUPAC system is to choose the largest chain of carbon atoms in the molecule and call it parent compound. Thus,




(a) has a 6 carbon parent chain with methyl groups bonded to the second and fourth carbon atoms of the parent chain. There exists, also, an ethyl group bonded to the third carbon atom.

As such, the name of this organic molecule is 2,4-dimethyl-3-ethylhexane.

(b) has 5 carbon parent molecule and 2 methyl groups on carbon number 3. Therefore, the name is 3,3-dimethylpentane.


(c) has a 5 carbon parent molecule with 2 methyl groups bonded to the third carbon of the parent chain. Therefore, the name of this structure becomes 3,3-dimethylpentane.

(d) has a 5 carbon parent molecule, and 2 methyl groups attached to carbon numbers 2 and 3.



Their names are (d) 1-chloro-2-methylbutane, (e) 1-chloro -3-methylbutane, (f) 2-chloro -2-methylbutane, and (g) 2-chloro -3-methylbutane.

Third, one writes the structure of the isomer with three carbons for a parent molecule, in which three hydrogen atoms of the three carbon chain have been replaced by a chlorine atom and two methyl groups:








This compound's name is 1-chloro -2,2-dimethylpropane or 1-chloro neopentane.

Hence, there are eight structural isomers with the molecular formula $C_5H_{11}Cl$.

A chemist has at his disposal the following hydrocarbons: (a) 3,3-dimethylpentane, (b) n-heptane, (c) 2-methylheptane, (d) n-pentane, and (e) 2-methylhexane. Arrange these compounds in order of decreasing boiling points (without referring to tables).

Solution: To rank the boiling points of these substances, you must consider the rules which govern the property of boiling points.

Except for very small alkanes, the boiling point rises 20 to 30 degrees for every carbon that is added to the chain. (Alkanes are saturated hydrocarbons of the general formula C_nH_{2n+2} , where n = number of carbon atoms.) Alkanes of the same carbon number but different structures (isomers) will have different boiling points. A branched-chain isomer has a lower boiling point than a straight-chain isomer. This is because the shape of a branched-chain molecule tends to approach a sphere, decreasing the surface area, so that the intermolecular forces are easily overcome at a lower temperature. With this in mind, you can proceed to solve the problem. Because these rules pertain only to structures, you must write them out.



c. has the greatest carbon content, 8 carbon atoms, which means it has the highest boiling point. a, b and e all have 7 carbon atoms; thus carbon content is equal. But only b is not branched, which means it has the highest boiling point out of this group. e is the next

You are given 3-bromohexane. Draw the stereoisomers of this compound; specify the R and the S enantiomers.

Solution: Non-superimposable mirror-image stereoisomers are called enantiomers. It is not enough that a molecule has a mirror image, for everything has a mirror image; the mirror images must not be able to be superimposed on each other. The arrangement of atoms that characterize a particular stereoisomer is called its configuration. You want to specify the configuration in terms of R and S. Let us see what this means.

In assigning configuration, you do the following after determining the chiral carbon: (1) Assign a sequence of priority to the four atoms or groups of atoms attached to the chiral center. (A chiral center consists of a carbon atom to which 4 different groups are attached.) (2) Visualize the molecule so that the group of lowest priority is directed away from you. Now, observe the arrangement of the remaining groups. If, in going from the group of highest priority to the group of second priority your eye travels in a clockwise direction, the configuration is R; if counterclockwise, the configuration is S. Now, how do you determine priority? If the four atoms bonded to chiral center are all different, priority depends on atomic number. The atom of higher atomic number receives the higher priority. If they are isotopes, the higher mass number has priority. Now, if you can't determine priority from this, because you have 2 or more atoms that are the same attached to the chiral center, make a similar comparison of the next atoms (i.e., their atomic numbers) as you move out from the chiral center. With this in mind, you can proceed as follows:

(a) 3-bromohexane. Recall, enantiomers are mirror image isomers so that you can say the enantiomers I and II, where the asterisk designates the chiral carbon.




Figure A

Solution: A polarimeter measures the rotation of plane polarized light, which is light whose vibrations take place in only one plane. An optically active substance has the ability to rotate the plane of polarized light. In essence, then, a polarimeter, measures optical activity. Species that possess optical activity have a specific rotation. It is defined in the following equation:

$$\text{specific rotation} = \frac{\text{observed rotation (degrees)}}{\text{length (dm)} \times \text{g/cc}}$$

The length of the polarimeter tube must be converted to decimeters. (1 cm = 0.1 dm) Thus, 5 cm = .5 dm. For water, 100 ml = 100 cc. Substituting, you obtain,

$$\text{specific rotation} = \frac{-1.2^\circ}{.5 \times \frac{6.15}{100}} = -39.02^\circ.$$

ALKENES AND ALKYNES

• PROBLEM 732

Classify each of the following as a member of the methane series, the ethylene series, or the acetylene series:
 $C_{12}H_{26}$, C_9H_{16} , C_7H_{14} , $C_{26}H_{54}$.

Solution: Before beginning this problem, one should first know the general formulas for each of the series. For the alkanes (methane series), the general formula is C_nH_{2n+2} , where n is the number of carbon atoms and $2n + 2$ is the number of hydrogens. Molecules of the ethylene series, also called the alkene series, have two adjacent carbon atoms joined to one another by a double bond. Any member in this series has the general formula C_nH_{2n} . The acetylene series, commonly called the alkyne series, has two adjacent carbon atoms joined to one another by a triple bond. The general formula for this series is C_nH_{2n-2} . With this in mind one can write:

$C_{12}H_{26}$:	C_nH_{2n+2}	:	alkane series
C_9H_{16}	:	C_nH_{2n-2}	:	acetylene series
C_7H_{14}	:	C_nH_{2n}	:	ethylene series
$C_{26}H_{54}$:	C_nH_{2n+2}	:	alkane series

Draw the structure of 4-ethyl-3,4-dimethyl-2-hexene.


Solution: To draw the structure of more complex compounds, such as this one, certain steps must be followed.

(1) Identify the parent compound that associated with the longest carbon chain that contains the functional group. In 4-ethyl-3,4-dimethyl-2-hexene, the parent compound is hexene.


(2) Draw the parent carbon skeleton, in this case, a six carbon chain. Do not put any hydrogen atoms in yet.




(3) Number the carbon atoms starting at either end. This is important; otherwise it may get confusing when one adds the functionality.




(4) Add the suffix functionality, in this case -2-ene. "Ene" tells one that a double bond is present, while "2" indicates that it is at the second carbon.



(5) Add the prefix functionality, starting at the beginning of the name and continuing until the parent name is reached. Here, the prefixes are 4-ethyl-3,4-dimethyl-.




Now, the hydrogen atoms can be added to give a complete structure.




Write condensed structural formulas for all the alkynes, i.e. unsaturated compounds with triple bonds, with a molecular formula of C_5H_8 .

Solution: A condensed structural formula provides all the information represented by other structural formulas (i.e. Lewis diagrams, bond diagrams), but it is not as cumbersome.

To solve this problem, first write all the structural isomers with five carbons in a continuous chain:




Second, write all the structural isomers with a four carbon parent molecule and one carbon in a branch:



No other structural formulas are possible.



Thus, there are only three structural formulas possible for C_5H_8 .

Next, one writes these formulas in the condensed form. Namely,






• PROBLEM 735

In the following pair of geometric isomers, designate which is cis and which is trans.



Solution: Isomers, i.e. compounds of the same molecular formula, that differ only in their geometry are termed geometric isomers. Geometric isomerism is one type of stereoisomerism; that is, isomerism due to the arrangement of atoms in a molecule in three dimensions. Geometric isomers are different compounds with different properties. Since the order of attachment of the atoms is the same, they are not structural isomers. The isomer that has the



Solution: All of these compounds are alcohols; they fit into the general formula R-OH , where R is any alkyl group.


In the systematic naming of any alcohol, the following rules should be followed:

(1) The longest chain that contains the hydroxyl group (OH) is considered the parent compound.

(2) The -e ending of the name of this carbon chain is replaced by -ol.


(3) The locations of the hydroxyl and any other groups are indicated by the smallest possible numbers.

Thus, compound (a)




has a 6-carbon chain and an hydroxyl group on carbon number 2. The name of this compound is 2-hexanol.

Compound (b) has a 5-carbon chain,



because this is the longest chain that contains the hydroxyl group. There is a methyl group on carbon number 3, and an hydroxyl group on carbon number 1. The name of this compound is, therefore, 3-methyl-1-pentanol.



Add the suffix functionality with the appropriate numbering: 4-hepten-2-ol. Add the prefix functionality, remembering to group together like prefixes. Then, double check to make sure a substituent has not been forgotten or one substituent has not been included twice. By following these steps, one arrives at the name for the structure:

4-n-butyl-3,5-diethyl-4-hepten-2-ol.

• PROBLEM 740

Which of the following, if any, are not alcohols derived from the methane series of hydrocarbons: $\text{C}_6\text{H}_5\text{OH}$, $\text{C}_{17}\text{H}_{33}\text{OH}$, $\text{C}_4\text{H}_8\text{OH}$, $\text{C}_9\text{H}_{19}\text{OH}$?


Solution: Alcohols are derived from molecules whose hydrogen atoms have been replaced by one or more hydroxyl ($-\text{OH}$) groups. The simplest alcohols are derived from the alkanes or the methane series and contain only one hydroxyl group per molecule. These have the general formula ROH , where R is an alkyl group of composition, $\text{C}_n\text{H}_{2n+1}$. Thus, the alcohols to be derived from the methane series will follow this formula.

(a) $\text{C}_6\text{H}_5\text{OH}$ has 6 carbons. Therefore, its hydrogen content should be $2n + 1 = 2(6) + 1 = 13$. Because there exist only 5 hydrogens, excluding the H from OH, it cannot be derived from the methane series.


Compound (b), $\text{C}_{17}\text{H}_{33}\text{OH}$, also does not fit the general formula ($n = 17$, $2n + 1 = 35$), and thus, is not derived from the methane series. If it did fit, it would possess 35 hydrogens instead of 33.

Compound (c), $\text{C}_4\text{H}_8\text{OH}$, does not fit the general formula ($n = 4$, $2n + 1 = 9$) and, thus, is not derived from the methane series. If it did fit, the alkyl group it would possess 9 H instead of 8.


Compound (d), $\text{C}_9\text{H}_{19}\text{OH}$, does fit the general formula ($n = 9$, $2n + 1 = 19$), and is derived from the methane series.



ethylbenzene



1,2 dimethylbenzene
(o-dimethylbenzene)



1,3 dimethylbenzene
(m-dimethylbenzene)



1,4 dimethylbenzene
(p-dimethylbenzene)

Solution: In establishing a consistent systematic scheme for the naming of benzene hydrocarbons several problems arise. The prefixes ortho, meta, and para (abbreviated o, m, and p) are commonly used to designate the 1,2-, 1,3-, and 1,4-, relationship of substituents on the benzene ring. The problem arises when, for example, the compound named systematically as 1,2-dimethylbenzene is also known as ortho-xylene or ortho-methyltoluene or ortho-dimethylbenzene.

Four isomeric compounds are possible with benzene hydrocarbons having the molecular formula C_8H_{10} . One is obtained by substituting an ethyl group for a hydrogen of benzene, whereas the other three are obtained by substituting methyl groups for two hydrogens.

● PROBLEM 744

Write structural formulas for the following: (2) 2-hexanone; (b) 2-methylbutanal; (c) O-methylbenzaldehyde; (d) methyl phenyl ketone.




Figure A





Figure B

Write a structural formula for each of the following molecular formulas. Name each structure that you draw. (a) CH_2O ; (b) $\text{C}_2\text{H}_6\text{O}_2$; (c) $\text{C}_2\text{H}_5\text{O}$; (d) $\text{C}_2\text{H}_4\text{O}_2$; (e) $\text{C}_2\text{H}_4\text{O}$.

Solution: In order to do this problem, one must know how carbon, hydrogen, and oxygen bond. Carbon can share its 4 valence electrons to form 4 covalent bonds, hydrogen can share its 1 valence electron to form 1 covalent bond, and oxygen can share its 6 covalent electrons to form only 2 covalent bonds. Thus, with this knowledge in mind, one can construct several types of molecular structures. A few of these structures have the following functional groups: Alcohols contain the hydroxyl group, $-\text{OH}$, acids contain the carboxyl group,




contain the carbonyl group, >C=O .

Structure (a), CH_2O , is an aldehyde,




and has the name formaldehyde.

Structure (b), $\text{C}_2\text{H}_6\text{O}_2$, is an alcohol




and has the name 1,2 dihydroxyethane or ethylene glycol.

Structure (c), $\text{C}_2\text{H}_5\text{O}$, is either an alcohol or an ether




and is named ethanol or dimethyl ether, respectively.

Structure (d) $\text{C}_2\text{H}_4\text{O}_2$, is an acid



and has the name acetic acid.

Structure (e), C_2H_4O , is an aldehyde




and has the name acetaldehyde.

• PROBLEM 746


Ethanol and methyl ether are isomers, having the molecular formula C_2H_6O . Based on structure, which one would be expected to have the higher vapor pressure? The higher boiling point? The greater solubility in water? Explain why in each case.

Solution: The ethers are a class of compounds that contain two hydrocarbon groups bound to an atom of oxygen. The general formula for ethers is ROR' where R and R' are alkyl groups and may or may not be the same

Both alcohols and ethers derived from alkanes have the same general formula $C_nH_{2n+2}O$ (ethanol is isomeric with methyl ether). Unlike alcohols, ethers do not form associated molecules by means of hydrogen bonds. Thus, the boiling point of an ether is considerably below the boiling point of its isomeric alcohol. Methyl ether is a gas, bp -23°C , whereas ethanol is a liquid, bp $+78^\circ\text{C}$. Therefore, it



ethanol



methyl ether

is reasonable to expect that methyl ether has a higher vapor pressure than ethanol.

Weak hydrogen bonding between water molecules and ether molecules leads to slight water solubility. The functional group that is characteristic of ethers, the C O C group, is chemically inactive compared to the hydroxyl group of alcohols. Thus, ethanol has the greater solubility.

• PROBLEM 747

Name and classify each of the following: $\text{CH}_3\text{COC}_2\text{H}_5$; $\text{C}_2\text{H}_5\text{OH}$; CH_3NH_2 ; $(\text{C}_2\text{H}_5)_2\text{O}$; $\text{CH}_3\text{COOC}_3\text{H}_7$; CH_3COOH .

Solution: All of these compounds are derivatives of hydrocarbons, i.e. compounds that contain only hydrogen and carbon atoms. The derivative of a hydrocarbon is the substitution of elements or radicals for one or more hydrogen atoms. These derivatives may be classified into functional groups, which, to a great extent, determine the properties of the compound. This is the key to naming these compounds, identify the functional groups and name it accordingly. First one must describe these functional groups.

Alcohols: They have the general formula $R-OH$, where R is any alkyl group.

Amines: They are derivatives of ammonia (NH_3), where one or more H 's have been replaced by an alkyl group. In other words, $R-NH_2$, $R-NH$, or $R-N-R$.

Ethers: General formula is $R-O-R'$, where R may be the same as R' .

Aldehydes: General formula is $R-\overset{\overset{O}{||}}{C}-H$

Ketones: General formula is $R-\overset{\overset{O}{||}}{C}-R'$, where R can equal R' . In both, ketones are aldehydes, $\text{C}=\text{O}$ is called the carbonyl group.

Carboxylic acids: General formula is $R-\overset{\overset{O}{||}}{C}-OH$, where $\text{C}-OH$ is called the carboxyl group.

Esters: General formula is $R-\overset{\overset{O}{||}}{C}-OR'$, where R can equal R' . With this in mind, proceed as follows:

$CH_3COC_4H_9$ can be written as $CH_3-\overset{\overset{O}{||}}{C}-C_4H_9$. This, then fits into the general formula of a ketone and is called methyl n-butyl ketone.

C_2H_5OH fits into the alcohol classification group. It is named ethanol or ethyl alcohol.

CH_3NH_2 fits the description of an amine and is named methylamine.

$(C_2H_5)_2O$ can be written as $H_5C_2-O-C_2H_5$, which fits the description of an ether. It is named ethyl ether.

$\text{CH}_3\text{COOC}_3\text{H}_7$ indicates that an ester is present, once it is rewritten as $\text{CH}_3-\overset{\text{O}}{\underset{\parallel}{\text{C}}}-\text{OC}_3\text{H}_7$. It is named as n-propyl acetate.


CH_3COOH can be written as $\text{CH}_3-\overset{\text{O}}{\underset{\parallel}{\text{C}}}-\text{OH}$. The carboxyl group is present, which tells you that it is an organic acid. The specific name is acetic acid.

• PROBLEM 748


Write the structure for methyl-6-chloro-5,8-diethyl-8-iodo-3,4,4-trimethyl-2,6-decadienoate.

Solution: To draw this structure one should follow certain rules.

(1) Identify and draw the parent name of the compound. The parent is decane,




(2) Add the suffix functionality. Note that in this compound two different types of functional groups are designated in the suffix endings, double bonds and an ester. Start with the double bonds: -2,6-decadienoate.




Now consider the ester group $\left(\overset{\text{O}}{\parallel} \text{C} - \text{O} - \right)$. The carbonyl ($\text{C} = \text{O}$) of the ester must be at carbon number 1.

The alkyl portion of the ester precedes the name (and the prefixes) as a separate word. In this case it is methyl (CH_3) so that the structure becomes



(3) Add the prefix functionality in the order given, 6-chloro-5,8-diethyl-8-iodo-3,4,4-trimethyl-.



In (d), 2-hydroxymethylcyclohexanol, the compound has no optical isomer, since two of the four groups attached to the central carbon atom are the same.


• PROBLEM 750

Which of the following organic compounds would you predict to be associated liquids: (a) CH_3OH ; (b) CH_3OCH_3 ; (c) CH_3Cl ; (d) CH_3NH_2 .

Solution: To answer this question, one should first understand the term, associated liquids. Liquids whose molecules are held together by hydrogen bonding are associated liquids. Hydrogen bonding is an especially strong kind of attraction between a hydrogen atom (that is bound to a highly electronegative atom) and another highly electronegative atom. This type of bond is formed by purely electrostatic forces between the positive end of one polar molecule and the negative end of another polar molecule. For hydrogen bonding to be important, the electronegative atoms must be F, O, or N.


Thus, once the structures of the species are known, one can tell if they are associated liquids.

(a) CH_3OH , (methanol);




It is an associated liquid because oxygen is bonded to the hydrogen atom. And the hydrogen atom thus forms a bond with another oxygen molecule of methane.

Liquid (b), CH_3OCH_3 , (dimethyl ether);



is not an associated liquid; the oxygen is not bonded directly to any hydrogen atom and therefore cannot form any hydrogen bonds.

Liquid (c), CH_3Cl , (methyl chloride);



Step-by-Step Solutions to Problems in this Chapter, "Organic Chemistry II: Reactions"

ALKANES

• PROBLEM 751

4 liters of octane gasoline weigh 3.19 kg. Calculate what volume of air is required for its complete combustion at S.T.P.

Solution: To answer this problem, you need to write the balanced equation for the combustion of octane gasoline. This means knowing what the molecular formula of octane gasoline is and what is meant by combustion. Octane is a saturated hydrocarbon, i.e., it is an alkane. Now, by saturated hydrocarbon, you mean a compound that contains only single bonds between the carbon to carbon and carbon to hydrogen bonds. Alkanes have the general formula C_NH_{2N+2} , where N = number of carbon atoms. Since the prefix "oct" means eight, you know there are 8 carbon atoms, which indicates that 18 hydrogen atoms are present. Thus, gasoline octane has the formula C_8H_{18} . Now, by combustion you mean the reaction of an organic compound with oxygen to produce CO_2 and H_2O . With this in mind, you can write the balanced equation for the reaction as



To find out what volume of air is required for combustion, you need the volume of O_2 required, since 20% of air is oxygen (O_2). To find the amount of O_2 involved, use the fact that at S.T.P. (standard temperature and pressure) 1 mole of any gas occupies 22.4 liters. Thus, if you knew how many moles of O_2 were required, you would know its volume. You can find the number of moles by using stoichiometry. You are told that there exists 3.19 kg or 3190 g (1000 g = 1 kg). The molecular weight (MW) of octane is 114 grams/mole. Thus, since

mole = $\frac{\text{grams (weight)}}{\text{M.W.}}$, you have $\frac{3190}{114} = 27.98$ moles of

gasoline. From the equation's coefficients, you see that for every 2 moles of gasoline, 25 moles of O_2 are required. Thus, for this number of moles of gasoline, you need $(27.98) \frac{25}{2} = 349.78$ moles of O_2 .

Recalling that 1 mole of gas occupies 22.4 liters at S.T.P., 349.78 moles of O_2 occupies $(349.78)(22.4) = 7835.08$ liters. Oxygen is 20% of the air. Thus, the amount of air required is


$$\frac{5 \text{ liters air}}{1 \text{ liter } O_2} \times 7835.08 \text{ liters } O_2 = 39,175.40 \text{ liters air.}$$

• PROBLEM 752

What is your understanding of a substitution reaction mechanism? Illustrate your answer with a specific example.

Solution: A substitution reaction is one in which one atom or group of atoms replaces another. In this type of reaction, the overall change taking place in the alkane is that one sigma bond is broken and a new sigma bond is formed. At elevated temperatures, or in the presence of ultraviolet light, one or more hydrogen atoms in an alkane molecule may be replaced by atoms of chlorine, bromine, or other similar atoms.

The substitution of a chlorine atom for hydrogen in alkane molecules proceeds by a free-radical mechanism and is a chain reaction. In the initiation step of chlorination of methane, for example, high temperature or light furnishes energy for dissociating chlorine molecules into chlorine radicals as shown (unpaired electrons are shown with a dot):



The chlorine atom collides with a methane molecule to form a methyl radical and a molecule of hydrogen chloride.



Subsequent collision of a methyl radical with a chlorine molecule produces the desired compound, CH_3Cl , and another chlorine atom



so that the whole chain may be repeated.

Equations 2 and 3 are called propagation steps in the chain reaction. Termination steps of this chain re-

action are 4 and 5:




These chain-terminating reactions tend to stop the reaction between methane and chlorine by consuming radicals necessary for the propagation steps.

• PROBLEM 753

A chemist has tert-butane. If he adds Br_2 and light, what are the monobromo substitution products expected? Which one will predominate and why? How would you prepare the predominating compound by an addition reaction?


Solution: The first question concerns substitution in alkanes. You are performing a halogenation substitution reaction and are asked to predict the monohalogenation products expected for tert-butane. This necessitates the writing of its structure and analyzing its hydrogens, since they will be replaced by the bromine atoms. Tert butane can be written




All the hydrogens on the methyl groups are equivalent. This means that if you substitute a bromine atom for one hydrogen atom from a methyl group, it is the same as substituting for any other hydrogen on one of the other methyl groups. The other hydrogen (the one attached to the 3° carbon) is not equivalent to the hydrogens bound to the 1° carbons. Thus you expect two monobromo products. They are




could have placed the Br on the other two methyls and the same molecule would be obtained, for Example,




Solution: You are told about isomerism in the compounds A and B, and their chlorinated products. When two compounds are isomers they have the same molecular formula, but differ in the sequence of linkages of atoms. These are structural isomers. Ethyl chloride has the structural formula




The equation for chlorination is its reaction with Cl_2 in the presence of light. Note that there exist two possible carbons on which the chlorines can bond. These are the two isomers A and B. Thus you can write




The question is which one is A, as the other must be B. You are told that A gives 2 isomers, while B yields 1, if they are chlorinated again. Let's start with






If you chlorinate it, you would obtain



These are two different compounds in the arrangement of atoms. They are structural isomers. Thus, isomer A must be




If this is correct, then the chlorination of B should yield only one product. Let's see if this is true.




• PROBLEM 755

The ease of formation of carbonium ions is $3^\circ > 2^\circ > 1^\circ > \text{CH}_3^+$. Explain why.


Solution: A carbonium ion is a group of atoms containing a carbon atom with only six electrons around it. Thus the carbonium ion is positively charged. The carbonium ions are classified as primary (1°), secondary (2°), and tertiary (3°) after the carbon bearing the positive charge. For example,



primary




secondary



tertiary

where R can be any alkyl group.

To understand why the ease of formation is $3^\circ > 2^\circ > 1^\circ > \text{CH}_3^+$, you need to understand some of the properties of a charged system. According to the laws of electrostatics, the stability of a charged system is increased by the distribution of the charge. Alkyl groups, for example, tend to donate electrons, so that any positive charge is dispersed. Thus, the more alkyl groups donating electrons, the greater the distribution of positive charges in a carbonium ion, and, as such, the more stable it is. The greater the stability of a substance, the easier it is to form. In our example, the 3° carbonium ion has 3 alkyl groups, the 2° ion has 2 alkyl groups, and the 1° ion has one alkyl group. Note CH_3^+ can be written



Therefore, $3^\circ > 2^\circ > 1^\circ > \text{CH}_3^+$ in ease of formation, since this sequence is also the sequence for the number of alkyls that distribute the positive charge, which results in increased stability.

ALKENES AND ALKYNES

• PROBLEM 756


What is your understanding of an addition reaction mechanism? Illustrate your answer with a specific example.

Solution: An important difference between saturated and unsaturated hydrocarbons is the general type of reaction that each will undergo. Alkanes react by substitution, while alkenes react by addition reactions (as well as substitution). In an addition reaction, the pi bond of the double bond is destroyed and two sigma bonds are formed, one to each of two carbon atoms.

A mixture consisting of only hydrogen and an alkene undergoes no detectable addition reaction even at high temperature and pressure. However, if a small amount of finely divided metal, such as platinum, palladium, or nickel is added as a catalyst, addition of hydrogen to the alkene occurs quite readily even at room temperature. (The same is true for alkynes, i.e. unsaturated compounds containing a triple bond.)

According to the commonly accepted view, alkenes and alkynes, with relatively loosely held electrons in their pi bonds, act as electron-pair donors, that is, Lewis bases, and protons from addition reagents, HX (where X is H, Cl, Br, ...), act as Lewis acids, that is, electron-pair acceptors. The first step, then, in the addition of HX to an alkene is the addition of a proton in a Lewis acid-base reaction. The process involves the formation of alkyl cations and chlorine anions as reaction intermediates. Such alkyl cations are called carbonium ions; they are groups of atoms that contain a carbon atom having only six electrons.

An example of an addition reaction would be





• PROBLEM 75

Write the structural formulas of the hydrogenation products of the following olefins: (a) propylene; (b) 2-methyl-2-pentene; (c) 1-methyl cyclohexene.


Solution: The most important reactions of alkenes (olefins) involve addition of reagents to the double bonds. An alkene can be converted to an alkane by addition of hydrogen to the double bond. These reactions are usually carried out by using a high pressure of hydrogen gas in the presence of a catalyst such as finely divided platinum, palladium, or nickel.

To solve this problem, draw the structural formulas of each alkene. The addition of hydrogen breaks the double bond, and the H_2 molecule attaches itself to the carbon atoms involved. Thus,




What are the products when the following hydrocarbons are oxidized with acidic permanganate solutions? (a) 2-butene; (b) 3,4-dimethyl-3-hexene; (c) 2-methyl-2-butene.


Solution: Alkenes readily react with a number of oxidizing agents. A test for the presence of an olefin is the reaction with an acidic solution of permanganate ion (MnO_4^-). The purple color of permanganate ion disappears as the olefin is oxidized. The course of such a reaction is that the olefin cleaves into two oxidized fragments. A carbon atom with two alkyl groups attached is converted to the carbonyl group of ketone, while a carbon atom with one attached hydrogen becomes the carboxyl group of an acid. This is summarized by the following




To solve this problem, write the structural formula of each hydrocarbon, find which alkyl groups correspond to R , R' , and R'' in the general formula, and then substitute into the general formula.



The product of this reaction is acetic acid.



The product of this reaction is methyl ethyl ketone.






The products of this reaction are acetic acid and acetone.


ALCOHOLS

• PROBLEM 763


Perform the following conversions (synthesis) using any inorganic reagents you require:




Solution: In any synthesis problem you can use a step by step approach. Try to find reactions that lead to intermediates from which the product can be obtained. In (A) you want to shift the hydroxyl group from the 1° carbon to the 2° carbon. You can do this through the corresponding alkene. Once the alkene is obtained, the addition of acid and water will generate the desired product. To obtain the intermediate, add acid to the starting material. Because it is an alcohol, it will dehydrate to the alkene. You have, then,



In (B) you want to switch the Cl from the 2° carbon to the 1° carbon. Convert the starting material to an alkene and from this the product can be obtained. You have, therefore, the following process:




Potassium hydroxide has the ability to dehalogenate an alkyl halide to an alkene.



With peroxides, the hydrogen from the HCl adds to the carbon with the least number of carbons attached. You


are performing an addition reaction across a double bond. In (C), you are converting the OH group to a NH_2 group. You can convert the OH to Cl, first, since upon addition of NH_3 , the NH_2 group is obtained. To obtain the chlorine, dehydrate the alcohol with acid to form the alkene, then add HCl. You have the following sequence:



• PROBLEM 764

Synthesize $\text{CH}_2\text{BrCH}_2\text{Br}$ from $\text{CH}_3\text{CH}_2\text{OH}$.

Solution: You are given the alcohol, ethanol, and are asked to synthesize $\text{CH}_2\text{BrCH}_2\text{Br}$, 1,2 dibromoethane. The fact that it is a dibromo compound suggests that the bromine added across a double bond. Therefore, if you could convert the alcohol to an olefin, namely, ethylene, you could add Br_2 in the solvent CCl_4 , and via an addition reaction, obtain the desired product. To obtain ethylene, remember that the alcohol can be dehydrated to the corresponding olefin by the addition of acid. This well-documented mechanism is as follows:



When you add the acid, you protonate the alcohol. Water is expelled and you obtain a carbonium ion, CH_3CH_2^+ . The carbonium expels a proton (H^+), so that ethylene is obtained. Now that you have $\text{CH}_2=\text{CH}_2$, add Br_2 and CCl_4 , and you obtain $\text{CH}_2\text{BrCH}_2\text{Br}$, your desired product. The Br_2 is added across the double bond in an addition reaction.

• PROBLEM 765

Ethanol, $\text{C}_2\text{H}_5\text{OH}$, reacts with HBr, hydrogen bromide, to give ethyl bromide, $\text{C}_2\text{H}_5\text{Br}$, and water. Ethanol, however, does not react with NaBr. Explain. Suggest a mechanism for the reaction with HBr.

A chemist dissolves methyl alcohol in n-octane. He adds this to excess methylmagnesium iodide, which is dissolved in the high-boiling solvent, n-butyl ether. A gas is evolved, which is collected at STP. It occupies a volume of 1.04 cc. What is the gas and how many grams of methyl alcohol were present?

Solution: Grignard reagents, which have the general formula of RMgX (R is an alkyl or aryl group, Mg is magnesium, and X is a halogen atom), are extremely reactive. In the presence of any acid stronger than an alkane, the grignard reagent is displaced to form RH. This happens because the grignard reagent may be considered the magnesium salt of a weak acid RH. Thus, a stronger acid displaces RH from its salt. Alcohols are stronger acids than alkanes. Therefore, when methyl alcohol (CH_3OH) is added to methyl magnesium iodide (CH_3MgI), the following reaction occurs:



At S.T.P. (0°C and 1 atm), CH_4 exists as a gas. To find the amount of CH_3OH present, remember that at STP a mole of gas occupies 22.4 liters. The gas produced here occupies only 1.04 cc or 0.00104 liters (1000 cc = 1 liter). Let x = moles of methane gas produced. Thus, you can set up the proportion

$$\frac{22.4 \text{ liters}}{1 \text{ mole}} = \frac{.00104 \text{ liters}}{x \text{ mole}}$$

Solving, $x = 4.64 \times 10^{-5}$ moles. From the stoichiometry of the equation, 1 mole of CH_3OH produces 1 mole of CH_4 . Thus, you have 4.64×10^{-5} moles of CH_3OH . The molecular weight (MW) of CH_3OH is 32 g/mole. By definition, 1 mole equals the weight in grams divided by MW. Substituting and solving for weight in grams,


$$4.64 \times 10^{-5} \text{ moles} = \frac{\text{grams of } \text{CH}_3\text{OH}}{32 \text{ g/mole}};$$

$$\begin{aligned} \text{grams of } \text{CH}_3\text{OH present} &= 32 (4.64 \times 10^{-5}) \\ &= 1.48 \times 10^{-3} \text{ grams.} \end{aligned}$$


How would you distinguish between the following pairs:
(a) isobutane and isobutylene (b) sec-butyl alcohol and n-heptane.

Solution: To distinguish between two compounds, try to

select for a characterization test. Begin by writing the structures of the compounds involved.




You see that isobutylene is an alkene, and that isobutane is an alkane. Thus, you should be able to distinguish isobutylene from isobutane by an addition reaction. Addition reactions are characteristic of alkenes and other unsaturated compounds. When Br_2/CCl_4 , a reddish brown solution, is in contact with an alkene, the following reaction occurs:




When the alkene consumes the Br_2 , the red color disappears. Thus, alkenes can be detected by their ability to decolorize the red solution of Br_2/CCl_4 .


You can also use a cold, dilute, neutral permanganate solution, the Baeyer test, instead of Br_2/CCl_4 to detect the presence of an alkene. Permanganate solution is purple colored, upon reaction with an alkene, the purple disappears and is replaced by a brown manganese dioxide precipitate. The reaction can be written



Alkenes do give this reaction. Thus, you now have two simple tests to distinguish isobutane from isobutylene.




These two can be distinguished by using a chromic anhydride reagent. Secondary alcohols, such as sec-butyl alcohol, are oxidized by chromic anhydride, CrO_3 , in aqueous sulfuric acid. The clear orange solution turns opaque, blue-green when the alcohol is added. In other words,



N-heptane is an alkane, and, as such, does not turn the solution blue green. Thus, you can distinguish sec-butyl alcohol from n-heptane by adding chromic anhydride.

Explain why benzene is a planar molecule, while cyclohexane, C_6H_{12} , contains a puckered ring of six carbon atoms.

Solution: The best way to approach this problem is to consider the geometry of the carbon-carbon bonding. To do this, consider its hybridization. Benzene can be written as shown in figure A.



Benzene

Figure A

Notice that each carbon has 4 bonds, and is bonded to three other elements (2 other carbons and 1 hydrogen). This suggests sp^2 hybridization. sp^2 hybridization allows bonding to three other elements, and leaves a "free" p orbital. This p orbital can overlap with another p orbital to give a pi bond, which serves to form the double bond. With sp^2 hybridization, the molecule must assume a geometry where the orbitals are 120° apart. This is the only way that the repulsion which occurs when like charges are brought together can be avoided. See figure B.




Figure B

An angle of 120° creates a planar geometry. This is true because a planar molecule with bond angles of 120° is a configuration where the 3 atoms involved are the farthest apart. Consider cyclohexane. It can be written as shown in figure C.

Each carbon atom is bound to four other atoms (2 other carbons and 2 hydrogens). This suggests sp^3 hybridization. With sp^3 hybridization, a molecule must assume a geometry that permits $109^\circ 28'$ bond angles. The atoms of the bond are arranged in a tetrahedral formation, this allows

The attack on the benzene ring is generally accomplished by an electron-deficient species such as a positive ion, or the positive end of a polar molecule. The electron withdrawing nitro group makes it more difficult for an electrophilic species to add onto the benzene ring; vigorous conditions are required for further nitration of nitrobenzene. The nitro group will "direct" the substitution of additional nitronium ions in a meta orientation.

With this in mind, the structure of the isomer of trinitrobenzene is at once known. Each nitronium ion is directed to the meta position of the other ion. Thus, the structure is that of figure B




Figure B


and has the formula 1,3,5-trinitrobenzene.

• PROBLEM 774


Explain why the hydrolysis of an ester to yield alcohol and acid is best carried out in basic rather than acidic solution. Use equations to illustrate your answer.

Solution: The best way to solve this problem is to consider what products will be formed with acid and base.


The general formula of an ester is $R' - \overset{\overset{O}{\parallel}}{C} - OR$.



The carboxylic acid and alcohol are generated. But esters are generated from carboxylic acid and alcohols by the use of H^+ . In other words, there exists nothing to prevent the reaction from going back in the other direction. That is, you have the following equilibrium:



Let us consider the reaction with base.




If the temperature is raised above 160°C, a different set of reactions occur. Again, the acid protonates the

alcohol to form $\text{C}_2\text{H}_5\text{OH}^+$. Water departs to yield C_2H_5^+ .


Let us write this as $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\text{H}$. This time, however,

it does not react with another alcohol molecule but expels a proton immediately to form the alkene, ethylene or ethene.




This is the final product, ethylene, at a temperature greater than 160°C.

(2) Here you have citric acid reacting with a base, KOH. This suggests the formation of a salt and water. To predict which salt, write out the structure of citric acid. The structure is,



To understand how it reacts, you need to know the origin of its acidic nature. The acidic nature is determined by the H of the carboxylic group, COOH. Thus, you can anticipate the following reaction:




• PROBLEM 779

Indicate how you would prepare (1) acetone from acetic acid and (2) acetic acid from ethyl alcohol.

Solution: This problem is concerned with synthesis of organic compounds. In such problems you want to write out the structures of the initial and final products involved, so that you can identify functional groups. From this, you employ reactions that change the functional groups until the desired product is obtained.


(1) Acetone : $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$; Acetic Acid : $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$

If you had calcium acetate, $\left(\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}\right)_2\text{Ca}$, you could obtain acetone by heating it, since calcium acetate forms calcium carbonate and acetone upon the addition of heat. To obtain calcium acetate, add $\text{Ca}(\text{OH})_2$, which is a base. Because it is a base, the acid reacts with it to produce the calcium acetate, which is the salt, and water. Thus, the sequence of reactions becomes



(2) Acetic acid : $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$

Ethyl alcohol : $\text{CH}_3\text{CH}_2\text{OH}$. To prepare acetic acid from ethyl alcohol, you must know one important fact: Ethyl alcohol is a primary alcohol, i.e., one where the OH group is located on a carbon atom bonded to only one other carbon atom. Primary alcohols can be directly oxidized to the corresponding carboxylic acid by the addition of KMnO_4 , potassium permanganate. Thus, you perform the following reaction:



• PROBLEM 780

Explain how you would prepare benzoic acid from benzene. Use any inorganic reagents.




Figure A

Solution: One of the most common ways of preparing benzoic acid from benzene is through an intermediate grignard. To plan this synthesis, first write your starting material and desired product. Then, think of inorganic reagents that lead you step by step through intermediates until you obtain the product.

A grignard reagent can be represented as $R-MgX$, where R is any aryl or alkyl group and X is a halogen. To obtain this grignard intermediate, you must halogenate the benzene ring. This can be accomplished by using a solution of halogen and FeX_3 , where X is the halogen. Let the halogen be Br . Thus, you have the reaction in Figure B.




Figure B

This is electrophilic substitution of a hydrogen in the ring by a bromine atom. By electrophilic, you mean electron seeking. By addition of Mg , in dry ether, you obtain the grignard reagent. See figure C.




Figure C

Now that you have the grignard, add carbon dioxide. The reaction proceeds as shown in figure D.




Figure D

With the addition of the acid, HBr , the reaction in figure E proceeds to the desired product benzoic acid.




Figure E

Show the mechanism for the transesterification of methyl benzoate with ethanol in an acidic solution.

Solution: Begin this problem by defining transesterification. An ester is an organic compound

of general formula $R-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{OR}'$, where R and R' are alkyl groups that can be the same or different. In transesterification, the alcoholic portion of the ester is exchanged for another alcoholic group. You are asked to give the mechanism for the reaction shown in figure A.




Figure A

You will note that you exchange the methyl alcohol portion with ethyl alcohol. From radioactive isotopes, the following mechanism shown in figure B has been established.




Figure B

Due to the electronegativity of oxygen, the oxygen of the carbonyl group, >C=O , has a slightly negative charge and the carbon has a slightly positive charge. The proton from the acid is attracted to the oxygen as in figure C. These two are in equilibrium with each other.




Figure C


CHAPTER 22

BIOCHEMISTRY I

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 755 to 788 for step-by-step solutions to problems.

Biochemistry is that branch of chemistry which deals with the chemistry in living cells and organisms. Because living systems are organic, biochemistry is heavily dependent on organic chemistry. However, it is a very specialized part of organic chemistry, and the two should not be confused as being equivalent. Biochemistry, like organic chemistry, is a huge area with many very complicated reactions and mechanisms of importance to living systems. Only the simplest of the ideas can be introduced in this chapter.

Amino acids — molecules that contain both an amino (NH_2) group and a carboxylic acid (CO_2H) group — are the building blocks of proteins in living systems. These acids have two acid-base sites per molecule — one is the carboxylic acid hydrogen, the other the basic amino site. These sites can ionize independently or together according to the overall reaction.



where R group is an arbitrary organic group that will be different for different amino acids. The ion on the right side of Equation 22-1, with both a positive and negative charge, is called a zwitterion. There are two ionization constants for this ion — one for the ionization of the CO_2H group and one for the NH_3^+ group. The pK_1 and pK_2 constants, shown in Problem 792 for several amino acids, are the negative logarithms of the ionization constant for these reactions. The pH that exists when there is no net charge on the ion is called the isoelectric pH and is written as pI.

Step-by-Step Solutions to Problems in this Chapter, "Biochemistry I"

CELLULAR CONSTRUCTION AND DIMENSIONS

• PROBLEM 784

A chemist has an *E. coli* bacterium cell of cylindrical shape. It is 2μ (microns) long and 1μ in diameter and weighs 2×10^{-12} g. (a) How many lipid molecules are present, assuming their average molecular weight to be 700 and lipid content to be 2%? (b) If the cell contains 15,000 ribosomes, what percent of the volume do they occupy? You may assume them to be spherical with a diameter of 180 Å; also 10^4 Å = 1 micron.

Solution: (a) Because you are told the molecular weight of an average lipid molecule, you need only compute their total weight in the *E. coli* cell; and then find the number of moles present. A mole is defined as weight in grams/molecular weight (MW). From this, you use the fact that in one mole of any substance, 6.02×10^{23} molecules (Avogadro's number) exist.

The lipid content is given as 2%. If the total weight of the cell is 2×10^{-12} g, the lipid molecules must have a total weight of $(.02)(2 \times 10^{-12})$ or 4×10^{-14} grams. The average molecular weight of a lipid molecule is given as 700 g/mole. Thus, you have $4 \times 10^{-14}/700 = 5.71 \times 10^{-17}$ moles of lipid molecules. If 6.02×10^{23} molecules are in a mole, then in 5.71×10^{-17} moles, there are $(6.02 \times 10^{23})(5.71 \times 10^{-17}) = 3.44 \times 10^7$ lipid molecules.

(b) Calculate the volume of both the *E. coli* cell and the total volume occupied by the ribosomes.

Volume of the *E. coli* cell: It is of cylindrical shape. The volume of a cylinder can be found from the product of its length and the area of its base. The area of the base, a circle, is πr^2 , where r = radius. Because

the diameter = $2r$, the radius of the cylindrical E. coli cell is 1μ divided by 2 or 0.5μ . This is converted to $5 \times 10^3 \text{ \AA}$, by the conversion factor $10^4 \text{ \AA} = 1 \text{ micron}$. Similarly, $2\mu = 2 \times 10^4 \text{ \AA} = \text{length of E. coli cell}$. Its total volume = (area of circle) (length) = $[\pi(5 \times 10^3)^2 \times 2 \times 10^4] = 1.57 \times 10^{12}$.

Volume of ribosomes: Each ribosome has a diameter of 180 \AA or a radius of 90 \AA . It is given that it is a sphere. The volume of a sphere is given by the formula $\frac{4}{3} \pi r^3$, where $r = \text{radius}$. Substituting the volume of one ribosome = $\frac{4}{3} \pi (90)^3 = 3.05 \times 10^6$. If there exists 15,000 ribosomes, their total volume must be $(15,000)(3.05 \times 10^6) = 4.58 \times 10^{10}$. Therefore, the percentage volume it occupies =

$$\frac{\text{total volume of ribosomes}}{\text{volume of E. coli cell}} \times 100 = \frac{4.58 \times 10^{10}}{1.57 \times 10^{12}} \times 100 = 2.9\%$$

• PROBLEM 785

An E. coli culture will grow to a limiting concentration of about 10^9 cell per cm^3 . At this concentration what percentage of the total volume of the culture medium is occupied by the cells? Assume that an E. coli cell is cylindrical and that it is 2.0μ long and has a 1.0μ diameter.

Solution: This percentage can be found by first determining the volume that one E. coli cell occupies and then the volume that 10^9 cells occupy. This volume divided by 1 cm^3 multiplied by 100 equals the percent of the total volume that the cells occupy.

The volume of a cylinder is equal to its length multiplied by the area of its cross-section. Area of a circle = πr^2 , where $r = \text{radius of the circle}$. $1\mu = 10^{-4} \text{ cm}$

$$\begin{aligned} \text{Volume of 1 E. coli cell} &= (2.0 \times 10^{-4} \text{ cm}) (\pi) \\ &\quad \times (0.5 \times 10^{-4} \text{ cm})^2 \\ &= 1.57 \times 10^{-12} \text{ cm}^3 \end{aligned}$$

There are 10^9 cells per cm^3 of solution, therefore the volume that the cells occupy in 1 cm^3 of solution is $10^9 \times 1.57 \times 10^{-12} \text{ cm}^3$ or $1.57 \times 10^{-3} \text{ cm}^3$.

$$\begin{aligned} \text{Percent of total volume} \\ \text{that the E. coli cells} \\ \text{occupy} &= \frac{1.57 \times 10^{-3} \text{ cm}^3}{1.0 \text{ cm}^3} \times 100 = .157\% \end{aligned}$$

a) Calculate the ratio of the volume of a hepatocyte to that of an E. coli cell. Assume the hepatocyte to be a cube 20μ on an edge. b) Calculate the ratio of their surface areas. c) Calculate the surface/volume ratios for each.

Solution: A hepatocyte is an eucaryote, a large complex cell, and an E. coli cell is a procaryote, a small simple cell. An E. coli cell is cylindrical with a length of 2.0μ and a cross-sectional diameter of 1.0μ . The volume of a cylinder is equal to the product of its length and the area of its cross-section.

$$\text{Volume of an E. coli cell} = 2.0\mu \times \pi \times (0.5\mu)^2 = 1.57\mu^3$$

$$\text{Volume of a hepatocyte} = (20\mu)^3 = 8.0 \times 10^3\mu^3$$

The volume ratio is

$$\frac{\text{hepatocyte}}{\text{E. coli}} = \frac{8.0 \times 10^3\mu^3}{1.57\mu^3} = 5.1 \times 10^3 : 1.$$

b) The surface area of a solid cylinder is equal to $2\pi rl + 2\pi r^2$, where r is the cross-sectional radius and l is the length.

$$\begin{aligned} \text{Surface area of an E. coli} &= 2\pi(0.5\mu)(2.0\mu) + 2\pi(0.5\mu)^2 \\ &= 6.28\mu^2 + 1.57\mu^2 = 7.85\mu^2 \end{aligned}$$

The surface area of a cube is $6s^2$, where s is the length of the edge.

$$\text{Surface area of a hepatocyte} = 6(20\mu)^2 = 2.40 \times 10^3\mu^2$$

The ratio of their surface areas is

$$\frac{\text{hepatocyte}}{\text{E. coli}} = \frac{2.40 \times 10^3 \mu^2}{7.85 \mu^2} = 306 : 1.$$


c) Surface to volume ratios:

$$\text{E. coli; } \frac{\text{surface}}{\text{volume}} = \frac{7.85 \mu^2}{1.57 \mu^3} = 5 \mu^{-1}$$

$$\text{Hepatocyte; } \frac{\text{surface}}{\text{volume}} = \frac{2.40 \times 10^3 \mu^2}{8.0 \times 10^3 \mu^3} = 0.3 \mu^{-1}.$$

• PROBLEM 787

The hydrated K^+ and Cl^- ions are approximately spherical in shape, with a diameter of 6.0 Å. What fraction of the total volume of the water phase of an E. coli cell is occupied by these ions?



KCl dissolving
in water.

Solution: A hydrated ion is an ion that is completely surrounded by water molecules. Water gathers around ions because of the strong electrostatic attraction between the water dipoles and the ions. The O atom of a water molecule is slightly negative and is attracted to the positive K^+ ion and the H atom of the water is slightly positive and is attracted to the negative Cl^- ion.

The concentration of KCl in an E. coli cell is 150×10^{-3} M. Because KCl dissociates into K^+ and Cl^- ions, the ion concentration in the cell is $2(150 \times 10^{-3} \text{ M})$. If one knows the volume of the aqueous phase of an E. coli cell, one can determine the number of moles of the ions present. An E. coli is cylindrical; it is 2.0μ long and has a diameter of 1.0μ . The volume of a cylinder is equal to $\pi r^2 l$, where l is the length of the cylinder and r is the radius of its cross-section. $1 \mu = 10^{-6} \text{ cm}$.

The volume of a cylinder is equal to its length times the area of its cross-section.

$$l \times \pi r^2,$$

where l is the length of the cylinder and r is the radius of the cross-section. $1\mu = 10^{-4}$ cm.

$$\begin{aligned}\text{Volume} &= 1.8 \times 10^{-4} \text{ cm} \times \pi \times (0.5 \times 10^{-4} \text{ cm})^2 \\ &= 1.4 \times 10^{-12} \text{ cm}^3 = 1.4 \times 10^{-15} \text{ l}\end{aligned}$$

Solving for the number of moles of H^+ ion,

$$\begin{aligned}\text{No. of moles} &= 10^{-6} \text{ moles/liter} \times 1.4 \times 10^{-15} \text{ l} \\ &= 1.4 \times 10^{-21} \text{ moles}\end{aligned}$$

There are 6.02×10^{23} ions per mole.

$$\begin{aligned}\text{No. of } \text{H}^+ \text{ ions} &= 1.4 \times 10^{-21} \text{ moles} \times 6.02 \times 10^{23} \frac{\text{ions}}{\text{mole}} \\ &= 843 \text{ H}^+ \text{ ions.}\end{aligned}$$

• PROBLEM 791

At pH values of 4 to 9, natural amino acids exist as polar or zwitter ions: $\text{H}_2\text{N}^+\text{CHZCO}_2^-$. At a pH of 12, what would be the predominant ionic type? What would it be at a pH of 2? Can the uncharged molecular form, $\text{H}_2\text{NCHZCO}_2\text{H}$ ever predominate at any pH?

Solution: To answer these questions, you need to consider the chemical composition of amino acids.

Amino acids are compounds whose molecules possess both the amino ($-\text{NH}_2$) and the carboxy ($-\text{CO}_2\text{H}$) functional groups. An amino group is basic - if it is an electron donor. The carboxylic group is acidic; it releases protons (H^+). In a zwitter ion, as indicated, the carboxylic group has donated its proton to become $\text{COO}^- \equiv \text{CO}_2^-$ and NH_2 has received a proton to become NH_3^+ . The presence of such an ion is dependent on pH, which is a measure of the H^+ or hydronium ion (H_3O^+) concentration. Thus, pH gives an indication of the acidity or basicity of a solution. The lower the pH, the more acidic the solution, and, as such, the greater the concentration of H^+ ions. The higher the pH, the more basic the solution, and, the lower the concentration of H^+ ions. One can now proceed as follows:

pH = 12. At this pH, which is high, the H^+ concentration is low; the solution is basic. Thus, the amino group (NH_2) will probably not be protonated. The carboxylic


Paper electrophoresis at pH 6.0 was carried out on a mixture of glycine, alanine, glutamic acid, lysine, arginine and serine. (a) Which compound moved toward the anode? (b) Which moved toward the cathode? (c) Which remained at the origin?

Solution: A method of separating amino acids is paper electrophoresis. A drop of the solution containing the amino acids is placed on a filter paper sheet, which is then moistened with a buffer of a given pH. The ends of the sheet dip into electrode vessels, and a high-voltage electrical field is applied while cooling. Because of their different pK^1 values, the amino acids migrate in different directions and at different rates, depending on the pH of the system and the emf (electromotive force) applied. To determine whether an amino acid will move towards the anode (positive end) or the cathode (negative end), one must determine the charge on the particular amino acid at pH = 6. This is done by calculating the isoelectric pH_I (the pH, where the amino acid is electrically neutral) of each amino acid. $pH_I = \frac{1}{2}(pK^1 + pK^2)$. If pH_I is less than pH = 6, then the amino acid will be negatively charged at pH = 6 and will move towards the cathode. When the pH_I of an amino acid is near 6, the amino acid will not move.


glycine: $pH_I = \frac{1}{2}(2.34 + 9.6) = 5.97$. This is very near pH = 6, and glycine will remain in the center.

alanine: $pH_I = \frac{1}{2}(2.34 + 9.69) = 6.02$. This is also very close to 6 and alanine will not move from the center.

glutamic acid: As glutamic acid is titrated with base, the following species come about:



Species B is the neutral species, here. Therefore, $pH_I = \frac{1}{2}(pK^1 + pK^2_R) = \frac{1}{2}(2.19 + 4.25) = 3.22$. 3.22 is much less than 6 and therefore glutamic acid will be negatively charged at pH = 6 and will move towards the anode.



A cytochrome.
R' and R" are
different alkyl
groups.

Solution: From the accompanying figure, one can see that there is one iron atom in each cytochrome molecule. Thus, there is 1 mole of Fe atoms in 1 mole of cytochrome C. If each molecule is 0.45 percent iron by weight, then in 1 mole of cytochrome C, 0.45 percent of the total molecular weight of cytochrome C is equal to the weight of one mole of Fe. MW of Fe =

$$(4.5 \times 10^{-3}) (\text{MW of cytochrome C}) = 55.8 \text{ g/mole}$$

$$\text{MW of cytochrome C} = \frac{55.8 \text{ g/mole}}{4.5 \times 10^{-3}} = 12,400 \text{ g/mole.}$$

• PROBLEM 804

The hard shell of crustaceans (lobsters, etc.) and insects (roaches, etc.) is a polysaccharide called chitin. On enzymatic hydrolysis of chitin, N-acetylglucosamine is obtained. This molecule resembles glucose except that at

C-2 a $\text{-N}-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-\text{CH}_3$ is attached instead of -OH . (a) Write an open chain formula for N-acetylglucosamine.

(b) The structure of chitin is analogous to that of cellulose. Draw a formula containing two joined N-acetylglucosamine units. (c) If the molecular weight of chitin is 150,000, how many units are in the polymer?

Solution: A polysaccharide is a chain made up of many simple sugars. Chitin is a long unbranched chain of many molecules of the sugar, N-acetylglucosamine.

(a) One is given that the only difference between glucose and N-acetylglucosamine is that a $\text{-N}-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-\text{CH}_3$ group replaces the -OH on C-2. The open-chain formula of N-acetylglucosamine can therefore be written as shown in Figure A.




Figure D

Because the O in the - OH on C-1 joins to a C-4 atom on the adjacent molecule and the ring structures are called glycosides, this is termed a 1,4-glycosidic linkage.


(c) To determine the number of units in one strand of chitin, one can divide the total weight of the molecule by the weight of one glycoside. From the ring structure of a glycoside joined to another glycoside, one sees that there are 8C's, 5 O's, 13 H's and 1 N. The weight of one mole of this glycoside is $(8 \times 12) + (5 \times 16) + (13 \times 1) + (14) = 203$ g/mole.

$$\text{No. of units} = \frac{150,000 \text{ g}}{203 \text{ g/mole(unit)}} = 739 \text{ units.}$$

• PROBLEM 805

Calculate the density of the tightly coiled tropocollagen molecule, which may be considered to be a cylinder 2,800 Å long and 14 Å in diameter. It contains three polypeptide chains of 1,000 amino acid residues each.

Figure A



Conformation of polypeptide chains in triple-stranded tropocollagen molecule. Each chain is a coil with many repeating sequences of Gly-Pro-Hyp.

$$M_n = \frac{\sum_i n_i M_i}{\sum_i n_i} = \sum_i X_i M_i$$

Here, n_i is the number of molecules of molecular weight M_i per gram of dry polymer and X_i is the mole fraction of each component.

Solving for M_n :

$$\begin{aligned} M_n &= 0.5(100,000) + 0.5(200,000) \\ &= 50,000 + 100,000 = 150,000 \text{ g/mole.} \end{aligned}$$

(b) In the weight-average M_w , each item counts not as a single unit but in proportion to its weight. The molecular weight M_i is multiplied by the weight $n_i M_i$ of material of that molecular weight rather than by the number of molecules. The weight average molecular weight M_w is defined

$$M_w = \frac{\sum_i n_i M_i^2}{\sum_i n_i M_i}$$

$$\text{Because } \frac{\sum_i n_i M_i}{\sum_i n_i} = \sum_i X_i M_i; \quad M_w = \frac{\sum_i X_i M_i^2}{\sum_i X_i M_i}$$

Solving for M_w :

$$\begin{aligned} M_w &= \frac{0.5(100,000)^2 + 0.5(200,000)^2}{0.5(100,000) + 0.5(200,000)} \\ &= \frac{(1.0 \times 10^5)^2 + (2.0 \times 10^5)^2}{(1.0 \times 10^5) + (2.0 \times 10^5)} = \frac{5.0 \times 10^{10}}{3.0 \times 10^5} \\ &= 1.67 \times 10^5 \text{ g/mole.} \end{aligned}$$

• PROBLEM 809

For a condensation polymerization of a hydroxyacid with a residue weight of 200, it is found that 99% of the acid groups are used up. Calculate a) the number average molecular weight and b) the weight average molecular weights.

Solution: a) The number average molecular weight is the average weight of all particles present, where each unit

is given equal weight. Thus, the number average molecular weight is equal to the individual residue weight times the average number of residues. This number can be calculated from the fact that 99% of the acid groups have reacted to form the polymer chain. When monomers join to form a polymer, the last monomer is left unreacted at the end of the chain. If 99% of the acid reacts then 1% is left unreacted, so that 1% of the average number of monomers is equal to 1. Let the number average of the monomers equal \bar{X}_n . Then,

$$.01 \bar{X}_n = 1$$

$$\bar{X}_n = 100$$

One can now solve for the number-average molecular weight, M_n .

$$\begin{aligned} \text{number-average mol.wt} &= \text{mol.wt of a monomer} \times \bar{X}_n \\ &= 200 \text{ g/mole} \times 100 \\ &= 2.0 \times 10^4 \text{ g/mole.} \end{aligned}$$

b) The weight average molecular weight, M_w , weights molecules proportionally to their molecular weight in the averaging process. It is found from some complex derivations that

$$M_w = M_n(1 + p), \quad \text{where } p \text{ is the extent of the re-}$$

action. In this problem, p is .99. From this one can solve for M_w

$$\begin{aligned} M_w &= (2.0 \times 10^4 \text{ g/mole})(1 + 0.99) \\ &= 3.98 \times 10^4 \text{ g/mole.} \end{aligned}$$

• PROBLEM 810

For a condensation polymerization of a hydroxyacid in which 99% of the acid groups are used up, calculate (a) the average number of monomer units in the polymer molecules, (b) the probability that a given molecule will have the number of residues given by this value, and (c) the weight fraction having this particular number of monomer units.

Solution: (a) The average number of monomer units is equal to

$$\bar{X}_n = \frac{1}{1 - p},$$

PHYSICAL ASPECTS OF BIOCHEMISTRY

• PROBLEM 811

Calculate the percentage of the total wet weight of a hepatocyte that is contributed by the cell membrane. Assume the hepatocyte to be a cube 20μ on an edge and that the membrane is 90 \AA thick and that its specific gravity, as well as that of the entire cell, is 1.0.

Solution: Because the specific gravity is the same for the membrane and the rest of the cell, the weight percentage is equal to the volume percentage. If the hepatocyte is assumed to be a cube, its volume is equal to s^3 , where s is length of the side. The membrane is composed of six rectangular solids. Each solid is 90 \AA by $20\mu \times 20\mu$ and one is found on each face of the cube. The volume of the membrane is equal to the length times the product of the height and width times 6. The percentage of the total weight of the cell that is taken up by the membrane is the quotient of the volume of the membrane divided by the total volume of the cell multiplied by 100. Since $1 \text{ \AA} = 10^{-4} \mu$

$$\text{Volume of cell} = (20\mu)^3 = 8.0 \times 10^3 \mu^3$$

$$\begin{aligned} \text{Volume of the membrane} &= (9.0 \times 10^{-3}\mu)(20\mu)(20\mu) \times 6 \\ &= 21.6 \mu^3 \end{aligned}$$

Percent of weight taken up by the membrane =

$$= \frac{21.6 \mu^3}{8.0 \times 10^3 \mu^3} \times 100 = .27\%.$$

• PROBLEM 812

Calculate the molecular weight of a pure isoelectric protein if a 1% solution gives an osmotic pressure of 46 mm of H_2O at 0°C . Assume that it yields an ideal solution.

Solution: The relationship between molecular weight and osmotic pressure is

$$M = \frac{C}{\pi} RT,$$

where M is the molecular weight, c the concentration in grams per liter, R the gas constant (0.082 liter atm deg⁻¹ mole⁻²), T the absolute temperature, and π the osmotic pressure in atmospheres. One is given these quantities in the problem, but they are expressed in different units than those needed to solve for molecular weight. Converting these quantities:

Concentration: The solution is 1% by weight protein. One liter of water weighs 1000 g; 1% of 1000 is 10 so that c = 10.

Osmotic pressure: The osmotic pressure is 46 mm H₂O. To convert to atmospheres, 46 mm H₂O must be converted to mm Hg.

$$\text{mm Hg} = 46 \text{ mm H}_2\text{O} \times \frac{.9970 \text{ density of H}_2\text{O}}{13.534 \text{ density of Hg}} = 3.39.$$

There are 760 mm Hg in 1 atm. Therefore, the number of atmospheres is equal to

$$\frac{3.39 \text{ mm Hg}}{760 \text{ mm Hg/atm}} = 4.45 \times 10^{-3} \text{ atm.}$$

$$\text{Temperature: } T = 0^\circ\text{C} + 273 = 273^\circ\text{K.}$$

Solving for M:

$$M = \frac{10 \text{ g/liter}}{4.45 \times 10^{-3} \text{ atm}} \times 0.082 \frac{\text{liter-atm}}{^\circ\text{K-mole}} \times 273^\circ\text{K} = 50,300 \text{ g/mole}$$

• PROBLEM 813

The protein human plasma, albumin, has a molecular weight of 69,000. Calculate the osmotic pressure of a solution of this protein containing 2 g per 100 cm³ at 25°C in (a) Torr and (b) millimeters of water. The experiment is carried out using a salt solution for solvent and a membrane permeable to salt.

Solution: When a semipermeable membrane separates a solution of a protein in water from pure water, osmosis will occur. The concentration - or the thermodynamic activity - of water molecules in the protein solution is less than in pure water, and the system will compensate for this difference by net movement of the water from the pure water compartment into that containing the protein solution, until the concentration of the water is the same on both sides of the membrane. Osmotic pressure is the force that must be applied to just prevent such osmotic flow. Osmotic pressure can be determined by using the following equation.


$$\pi = \frac{c R T}{M},$$

$$\frac{[\alpha]_D^{25} \times \text{optical path length (dm)} \times \text{concentration (g/ml)}}{100}$$

$$\text{observed rotation} = \frac{+1.8 \times 2.5 \text{ dm} \times 9.79}{100} = 0.44^\circ.$$

• PROBLEM 815

A student wanted to produce a sample of lactic acid. He carried out the following synthesis: $\text{CH}_3\text{CH}_2\text{CO}_2\text{H} \rightarrow \text{CH}_3\text{CHClCO}_2\text{H} \rightarrow \text{CH}_3\text{CHOHCO}_2\text{H}$. He obtained a product that appeared to be lactic acid, and yet, it was optically inactive. Does this mean the product was not truly lactic acid?




Models showing the two possible arrangements in space of the four groups around the asymmetric carbon atom of lactic acid. Note that the two forms have an object-mirror image relationship and that one cannot be superimposed on the other for all points to coincide.

Solution: A carbon atom that has four different atoms or groups attached to it is termed an asymmetric carbon atom or a chiral center. These four groups can give two spatial arrangements, whose relationship to each other is that of an object and its mirror image. For example, see the accompanying figure. The two arrangements are not identical, because they are not superimposable. In such cases, these molecules, i.e. those which differ only in the positioning of groups around asymmetric carbon atoms, are called optical isomers.

Such isomers can rotate plane polarized light. Plane polarized light vibrates in only one plane. When substances can rotate plane polarized light, they are optically active. Each of the two optical isomers rotates the light in a different direction. The dextro-rotary isomer rotates it to the right, while the levo-rotary rotates it to the left. If both isomers are present in equal amounts, the light will not be rotated at all. For one rotates it to the left and the other rotates it to the right, which means there is a cancellation of the rotations. When plane-polarized light is not rotated, the solution is optically inactive. With this information, one can now answer the question.

The student formed $\text{CH}_3\text{CHOHCO}_2\text{H}$, which he believed was lactic acid. The solution was optically inactive, and yet, it is known that lactic acid is optically active. The only possible explanation for this is that the student formed equal amounts of the two possible optical isomers. As mentioned, this results in optical inactivity. There is every reason to believe that



formed in equal amounts. The star (*) indicates the asymmetric carbon. Thus, the student did form lactic acid. It is just that a mixture of the two optical isomers in equal proportions is present.

• PROBLEM 816

Insulin controls the glucose level in the blood. In 1969, Hodgkin announced the secondary structure of insulin, the first protein hormone to be resolved by X-ray diffraction. (a) Chemical analysis shows insulin is 2.15% zinc by weight. Calculate the minimum molecular weight. (b) Osmotic pressure studies show the molecular weight of insulin is about 6000. How many zinc atoms does the molecule contain?

Solution: The minimum molecular weight of insulin can be determined by assuming that there is one zinc atom per molecule. This means that in 1 mole of insulin there exists 1 mole of zinc. From the knowledge that zinc is 2.15% by weight of insulin, and that the MW of zinc is 65.3, one can determine the minimum MW of insulin by using the following relationship.

$$(.0215)(\text{MW of insulin}) = (\text{MW of zinc})$$

$$\text{MW of insulin} = \frac{\text{MW of Zn}}{(.0215)} = \frac{65.3 \text{ g/mole}}{(.0215)} = 3037 \text{ g/mole}$$

(b) If the true molecular weight of insulin is about 6000, then two of the units weighing 3037 g/mole make up each molecule of insulin. Thus, there are 2 atoms of Zn per molecule.

• PROBLEM 817

The molar absorptivity of phytochrome (a light-sensitive pigment thought to control the process of flowering in plants) is 76,000. What will be the absorbance of a 5.0×10^{-6} molar aqueous solution of phytochrome if the

Solution: For spherical molecules, the molecular weight M is related to the diffusion coefficient D by the following equality.

$$D = \frac{RT}{N_A 6\pi\eta} \left(\frac{4\pi N_A}{3M\bar{v}} \right)^{1/3},$$

where R is the gas constant in joules ($8.314 \text{ J/mole } ^\circ\text{K}$), N_A is Avogadro's Number (6.02×10^{23}), η is the viscosity of the solvent, and \bar{v} the partial specific volume of the protein.

Solving for M ;

$$D = 4.0 \times 10^{-11} \text{ m}^2/\text{s}$$

$$= \frac{(8.314 \text{ J/mole } ^\circ\text{K})(293^\circ\text{K})}{(6.02 \times 10^{23}/\text{mole})6\pi(.001005 \text{ Js/m}^2)} \times \left(\frac{4\pi(6.02 \times 10^{23} \text{ mole})}{3 \times M \times 0.75 \times 10^{-6} \text{ m}^3/\text{s}} \right)^{1/3}$$

$$4.0 \times 10^{-11} \text{ m}^2/\text{s} = (2.14 \times 10^{-19} \text{ m}^2/\text{s}) \times (3.36 \times 10^{30}/M)^{1/3}$$

$$1.87 \times 10^8 = (3.36 \times 10^{30}/M)^{1/3}$$

$$6.53 \times 10^{24} = 3.36 \times 10^{30}/M$$

$$M = 5.15 \times 10^5 \text{ g/mole.}$$

• PROBLEM 821

The sedimentation and diffusion coefficients for hemoglobin corrected to 20° in water are $4.41 \times 10^{-13} \text{ sec}$ and $6.3 \times 10^{-11} \text{ m}^2/\text{s}$, respectively. If $\bar{v} = .749 \text{ cm}^3/\text{g}$ and $\rho_{\text{H}_2\text{O}} =$

0.998 g/cm^3 at this temperature, calculate the molecular weight of the protein. If there is 1 g-atom of iron per 17,000 g of protein, how many atoms of iron are there per hemoglobin molecule?

Solution: One can calculate molecular weight by using the Svedberg equation. This equation is written

$$M = \frac{RTs}{D(1 - \bar{v}\rho)},$$

where M is the molecular weight, R is the gas constant in ergs ($8.314 \times 10^7 \text{ ergs/mole } ^\circ\text{K}$), T is the absolute temperature, s is the sedimentation coefficient, D is the diffusion coefficient, ρ is the density of the solvent,

CHAPTER 23


BIOCHEMISTRY II

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 789 to 807 for step-by-step solutions to problems.

Proteins are among the most important, if not the single most important, class of biochemical molecules. They are formed by the selective ordering of amino acids into huge molecules with molecular weights in the tens and hundreds of thousands. The peptide linkage, shown in Equation 23-1, is the basis of the bond that is used to form proteins or polypeptide chains.



This bond is hydrolyzed in acid to give back the amino and carboxylic acid groups.



Proteins are formed by polypeptide chains containing hundreds of amino acids, some of which can be found in an α -helix structure. To work those problems in this chapter that deal with the helical structure, the principles of stoichiometry can be applied directly as in problems in previous chapters.

Enzymes act as incredibly selective catalysts for biochemical reactions. They have the ability to promote one reaction while completely blocking competing reactions. The “lock and key” theory has been proposed as the mechanism for these very selective catalysts. Enzymes have a chemical


Step-by-Step Solutions to Problems in this Chapter, "Biochemistry II"

PROTEINS


• PROBLEM 822

If you hydrolyze a peptide bond in a protein molecule, what type of compounds are formed?

Solution: When proteins are boiled in dilute acids or bases they are hydrolyzed, i.e. degraded or broken down, to amino acids. Amino acids are compounds whose molecules possess both the amino ($-NH_2$) and the carboxyl ($-CO_2H$) functional groups. The general formula may be written as




The union of amino acids, which result in the protein, is due to bonds that come about as a result of the elimination of a hydrogen from the $-NH_2$ group and an OH from the $-CO_2H$ group. This linkage is termed a peptide bond. It is depicted as



peptide bond

When this substance is hydrolyzed, the following reaction occurs. In acid:



(back) the amino and carboxylic group.

Note: NH_2 is basic, so that in acid, you actually get $-\text{NH}_3^+$. Thus, you obtain a compound that possesses an amino and a carboxylic group, which is, as you recall, an amino acid. With base, the same result is obtained, but, you immediately obtain NH_2 and not NH_3^+ . The carboxylic portion, which is acidic, reacts with the OH^- from base to yield COO^- .

• PROBLEM 823

Calculate the length (in Å) of a polypeptide chain containing 105 amino acid residues if (a) it exists entirely in α -helical form, or (b) it is fully extended.

The average dimensions of the α -helix. The pitch and the rise per residue correspond to the major and minor periodicities of 5.4 and 1.5 Å, respectively. This drawing shows a left-handed α -helix.




Fig.A

Solution: (a) Peptide chains may assume the α -helix configuration spontaneously, because this form is stable and has the least free energy, providing there are no opposing interactions of the R groups or of the solvent. The α -helix is pictured in Figure A. As shown, when the peptide chain assumes this configuration each amino acid residue contributes 1.5 Å to the length of the chain. If a chain in the α -helical configuration contains 105 amino acid residues, its length is $1.5 \text{ Å} \times 105 = 157.5 \text{ Å}$.

that in each 5.41 Å of length, there are 3.6 residues. After converting 6 in to Å, one can determine the number of residues that are added per year to each strand. From this, the number of residues added per second can be calculated.

Solving for length:

$$\begin{aligned} 1 \text{ in} &= 2.54 \text{ cm}, & 1 \text{ Å} &= 10^{-8} \text{ cm} \\ \text{length in Å} &= 6 \text{ in} \times 2.54 \text{ cm/in} \times 1 \text{ Å}/10^{-8} \text{ cm} = \\ &= 1.52 \times 10^9 \text{ Å} \end{aligned}$$

Calculating the number of residues:

$$\begin{aligned} \text{number of residues} &= \text{length in Å} \times \frac{3.6 \text{ residues}}{5.41 \text{ Å}} \\ &= 1.52 \times 10^9 \text{ Å} \times \frac{3.6 \text{ residues}}{5.41 \text{ Å}} \\ &= 1.01 \times 10^9 \text{ residues} \end{aligned}$$

There are 1.01×10^9 residues produced per year.

Solving for the number of residues produced per second

residues produced per second =

$$\begin{aligned} 1.01 \times 10^9 \frac{\text{res}}{\text{year}} &\times \frac{1 \text{ year}}{365 \text{ days}} \times \frac{1 \text{ day}}{24 \text{ hrs}} \times \frac{1 \text{ hr}}{60 \text{ min}} \times \frac{1 \text{ min}}{60 \text{ sec}} \\ &= 32 \text{ res/sec.} \end{aligned}$$

• PROBLEM 826

There are some 25,000 ribosomes in an E. coli cell. If the structural proteins of these ribosomes were stretched out end to end as fully extended polypeptide chains, how many times could they encircle the E. coli cell? Assume that the ribosomes are 180 Å in diameter, with a specific gravity of 1.0, and that they contain 40% protein. Assume that the E. coli cell is a sphere 1 μ in diameter.

Solution: Ribosomes are complexes of nucleic acids and protein, and are found in the cytoplasm of the cell. This problem is solved by using the following series of steps.

(a) the volume of a ribosome

(b) the total weight of all the ribosomes present and the weight of the protein

(c) the number of amino acid residues in the protein

(d) the length of the extended protein chain and the number of times it will encircle an E. coli cell.

Solving:

(a) A ribosome is assumed to be spherical and, thus, its volume can be found by using the formula $\frac{4}{3} \pi r^3$, where r is the radius of the sphere. If the diameter of a ribosome is 180 Å its radius is 90 Å or 90×10^{-8} cm.

$$\begin{aligned}\text{volume of one ribosome} &= \frac{4}{3} \pi (90 \times 10^{-8} \text{ cm})^3 \\ &= 3.06 \times 10^{-18} \text{ cm}^3\end{aligned}$$

(b) Because the specific gravity of the ribosome is 1.0, one ribosome weighs 3.06×10^{-18} g. There are 25,000 ribosomes present.

$$\begin{aligned}\text{total weight} &= 3.06 \times 10^{-18} \text{ g/ribosome} \times 25,000 \text{ ribosomes} \\ &= 7.65 \times 10^{-14} \text{ g}\end{aligned}$$

The protein comprises 40% of the ribosome.

$$\begin{aligned}\text{weight of the protein} &= (.4) (7.65 \times 10^{-14} \text{ g}) \\ &= 3.06 \times 10^{-14} \text{ g}\end{aligned}$$

(c) The molecular weight of an average amino acid residue is 120. One can find the number of moles of amino acid residues present by dividing the weight of the protein by 120. The number of residues is found by multiplying the number of moles by 6.02×10^{23} residues/mole

$$\begin{aligned}\text{no. of amino acid residues} &= \frac{3.06 \times 10^{-14} \text{ g}}{120 \text{ g/mole}} \\ &\quad \times 6.02 \times 10^{23} \frac{\text{residues}}{\text{mole}} \\ &= 1.54 \times 10^8 \text{ residues}\end{aligned}$$

(d) When a protein is fully extended, each amino acid residue contributes 3.6 Å or 3.6×10^{-8} μ to the length of the chain.

$$\begin{aligned}\text{length of the chain} &= 3.6 \times 10^{-8} \mu \times 1.54 \times 10^8 \\ &= 5.54 \times 10^0 \mu\end{aligned}$$

The diameter of the E. coli is 1μ , which means its radius $(\frac{1}{2})\mu = 0.5\mu$. As such the circumference of the E. coli equals $2\pi r$, where r is radius, or $2\pi(0.5\mu) =$

no detectable reaction. One molecule of the enzyme amylase, containing the same amino acids, however breaks (hydrolyzes) 4000 -O- bonds per second. (a) Estimate the mass in picograms of glucose produced by an amylase molecule in 1 day. (b) The heat in calories liberated to a cell by the oxidation of the glucose: $\text{C}_6\text{H}_{12}\text{O}_6(\text{aq}) + 6\text{O}_2(\text{g}) \rightarrow 6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{l})$. $\Delta H = -686 \text{ Kcal}$. How many calories does the cell obtain from the mass of glucose calculated in part (a)?

Solution: (a) Five steps can be used to solve for the number of picograms of glucose liberated from the amylose by 1 amylase in a day.

- (1) find the number of seconds in one day
- (2) calculate the number of glucose molecules formed per day
- (3) determine the weight of one molecule of glucose
- (4) solve for the number of grams of glucose formed per day
- (5) convert from grams to picograms.

Solving:

(1) There are 60 seconds in a minute, 60 minutes in an hour and 24 hours in a day. One can solve for the number of seconds in a day by multiplying these factors together. The units will cancel out.

$$60 \text{ sec/min} \times 60 \text{ min/hour} \times 24 \text{ hours/day} = 8.64 \times 10^4 \text{ sec/day}$$

(2) The problem states that one amylase molecule liberates 4000 molecules per second. Therefore the number of molecules liberated per day can be found by multiplying the number of seconds in a day by the number of molecules formed per second.

$$\begin{aligned} \text{no. of glucose molecules formed per day} &= \\ &= 8.64 \times 10^4 \text{ sec/day} \times 4000 \text{ molecules/sec} \\ &= 3.456 \times 10^8 \text{ molecules/day} \end{aligned}$$

(3) The weight of one molecule of glucose is found by dividing the molecular weight of glucose by the number of molecules per mole. (MW of $\text{C}_6\text{H}_{12}\text{O}_6 = 180 \text{ g/mole}$, Avogadro's Number = $6.02 \times 10^{23} \text{ molecules/mole}$)

weight of one molecule =

$$= \frac{180 \text{ g/mole}}{6.02 \times 10^{23} \text{ molecule/mole}} = 2.99 \times 10^{-22} \text{ g/molecule}$$

(4) After one knows the number of molecules formed per day and the weight of one molecule, one can find the weight

in grams of the glucose formed in one day by one amylase.

no. of grams of glucose formed per day =

$$= 2.99 \times 10^{-22} \text{ g/molecule} \times 3.456 \times 10^8 \text{ molecules/day}$$

$$= 1.03 \times 10^{-13} \text{ g/day.}$$

(5) There are 10^{12} picograms in one gram. Grams can be converted to picograms by multiplying the number of grams by the conversion factor 10^{12} pg/g.

$$\text{no. of pg produced} = 1.03 \times 10^{-13} \text{ g/day} \times 10^{12} \text{ pg/g}$$

$$= 1.03 \times 10^{-1} \text{ pg/day.}$$

(6) From the equation given one knows that 686 Kcal are liberated when 1 mole of glucose is oxidized. One can find the number of calories liberated by

1) determining the number of moles of glucose present

2) solving for the number of Kcal produced

3) converting Kcal to calories.

Solving.

1) One has solved for the number of molecules produced per day previously (part (a)(2)). Therefore one can find the number of moles produced by dividing the number of molecules produced by the number of molecules in one mole.

$$\frac{3.456 \times 10^8 \text{ molecules/day}}{6.022 \times 10^{23} \text{ molecules/mole}} = 5.74 \times 10^{-16} \text{ moles/day.}$$

2) There are 686 Kcal liberated per mole of glucose oxidized, therefore the number of Kcal formed when 5.74×10^{-16} moles are oxidized is equal to

$$5.74 \times 10^{-16} \text{ moles/day} \times 686 \text{ Kcal/mole} =$$

$$= 3.94 \times 10^{-13} \text{ Kcal/day.}$$

3) There are 1000 cal in 1 Kcal, thus Kcal can be converted to cal by multiplying by the conversion factor 1000 cal/Kcal

$$\text{no. of cal produced} = 3.94 \times 10^{-13} \text{ Kcal/day} \times 1000 \text{ cal/Kcal}$$

$$= 3.94 \times 10^{-10} \text{ cal/day.}$$

• PROBLEM 832


Catalase, the enzyme that assists the breakdown of hydrogen

Solution: This question deals with proteolytic enzymes, which may be defined as protein degradation catalysts. Two factors that affect enzymic activity are pH, and temperature. Whether these enzymes damage the stomach or not will depend, to a great extent, on the conditions in the stomach. The accompanying figure shows how pH can alter enzyme activity of pepsin and trypsin. From the diagram, you see that these enzymes do not function well at extremely low pH. Due to the presence of HCl in the stomach as a hydrolysis agent, the acidity of the stomach is high, which means an environment of low pH exists. It becomes doubtful that these proteolytic enzymes would have much activity at the low pH of the stomach. This means that little, if any, damage would be done to the stomach. Heating alters enzymic activity, also. It is likely that most of the enzyme would be inactivated by the cooking of the meat.


CARBOHYDRATES

• PROBLEM 834

Explain why it has been necessary to postulate the existence of a cyclic form of glucose?



Solution: Glucose ($C_6H_{12}O_6$), is one of the isomeric aldohexoses - a carbohydrate or sugar. Glucose exists in two forms, one of which is the mirror image of the other. Its configurational formulas may be written



At one time, it was believed that these glucose isomers existed in this linear form. Now, however, it is believed that cyclic forms exist, as indicated in the accompanying diagram. This has been postulated for the following reasons: It has been found that when you take a solution of pure (+) - glucose or a solution of pure (-) - glucose form, a pair of optical isomers are obtained, i.e., ones with the ability to rotate plane-polarized light. For this to occur, in a solution of pure (+) or (-) glucose, another asymmetric carbon must be present.

An asymmetric carbon is one with 4 different atoms or molecules attached. Asymmetric carbons can generate optical isomers. In the formation of cyclic structures, the carbon atom of the aldehyde group ($C = O$) must be the additional asymmetric carbon. This would account for the two optical isomers formed. In other words, that glucose exists in cyclic forms was deduced from the fact that only in such a structure is an additional asymmetric carbon generated, which would account for the optical isomers produced.


• PROBLEM 835

There exists a carbohydrate that is slightly soluble in water, does not have a sweet taste, but does dissolve slowly when heated with dilute HCl. To what class of carbohydrates does it probably belong?

Solution: Carbohydrates may be defined as simple sugars and the substances that hydrolyze to yield simple sugars. Carbohydrates are placed in three classes: (1) Monosaccharides - those that do not undergo hydrolysis; (2) Disaccharides - those that may be hydrolyzed to two monosaccharide molecules; and (3) Polysaccharides - those which form many monosaccharide molecules after hydrolysis.


It is given that the carbohydrate dissolves in dilute HCl. Thus, one can immediately eliminate monosaccharides as a possible case. The fact that it dissolves when HCl is present suggests that a hydrolysis reaction has taken place. Only disaccharides and polysaccharides have the ability to undergo such a reaction. Thus, the carbohydrate belongs to either the class of disaccharides or polysaccharides. To determine which one it is, use the information that it is not sweet or very soluble in water. The more carbons in a molecule the more unlikely it will be soluble in water. Carbon-carbon bonds are nonpolar, while water is a polar molecule. Thus, a polysaccharide, which possesses many more non-polar bonds than a disaccharide, should have a tendency to be less soluble

100% ^{15}N . But only 50% of the nitrogen is ^{15}N . This means, therefore, that half of the DNA from the parents, which is not newly synthesized, is used in the new generation. Half of the DNA is newly synthesized (^{15}N) and half already present (^{14}N). This implies the existence of two strands. This can be better seen if you write



of that type of nitrogen, and there are 2 strands, each type makes up only 50% of the total nitrogen.

(b) The new generation is reproducing on ^{14}N . The progeny will have one half of its DNA newly synthesized on ^{14}N and half of its DNA already present. You can picture it as the following:



The starred nitrogen 14 is that already present. The unstarred was that newly synthesized. From this, you see that half of the second new DNA generation would be expected to contain only ^{14}N strands (starred and unstarred), and the other half would contain 50% ^{14}N and 50% ^{15}N .

POLLUTION PROBLEMS

• PROBLEM 843

One of the major atmospheric pollutants emitted by fuel-combustion power stations is a sulfur oxide mixture generally designated as SO_x . It consists mainly of SO_2 but may contain anywhere from 1 to 10 % SO_3 computed as percent by weight of the total mass. What would be the approximate range to assign to x ?

Solution: One finds x by determining the number of O atoms present for each S present. For purposes of calculation assume that one always has 100 g of the mixture of SO_2 and SO_3 . Thus when 1 % of the mixture is SO_3 , there will be 1 g of SO_3 present and 99 g of SO_2 present. One then finds the number of moles present of each element. The ratio of S : O is then determined.

Solving for x when 1 % of the mixture is SO_3 :
(MW of SO_2 = 64, MW of SO_3 = 80). The number of moles present is equal to the number of grams present divided by the molecular weight.

$$\text{For } \text{SO}_3: \frac{1 \text{ g}}{80 \text{ g/mole}} = 1.25 \times 10^{-2} \text{ moles}$$

$$\text{For } \text{SO}_2: \frac{99 \text{ g}}{64 \text{ g/mole}} = 1.55 \text{ moles.}$$

Because there is one S present in each compound the total amount of S is 1.55 moles + 1.25×10^{-2} moles or 1.56 moles. In 1 mole of SO_3 there are 3 moles of O, therefore in 1.25×10^{-2} moles of SO_3 there is three times that amount.

$$\text{no. of moles of O in } \text{SO}_3 = 3 \times 1.25 \times 10^{-2} = 3.75 \times 10^{-2}$$

There are 2 moles of O in 1 mole of SO_2 , thus in 1.55 moles of SO_2 there are 3.10 moles of O. The total number of moles of O present is equal to the sum of the moles contributed by the SO_2 and the SO_3 .

$$\text{total no. of moles of O} = 3.75 \times 10^{-2} + 3.10 = 3.14 \text{ moles}$$

The formula for this compound is $\text{S}_{1.56}\text{O}_{3.14}$. The simplest formula is $\text{SO}_{2.01}$. When there is 1 % of SO_2 present, $x = 2.01$. $x = 3.14/1.56$.

When there is 10 % of SO_3 present, a similar method is used to find x.

- 1) Find the number of moles present of SO_2 and SO_3 .
- 2) Determine the number of moles of S and O contributed by the SO_2 and SO_3 .
- 3) write the formula and simplify.

Solving for x:

1) For a 100 g sample of gas where 10 % of the total weight is SO_3 , 10 g of the gas is SO_3 and 90 g is SO_2 .

$$\text{no. of moles of } \text{SO}_3 = \frac{10 \text{ g}}{80 \text{ g/mole}} = 1.25 \times 10^{-1} \text{ moles.}$$

$$\text{no. of moles of } \text{SO}_2 = \frac{90 \text{ g}}{64 \text{ g/mole}} = 1.41 \text{ moles}$$

2) The total number of moles of S present is equal to the sum of the number of moles of SO_2 and SO_3 present because each compound contributes 1 S per mole of compound present.

$$\text{no. of moles of S} = 1.25 \times 10^{-1} + 1.41 = 1.54 \text{ moles}$$

$$\begin{aligned} \text{no. of moles of O from } \text{SO}_3 &= 3 \times 1.25 \times 10^{-1} \\ &= 3.75 \times 10^{-1} \text{ moles.} \end{aligned}$$

$$\text{no. of moles of O from } \text{SO}_2 = 2 \times 1.41 = 2.82 \text{ moles}$$

$$\text{total no. of moles of O} = 3.75 \times 10^{-1} + 2.82 = 3.20 \text{ moles.}$$

3) The formula for this compound can be written $\text{S}_{1.54}\text{O}_{3.20}$. $x = 3.20/1.54 = 2.08$, thus the simplest formula is $\text{SO}_{2.08}$.

The range of x then is from 2.01 to 2.08.

• PROBLEM 844

Analysis of the exhaust composition of the supersonic transport for one hour of flight yielded the following information:

Compound	Molecular Weight (g/mole)	Mass (g)
H_2O	18	8.0×10^7
CO_2	44	6.6×10^7
CO	28	3.6×10^6
NO	30	3.6×10^6

Compute how many moles of H_2PO_4^- are present. A mole is defined as grams (weight)/molecular weight (M.W.). The M.W. of H_2PO_4^- = 96.97 g/mole. The number of moles of H_2PO_4^- is

$$= \frac{567,454 \text{ g}}{96.97 \text{ g/mole}} = 5852 \text{ moles.}$$

According to the reaction equation, one mole of H_2PO_4^- generates one mole of precipitate. The M.W. of the precipitate = 245 g/mole. Recalling the definition of a mole, the amount of $\text{Mg}(\text{NH}_4)\text{PO}_4 \cdot \text{H}_2\text{O}$ = $(5852)(245) = 1.433 \times 10^6 \text{ g} = 1.433 \times 10^3 \text{ kg}$.


• PROBLEM 8

The SO_2 content in air may be determined by passing the air through water to produce a solution of sulfurous acid, H_2SO_3 . The subsequent reaction of this solution with potassium permanganate, KMnO_4 , establishes the amount of SO_2 absorbed from the air. (a) Write the balanced equation for the reaction of SO_2 with water to produce H_2SO_3 . (b) Write the balanced equation for the reaction of H_2SO_3 (or SO_3^{2-}) with KMnO_4 (or MnO_4^-) to form MnSO_4 , K^+ and H_2SO_4 , among other products. (c) If 1000 liters of air are passed through water and the resulting solution requires 2.5×10^{-5} moles of KMnO_4 for complete reaction (part b), what is the concentration (in ppm) of SO_2 in the air. Assume STP conditions.

Solution: (a) An SO_2 molecule reacts with a water molecule to form an H_2SO_3 molecule according to the following equation:




However, H_2SO_3 is a diprotic acid and dissociates into 2 protons and a sulfite ion



Thus, for every mole of SO_2 consumed 1 mole of SO_3^{2-} are produced.

(b) This reaction is a typical redox reaction and the equation must be balanced using the standard rules.

From the problem, one knows initially that



This reaction is not balanced in terms of charge and elemental constituents. The first rule is to separate the above reaction into a reduction and an oxidation reaction.



Mn is reduced from +7 to +2 state.

2 moles of H_2O .

To solve this problem, calculate how many moles of SO_2 are polluted into the air by dividing its weight by its molecular weight (MW $\text{SO}_2 = 64 \text{ g/mole}$).

From the stoichiometry of the equation, determine how many moles of S would be produced, and then calculate the weight of S by multiplying the number of moles of S times its molecular weight (MW S = 32 g/mole). Thus,


$$\text{number moles of } \text{SO}_2 = \frac{650,000,000 \text{ g } \text{SO}_2}{64 \text{ g/mole}} = 1.02 \times 10^7 \text{ moles}$$

From the stoichiometry of the equation 1 mole $\text{SO}_2 + 3 \text{ mole S}$ or $1.02 \times 10^7 \text{ mole } \text{SO}_2 + 3.06 \times 10^7 \text{ mole S}$.

$$\begin{aligned}\text{weight of S} &= (3.06 \times 10^7 \text{ mole S})(32 \text{ g/mole}) \\ &= 97.9 \times 10^7 \text{ g} = 979,000 \text{ kg}.\end{aligned}$$

• PROBLEM 848

Iodine pentoxide (I_2O_5) is a very important reagent because it can oxidize carbon monoxide (CO), which is a major pollutant of the atmosphere, to carbon dioxide (CO_2) in the absence of water. The reaction involved is



How many kg of I_2O_5 is needed to completely clear 140,000 kg of CO from the air?

Solution: From the stoichiometry of this equation, 1 mole of I_2O_5 completely reacts with 5 moles of CO. To solve this problem, calculate the number of moles of CO that will react by dividing its weight by its molecular weight of 28 g/mole. Determine, also, how many moles of I_2O_5 will completely react with this, and then calculate the weight of I_2O_5 that will completely react by multiplying the number of moles of I_2O_5 by its molecular weight (MW of $\text{I}_2\text{O}_5 = 334 \text{ g/mole}$). Thus,

$$\text{number moles CO} = \frac{140,000,000 \text{ g CO}}{28 \text{ g/mole}} = 5.00 \times 10^6 \text{ mole}.$$

This means that one fifth of this number of moles will be the moles of I_2O_5 required. Recall, there will exist 5 moles of CO for every mole of I_2O_5 . Therefore,

$$\text{moles } \text{I}_2\text{O}_5 = \frac{1}{5}(5 \times 10^6) = 1 \times 10^6 \text{ moles}.$$

This means that

$$(1 \times 10^6 \text{ moles})(334 \text{ g/mole}) = 3.34 \times 10^8 \text{ grams of}$$

I_2O_5 is needed to completely clear 140,000 kg of CO from the air.

• PROBLEM 8

The average concentration of particulate lead in a city with highly polluted air is $5 \mu\text{g}/\text{m}^3$. If an adult respires 8500 liters of air daily and if 50 per cent of the particles below $1 \mu\text{m}$ in size are retained in the lungs, calculate how much lead is absorbed in the lungs each day. Of the particulate lead in urban air, 75 per cent is less than $1 \mu\text{m}$ in size.

Solution: To solve this problem (1) determine the total percentage of particulate lead remaining in the lungs, (2) determine the amount of lead absorbed into the lungs.

The problem states 75 % of particulate lead is less than $1 \mu\text{m}$ in size, therefore, 25 % of the lead is greater than $1 \mu\text{m}$ in size and does not remain inside the lungs. Of the 75 % particulate lead less than $1 \mu\text{m}$ in size, only 50 % of this is retained in the lungs. The total percentage of lead retained is 37.5 %.

The total volume of air-intake is 8500 liters, so that
total volume of air-intake = $(8500)(10^{-3} \text{ m}^3/\text{liter}) = 8.5 \text{ m}^3$

The total amount of lead-intake is
amount of lead = $(8.5 \text{ m}^3)(5 \mu\text{g}/\text{m}^3) = 42.5 \mu\text{g}$ lead.

The total amount of lead absorbed is
amount of lead absorbed = $(42.5 \mu\text{g})(0.375) = 15.9 \mu\text{g}$.

• PROBLEM 850

In 1932, many cities showed higher sunlight levels than in the previous year. Explain this observation.

Solution: To answer this problem, consider the effect of atmospheric particles on the amount of sunlight received and what happened in 1932.

By 1932 three years of slackened industrial activity had passed. Then as now, industries released massive amounts of particulate matter, i.e. solid and liquid aerosols, into the atmosphere. These particles have the ability to scatter and absorb light. Because industrial activity was drastically reduced, fewer particles were released that could scatter or absorb the sunlight. The cities, therefore,

received more sunlight in 1932 than in years when industry was thriving.


It is interesting to note that because of particulate matter, cities receive about 15 to 20 % less solar radiation than do rural areas.

• PROBLEM 851


The air pollutant, sulfur dioxide, is considered particularly foul. Explain why there is so much concern with this compound.

Solution: To solve this problem, consider the reactions that can occur between sulfur dioxide (SO_2) and other elements, and/or compounds, in the natural environment.


SO_2 can enter the atmosphere from both natural and man-made sources. Besides causing eye irritation and plant damage, it can cause other health problems because of its ability to form acids in the atmosphere. When SO_2 comes in contact with water (H_2O), the following reaction occurs.




SO_2 can also be oxidized in air to sulfur trioxide (SO_3) by the reaction



Sulfur trioxide, when exposed to water, produces sulfuric acid by the reaction



These two acids, sulfuric and sulfurous, account for the corrosion of metal parts exposed to the atmosphere. They are also responsible for the decay of stone statues and buildings. This occurs because most stones consist of CaCO_3 (calcium carbonate), which, upon addition of H_2SO_4 , undergoes the following reaction:



CaSO_4 is soluble in water and will wash away with the rain causing the wearing away of the stone.

• PROBLEM 852

If the carbon dioxide content of the air doubled, what

then reacts with oxygen (diatomic - O_2) to produce O_3 (ozone), a toxic element in the lower atmosphere. However, the ozone reacts with the NO that was previously produced to again form NO_2 , which allows the cycle to repeat.

Now, based on the above information, consider what happens at rush hour. The cars and busses release hydrocarbons. Among these fragments of hydrocarbons are free radicals which are extremely reactive. They proceed to react with NO , reducing its concentration. When the NO is depleted, there is little of it available to react with ozone to produce NO_2 and O_2 . Thus, O_3 builds up. It is found that this process takes 2 hours, which explains why the O_3 concentration increases after the peak hydrocarbon concentration.

• PROBLEM 854

Given that the earth's mean radius = 6.37×10^6 m, normal atmosphere pressure = 1.013×10^5 N/m², and the gravitational acceleration = 9.8 m/sec², what is the mass of the homosphere?

Solution: The homosphere is that part of the atmosphere that extends from the surface of the earth up to 80 kilometers. The constituent gases in the homosphere are well mixed. The proportions of the principal constituents at high altitudes do not differ to a great extent from those found at sea level. To find the mass of the homosphere, calculate the total mass of the atmosphere.

The homosphere is equal to 99.99% of the atmosphere and, hence, their masses are approximately equal.

The mass of the atmosphere can be derived from the atmospheric pressure. Since

$$\text{Pressure} = \frac{\text{Force}}{\text{Area}} \quad \text{and} \quad \text{Force} = \text{mass} \times \text{gravity}$$

Combining these two equations one obtains

$$\text{Pressure} = \frac{\text{mass} \times \text{gravity}}{\text{Area}}$$

Since one is solving for mass the equation can be rearranged to

$$\text{mass} = \frac{\text{Pressure} \times \text{Area}}{\text{gravity}}$$

One is given the atmospheric pressure and the force of gravity. The surface area of a sphere is equal to $4\pi r^2$. Substituting and solving for the mass of the homosphere: (N = kg-m/s²)

amounts of algae can lead to the death of marine life. This is one major problem associated with eutrophication.

A second problem also results from the depletion of oxygen. Remember that anaerobic bacteria, i.e. bacteria that can live without oxygen, still function effectively under the conditions described above. These bacteria feed on decaying algae. As a result of their metabolic process, they produce compounds such as hydrogen sulfide, which cause the putrid odor associated with decaying organic matter. The water becomes foul smelling and foul tasting.

A reduction of phosphate would alleviate eutrophication because it is one of the major nutrients of algae. A smaller amount of phosphate in the water would limit the growth of algae.

• PROBLEM 857

The combined effects of pollution have been known to bring about the phenomena of synergism and antagonism. Explain each of these terms. Provide specific examples of each.

Solution: The effects of individual pollutants can be disastrous to health, while the combined action of two or more pollutants can be completely different on health and/or the environment.

When the combined effects of two or more pollutants are more severe OR qualitatively different from the individual effects, a condition called synergism exists. For example, in 1966 and 1967, there was injury to peanut crops due to the synergistic action of ozone and sulfur dioxide.

A condition termed antagonism exists when the combined effects are less severe. For example, cyanides in industrial wastes are quite poisonous to aquatic life, however, in the presence of nickel, a nickel-cyanide complex is formed. This nickel-cyanide complex is not as toxic as the cyanides originally produced.

• PROBLEM 858

(A) Explain the purpose of a coliform bacteria count in natural water. (B) What is meant by hard or refractory organic compounds?

Solution: (a) One of the most important problems associated with polluted water is the possible presence of disease-causing bacteria. A test for the presence of bacteria is made by measuring the amount of common coliform bacteria in water. (Coliform bacteria are harmless

and reside in the large intestines of humans.) Coliform bacteria are present in plentiful quantities in feces, which become part of sewage wastes. Since coliform bacteria do not thrive in an aquatic environment, they give an indication of how recently and to what extent sewage pollution has occurred by their count (which is proportional to the amount of disease-causing bacteria present). If the count is high, human sewage contamination has taken place, possibly accompanied by the presence of dangerous bacteria.


(b) Hard or refractory organic compounds are nondegradable substances that decay very slowly in water. Degradation refers to the breaking down of substances into simpler forms. Examples of nondegradable substances are certain pesticides and detergents.

● PROBLEM 859


A chemist has a quantity of steam. He wants to determine whether or not it has been polluted with acidic mine drainage (AMD). How should he go about determining this?

Solution: AMD occurs yearly when millions of tons of sulfuric acid seep out from coal and copper mines to pollute waterways.


It begins by the oxidation of iron pyrite (FeS_2) to sulfate:




The ferrous ions are then oxidized to ferric ions:



This process can be sped up by microorganisms. The ferric ions can react with water to give hydrogen ions and a precipitate of hydrated ferric oxide:



Thus, acidic mine drainage is a diluted sulfuric acid solution carrying iron. If the steam has AMD, it must carry iron. Addition of thiocyanate to ferric (iron) solutions will yield an intensely red-colored complex ion. The reaction is as follows:



Thus, the chemist can detect for AMD pollution in the steam by adding thiocyanate. If a red-colored material is formed, he knows it is contaminated.



APPLIED GAS PROBLEMS

• PROBLEM 861

The pressure in an automobile tire is measured by a tire gauge in winter at 0°C as 30 pounds per square inch (psi) (2.0415 atm). The gauge measures the difference between the actual tire pressure and the atmospheric pressure (15 psi). Assume that the tire does not leak air and does not change its volume. What will the new gauge pressure be if the same tire is measured during the summer at 50°C?

Solution: We will solve this problem by employing a modified form of the ideal gas law. The ideal gas equation is $PV = nRT$, where P is the total pressure (P = atmospheric pressure + gauge pressure), V is the tire volume, n the number of moles of air, R the gas constant, and T the absolute temperature. Bearing in mind the fact that n , V , and R are constant, we may write this equation for the two separate sets of conditions, as

$$P_{\text{summer}} V = n R T_{\text{summer}} \quad \text{and} \quad P_{\text{winter}} V = n R T_{\text{winter}}$$

Dividing the first by the second gives

$$\frac{P_{\text{summer}} V}{P_{\text{winter}} V} = \frac{n R T_{\text{summer}}}{n R T_{\text{winter}}} \quad \text{or} \quad \frac{P_{\text{summer}}}{P_{\text{winter}}} = \frac{T_{\text{summer}}}{T_{\text{winter}}}$$

Solving for P_{summer} , we have

$$P_{\text{summer}} = \frac{T_{\text{summer}}}{T_{\text{winter}}} \times P_{\text{winter}}$$

$$P_{\text{winter}} = \text{atmospheric pressure} + \text{gauge pressure}$$

$$= 15 \text{ psi} + 30 \text{ psi} = 45 \text{ psi} \text{ (3.06 atm).}$$

$T_{\text{summer}} = 50^{\circ}\text{C} = 323^{\circ}\text{K}$, and $T_{\text{winter}} = 0^{\circ}\text{C} = 273^{\circ}\text{K}$. Hence,

$$P_{\text{summer}} = \frac{T_{\text{summer}}}{T_{\text{winter}}} \times P_{\text{winter}} = \frac{323^{\circ}\text{K}}{273^{\circ}\text{K}} \times 45 \text{ psi} = 53 \text{ psi.} \\ (3.606 \text{ atm})$$

The gauge pressure will be $P_{\text{summer}} - \text{atmospheric pressure}$, or $53 \text{ psi} - 15 \text{ psi} = 38 \text{ psi}$ (2.586 atm)

• PROBLEM 862

The pressure in interplanetary space is estimated to be of the order of 10^{-16} torr. This corresponds to a density of about 4 molecules per cubic centimeter. What temperature must be assigned to the molecules in interplanetary space for this to be true?

Solution: To solve this problem, one assumes that interplanetary space is an ideal gas at a low concentration and at a very low pressure. This means that the ideal gas law can be used, i.e.,

$$PV = nRT$$

The pressure, P , is known 10^{-16} torr = 10^{-16} torr \times (1 atm/760 torr) = 1.32×10^{-19} atm and R , the gas constant = 0.082 atm-liter/mole- $^{\circ}\text{K}$, is also known. The volume, V , and the number of moles, n , are unknown, but if n is divided by V , then this value becomes a density (moles/liter) and can be determined from the given density of 4 molecules/cm³. To do this, (1) convert the number of molecules to moles using Avogadro's number and (2) convert cm³ to units of l using the conversion 1 cm³ = 1 ml = 0.001 l. Thus, the density is

$$\left(\frac{4 \text{ molecules}}{1 \text{ cm}^3} \right) \left(\frac{1000 \text{ cm}^3/\text{l}}{6.02 \times 10^{23} \frac{\text{molecules}}{\text{mole}}} \right) \\ = 6.64 \times 10^{-21} \text{ moles/l}$$

Rearranging the ideal gas law from $PV = nRT$ to $\frac{P}{R} \cdot \frac{V}{n} = T$, and substituting the known values, yields

$$T = \frac{(1.32 \times 10^{-19} \text{ atm})}{\left(0.082 \frac{\text{atm-liter}}{^{\circ}\text{K-mole}} \right)} \times \left(\frac{1}{6.64 \times 10^{-21} \frac{\text{moles}}{\text{l}}} \right) \\ = 2.42 \times 10^2 \text{ }^{\circ}\text{K}$$

$$T = 242^{\circ}\text{K} = 242^{\circ}\text{K} - 273^{\circ} = -31^{\circ}\text{C}.$$

• PROBLEM 863

Assume that a thermonuclear explosion 1000 ft above ground quickly heats 10,000 ft.³ of air at standard pressure from 273°K to 1,092,273°K. The air would attempt to expand to what volume as an outgoing pressure-blast wave develops?

Solution: To solve this problem, Charles' Law is used,

$$\frac{V_1}{V_2} = \frac{T_1}{T_2} ,$$

where V = volume and T = absolute temperature. The pressure, (since the expansion is done in open space) remains at standard pressure of 1 atm. Thus,

$V_1 = 10,000 \text{ ft.}^3$, $T_1 = 273^\circ\text{K}$, $T_2 = 1,092,273^\circ\text{K}$
and V_2 is solved for

$$V_2 = \frac{V_1 T_2}{T_1} = \frac{(10,000 \text{ ft.}^3)(1,092,273^\circ\text{K})}{(273^\circ\text{K})} = 40,010,000 \text{ ft.}^3.$$

The air expands by a factor of

$$\frac{40,010,000 \text{ ft.}^3}{10,000 \text{ ft.}^3} = 4,001.$$

• PROBLEM 864

An aerosol spray at room temperature, 25°C, is under a pressure of 10 atm at a volume of 0.5 l. If someone releases all the contents to fill an empty 4.6 l box at standard pressure, what is the temperature of the contents inside the box?

Solution: To solve this problem the general gas law, assuming ideal gas conditions, for two different states is used.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} ,$$

where $P_1 = 10 \text{ atm}$, $V_1 = 0.5 \text{ l}$, $T_1 = 298^\circ\text{K}$, $P_2 = 1 \text{ atm}$, (standard pressure), $V_2 = 4.6 \text{ l}$, and T_2 is unknown. Solving for T_2 and substituting the known values into this equation gives

$$T_2 = \frac{P_2 V_2 T_1}{P_1 V_1} = \frac{(1 \text{ atm})(4.6 \text{ l})(298^\circ\text{K})}{(10 \text{ atm})(0.5 \text{ l})} = 274^\circ\text{K} = 1^\circ\text{C}.$$

The decrease in temperature is significant; the change is from 25°C to near freezing.

• PROBLEM 865

In the Van Slyke method for determining CO_2 capacity in blood, you place the sample over mercury in a closed flask. CO_2 is released from the sample by the addition of acid. The volume and pressure of the released is then measured. In a 0.2 ml sample of blood, the CO_2 released exerts a pressure of 162 mm Hg at a temperature of 27°C and occupies a volume of 0.5 cc. What is the corresponding volume of the CO_2 at standard temperature and pressure (STP)?

Solution: STP refers to conditions where the temperature = 0°C or 273°K and the pressure = 760 mm Hg or 1 atm. To find the volume of the CO_2 released under these conditions, apply the Combined Gas Law.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} = \text{constant}$$

where P = pressure, V = volume, and T = absolute temperature and the subscripts refer to different states.

To solve this problem substitute the known values into this equation:

$$P_1 = 162 \text{ mm}$$

$$P_2 = 760 \text{ mm}$$

$$V_1 = 0.5 \text{ cc} = 0.5 \text{ ml}$$

$$V_2 = ?$$

$$T_1 = 27^\circ\text{C} = 300^\circ\text{K}$$

$$T_2 = 273^\circ\text{K}$$

where P_2 and T_2 are the standard conditions at STP. Thus,

$$\frac{(162 \text{ mm})(0.5 \text{ ml})}{(300^\circ\text{K})} = \frac{(760 \text{ mm})V_2}{(273^\circ\text{K})}$$

Solve for V_2 , the volume of CO_2 at STP,

$$V_2 = \frac{(162 \text{ mm})(0.5 \text{ ml})(273^\circ\text{K})}{(760 \text{ mm})(300^\circ\text{K})} = 0.0969 \text{ ml} = 0.0969 \text{ cc.}$$

• PROBLEM 866

Carbon monoxide poisons humans by irreversibly binding to hemoglobin in the bloodstream. Although not dangerous in minute concentrations over long exposure times, a CO concentration of 0.4 % by volume is quickly lethal. If a defective automobile in a sealed garage having a volume of 4.1×10^4 liters produces 0.60 mole of CO per minute, how long will it take to reach this lethal concentration of CO? Assume that the volume remains constant at 27°C and the pressure at 1 atm.

Solution: This problem is solved by first converting the

concentration, 0.4 % by volume, to a partial pressure and determining the number of moles that will give rise to this pressure. The time it takes to reach this pressure is then determined by dividing the number of moles of CO corresponding to this pressure by the rate of CO formation.

When the concentration of CO is lethal, it occupies 0.4 % of the atmosphere by volume. Therefore, its partial pressure, p , is 0.4 % of the total pressure, or $p = 0.4 \% \times 1 \text{ atm} = 0.004 \text{ atm}$.

The ideal gas equation reads $pV = nRT$, or $n = \frac{pV}{RT}$ where n is the number of moles, V is the volume, R the gas constant, and T the absolute temperature. Since $V = 4.1 \times 10^4$ liters, $R = 0.082 \text{ liter-atm/mole-deg}$, and $T = 27^\circ\text{C} = 300^\circ\text{K}$, the number of moles of CO corresponding to a partial pressure of 0.004 atm is

$$n = \frac{pV}{RT} = \frac{0.004 \text{ atm} \times 4.1 \times 10^4 \text{ liters}}{0.082 \text{ liter-atm/mole-degree} \times 300^\circ\text{K}} = 6.7 \text{ moles.}$$

Since the automobile produces CO at a rate of 0.60 mole/min, it will take 6.7 moles/0.60 mole/min \approx 11 minutes for CO to reach the lethal concentration.

• PROBLEM 867

The average density of the Universe is very low, various estimates of the average density being between $1.0 \times 10^{-30} \text{ g/cm}^3$ and $3.0 \times 10^{-28} \text{ g/cm}^3$. Using an intermediate value for the average density, $1.5 \times 10^{-29} \text{ g/cm}^3$, and assuming that the Universe consists solely of hydrogen atoms, what is the average volume of space that contains a single hydrogen atom?

Solution: To solve this problem, we will convert the density (in g/cm^3) to a molar density (moles/ cm^3) and then express this as the number of atoms per unit volume (number density). Inverting the number density will give us the volume occupied by a single atom.

To convert from density to molar density, we divide by the atomic weight of hydrogen (1.0 g/mole). This stems from the fact that grams/atomic weight = mole. Hence,

$$\begin{aligned} \text{molar density} &= \text{density} \div \text{atomic weight} \\ &= 1.5 \times 10^{-29} \text{ g/cm}^3 \div 1.0 \text{ g/mole} \\ &= 1.5 \times 10^{-29} \text{ mole/cm}^3. \end{aligned}$$

To convert from molar density to number density, we must multiply by the number of atoms per mole (Avogadro's number). Hence

$$\begin{aligned}
 \text{number density} &= \text{molar density} \times \text{Avogadro's number} \\
 &= 1.5 \times 10^{-23} \text{ mole/cm}^3 \times 6 \times 10^{23} \text{ atoms/mole} \\
 &= 9.0 \times 10^{-6} \text{ atom/cm}^3.
 \end{aligned}$$

The average volume occupied by a single H atom is the reciprocal of the number density, or

$$\begin{aligned}
 \text{volume per atom} &= \frac{1}{\text{number density}} = \frac{1}{9.0 \times 10^{-6} \text{ atom/cm}^3} \\
 &= 1.1 \times 10^5 \text{ cm}^3/\text{atom}.
 \end{aligned}$$

Hence, on the average, each $1.1 \times 10^5 \text{ cm}^3$ (110 liters) is occupied by a single hydrogen atom.

• PROBLEM 868

An anaesthetic can be prepared by mixing gaseous cyclopropane (C_3H_6 , molecular weight = 42 g/mole), and oxygen (O_2 , molecular weight = 32 g/mole). If a gas cylinder is prepared with cyclopropane at a partial pressure of 170 torr (0.223 atm) and oxygen at a partial pressure of 570 torr (0.75 atm), calculate the ratio of the number of moles of cyclopropane to that of oxygen $\left(n_{\text{C}_3\text{H}_6} / n_{\text{H}_2\text{O}} \right)$.

Solution: We solve this problem by relating the ratio of moles, $n_{\text{C}_3\text{H}_6} / n_{\text{H}_2\text{O}}$, to the ratio of partial pressures, $P_{\text{C}_3\text{H}_6} / P_{\text{H}_2\text{O}}$. For a given volume V and fixed absolute temperature T , the ideal gas law reads:

$$P_{\text{C}_3\text{H}_6} = n_{\text{C}_3\text{H}_6} \frac{RT}{V}$$

for cyclopropane, and

$$P_{\text{O}_2} = n_{\text{O}_2} \frac{RT}{V}$$

for oxygen, where R is the gas constant. Dividing the first of these equations by the second gives

$$\frac{P_{\text{C}_3\text{H}_6}}{P_{\text{O}_2}} = \frac{n_{\text{C}_3\text{H}_6} \frac{RT}{V}}{n_{\text{O}_2} \frac{RT}{V}} = \frac{n_{\text{C}_3\text{H}_6}}{n_{\text{O}_2}}$$

Since $P_{\text{C}_3\text{H}_6} = 170 \text{ torr}$ and $P_{\text{O}_2} = 570 \text{ torr}$,

$$\frac{P_{\text{C}_3\text{H}_6}}{P_{\text{O}_2}} = \frac{170 \text{ torr}}{570 \text{ torr}} = 0.30 = \frac{n_{\text{C}_3\text{H}_6}}{n_{\text{O}_2}}$$

or, $n_{\text{C}_3\text{H}_6} / n_{\text{O}_2} = 0.30$.

The vapor pressure of water is 17.54 mm at 20°C and 9.21 mm at 10°C.

Solution: (A) Relative humidity is defined as the ratio of the partial pressure of water vapor in the air to the vapor pressure of water at that temperature of the air. Given that the partial pressure of water vapor at 20°C is 10 mm and that its vapor pressure is 17.54 mm at 20°C:

$$\text{relative humidity} = \frac{10 \text{ mm}}{17.54 \text{ mm}} = 0.57.$$

Expressed as a percent; relative humidity = 57 %.

(B) Most of the time, air is not saturated with water vapor. However, it can be saturated if it is cooled to a temperature where the vapor pressure of water becomes equal to the partial pressure of the water vapor in the air. This is the temperature of the dew point. To calculate the dew point in this problem, change the temperature until the vapor pressure equals 10 mm (the partial pressure). Given that the vapor pressure is 17.54 at 20°C and 9.21 at 10°C, the dew point must be between these two temperatures. To find the exact temperature (the dew point) the following proportion can be set up where 10 mm is the vapor pressure: Let x = dew point.

$$\frac{10 \text{ mm}}{17.54 \text{ mm}} = \frac{x}{20^\circ\text{C}}$$

Solving for x ; $x = 11.4^\circ\text{C}$.

• PROBLEM 871

A controlled environment for guinea pigs is made by constructing a 294 liter box to be maintained at 21°C and 40 % relative humidity, and in which the entire air volume is changed every minute. In order to maintain this relative humidity, what weight of water must be added to the dry air flow every minute. The equilibrium vapor pressure for water at 21°C is about 19 torr.

Solution: To solve this problem, we use the relative humidity and the equilibrium vapor pressure of water to obtain the partial pressure of water. The ideal gas law is then used to calculate the amount of water corresponding to this partial pressure.

The relative humidity is defined as the percent ratio of the partial pressure p to the equilibrium vapor pressure, p_e . Hence,

$$\text{relative humidity} = \frac{p}{p_e} \times 100 \%, \text{ or}$$

$$p = p_e \times \frac{\text{relative humidity}}{100 \%} = 19 \text{ torr} \times \frac{40 \%}{100 \%}$$

$$= 19 \times \frac{40}{100} \text{ torr} = 19 \times 0.4 \text{ torr}.$$

Converting to atmospheres, $p = 19 \times 0.4 \text{ torr} \times 1 \text{ atm}/760 \text{ torr} = 0.01 \text{ atm}$.

The ideal gas equation may be written as

$$pV = nRT = \frac{m}{MW} RT,$$

where V is the volume ($V = 294$ liters), R is the gas constant ($R = 0.082$ liter-atm/mole-degree), T is the absolute temperature ($T = 21^\circ\text{C} = 294^\circ\text{K}$), and n = numbers of moles = m/MW , where m is the mass, and MW is the molecular weight of H_2O ($MW = 18$ g/mole). Solving for m we obtain

$$m = \frac{PV(MW)}{RT} = \frac{0.01 \text{ atm} \times 294 \text{ liters} \times 18 \text{ g/mole}}{0.082 \text{ liter-atm/mole-degree} \times 294^\circ\text{K}} \approx 2.2 \text{ g}.$$

Thus, 2.2 g of H_2O must be supplied to the dry air flow every minute to maintain a relative humidity of 40 %.

• PROBLEM 872

The pistons in an automobile engine are driven by the following reaction between octane (C_8H_{18} , molecular weight = 114 g/mole), and oxygen (O_2 , molecular weight = 32 g/mole), which takes place in the cylinders.



If the cylinders have a total volume of 6.15 liters, and one-fifth of the air filling this volume at 1 atm and 27°C is oxygen, what weight of octane is necessary to combine exactly with the oxygen?

Solution: To solve this problem, we first determine the partial pressure of oxygen. Using this value and the ideal gas law, we then calculate the number of moles of oxygen present and finally, the number of moles of octane required to react with it.

One-fifth the total volume is occupied by oxygen. This corresponds to 20 % oxygen by volume ($1/5 \times 100 \% = 20 \%$). The partial pressure of oxygen, p_{O_2} , is equal to the percent oxygen multiplied by the total pressure, or $p_{\text{O}_2} = 20 \% \times 1 \text{ atm} = 0.20 \text{ atm}$.

The number of moles, n_{O_2} , of oxygen present is calculated from the ideal gas equation,

$$n_{O_2} = \frac{P_{O_2} V}{RT},$$

where V is the volume ($V = 6.15$ liters), R is the gas constant ($R = 0.082$ liter-atm/mole-degree), and T is the absolute temperature ($T = 27^\circ\text{C} = 300^\circ\text{K}$). Then,

$$n_{O_2} = \frac{P_{O_2} V}{RT} = \frac{0.20 \text{ atm} \times 6.15 \text{ liters}}{0.082 \text{ liter-atm/mole-deg} \times 300^\circ\text{K}} = 0.05 \text{ mole.}$$

From the equation for the reaction between octane and O_2 , we see that 2 moles of octane react with 25 moles of O_2 . Setting up the following proportion

$$\frac{2 \text{ moles } C_8H_{18}}{25 \text{ moles } O_2} = \frac{n_{C_8H_{18}}}{n_{O_2}}$$

and solving for the number of moles of octane, $n_{C_8H_{18}}$, we obtain


$$n_{C_8H_{18}} = \frac{2 \text{ moles}}{25 \text{ moles}} \times n_{O_2} = \frac{2 \text{ moles}}{25 \text{ moles}} \times 0.05 = 0.004 \text{ mole.}$$

The required mass of octane is obtained by multiplying the number of moles of octane by the molecular weight of octane, or

$$0.004 \text{ mole} \times 114 \text{ g/mole} = 0.46 \text{ g.}$$

• PROBLEM 873

SO_2 (sulfur dioxide) can be used as a preservative for meats, wine, and beer; as a bleaching agent for foods; in pulp and paper industry; and in tanning. Express the equilibrium constant, k_2 , in terms of the equilibrium constant k_1 , in the two gaseous equilibria involving SO_2 . Namely,



Solution: An equilibrium expression is an equation where the equilibrium constant is equated with the ratio of concentrations of products to reactants, each raised to the power of its coefficients in the chemical reaction. For k_1 ,

$$k_1 = \frac{[SO_3]}{[SO_2][O_2]^{\frac{1}{2}}}$$

$$\text{For } k_2, \quad k_2 = \frac{[O_2][SO_2]^2}{[SO_3]^2}.$$

Now that this is written, k_2 can now be expressed in terms of k_1 by solving these equations in terms of some common concentration. For example, solve the k_1 equation for $[SO_3]$.

$$[SO_3] = k_1 [SO_2] [O_2]^{\frac{1}{2}}.$$

$$[SO_3]^2 = k_1^2 \times [SO_2]^2 [O_2].$$

Substitute this into the k_2 equation,

$$k_2 = \frac{[O_2][SO_2]^2}{[SO_3]^2}$$

The result is $k_2 = \frac{[O_2][SO_2]^2}{k_1^2 [SO_2]^2 [O_2]}$. This reduces to

$$k_2 = \frac{1}{k_1^2}.$$

APPLIED LIQUID AND SOLUTION PROBLEMS

• PROBLEM 874

The maximum safe concentration of fluoride ion in drinking water is 2.0 ppm (parts per million). Express this concentration in milligram %.

Solution: Milligram percent is the number of milligrams of solute in 100 ml. of liquid. It is given that there are 2 g. of fluoride in 1.0×10^6 g H_2O . Because 1 ml of H_2O weighs 1 g, there are 2 g of fluoride in 1×10^6 ml of H_2O . Solving for mg %:

$$\text{milligram \%} = \frac{100 \text{ ml} \times 2 \text{ g} \times 1000 \text{ mg/g}}{1 \times 10^6 \text{ ml}} = .20 \text{ mg \%}$$

The concentrations of substances in body fluids such as urine or blood are rather low, and for this reason they are generally expressed in milligram %.

• PROBLEM 875

Suppose a rain cloud has a water volume of 10^5 liters. If the cloud forms over a chemical factory which puts SO_3 into the air and each SO_3 molecule bonds with one out of 10^6 H_2O molecules (to form H_2SO_4), what is the molarity of the H_2SO_4 in the cloud?

Solution: To solve this problem, the number of moles of H_2SO_4 must be determined. This is done by applying the stoichiometric rules to the equation of this particular reaction:



According to this equation for every mole of SO_3 consumed, one mole of H_2SO_4 is produced or, in other words, if the

number of moles of SO_3 consumed is known, then the number of moles of H_2SO_4 produced is also known. The number of moles of SO_3 can be found since it is the same number as H_2O (from the stoichiometry of the equation). The number of moles of H_2O consumed can be found by dividing its weight by its molecular weight ($\text{MW} = 18.02$) or

$$\begin{aligned}\text{number of moles of H}_2\text{O} &= \frac{(10^5 \text{ liter})(1000 \text{ g/liter})}{18.02 \text{ g/mole}} \\ &= 5.5 \times 10^6 \text{ moles}\end{aligned}$$

Note: 1000 g/liter is the density of water (rain).

However, the number of water molecules is needed and thus,

$$\begin{aligned}\text{number of H}_2\text{O molecules} \\ &= (5.5 \times 10^6 \text{ moles})(6.02 \times 10^{23} \text{ molecules/mole}) \\ &= 3.3 \times 10^{30} \text{ molecules.}\end{aligned}$$

The problem states that SO_3 molecules bond with one out of 10^4 H_2O molecules. From this, the number of SO_3 molecules is determined,

$$\frac{3.3 \times 10^{30} \text{ H}_2\text{O molecules}}{10^4 \frac{\text{H}_2\text{O molecules}}{\text{SO}_3 \text{ molecules}}} = 3.3 \times 10^{26} \text{ SO}_3 \text{ molecules.}$$

The number of moles of SO_3 can now be obtained.

$$\begin{aligned}\text{number of moles of SO}_3 &= \frac{(3.3 \times 10^{26} \text{ SO}_3 \text{ molecules})}{6.02 \times 10^{23} \frac{\text{molecules}}{\text{mole}}} \\ &= 5.5 \times 10^2 \text{ moles}\end{aligned}$$

or 5.5×10^2 moles of H_2SO_4 are formed.

The molarity of the H_2SO_4 produced is thus

$$\frac{5.5 \times 10^2 \text{ moles}}{10^5 \text{ liter}} = 5.5 \times 10^{-3} \text{ M.}$$

• PROBLEM 876

What is the molarity of ethanol in 90-proof whiskey?
Density of alcohol = 0.8 g/ml.

Solution: Molarity is defined as the number of moles of solute per liter of solution. By definition, 90-proof whiskey is 45 % ethanol by volume. Thus in 1 l (= 1000 ml)

of whiskey there is (45 %)(1000 ml) = 450 ml of ethanol. To determine the number of moles of ethanol in 1 l of whiskey first calculate the weight of ethanol in 1 l by multiplying the density of ethanol by the volume of ethanol in 1 l. Then, divide the weight of ethanol in 1 l by its molecular weight (MW ethanol = 46 g/mole). Therefore,


$$(450 \text{ ml ethanol})(0.8 \text{ g/ml}) = 360 \text{ g ethanol}$$

$$\text{molarity} = \frac{\text{number moles ethanol}}{1 \text{ l whiskey}} = \frac{360 \text{ g}/46 \text{ g/mole}}{1 \text{ l}} = 7.83 \text{ M.}$$

• PROBLEM 877

A nurse mistakenly administers a dilute $\text{Ba}(\text{NO}_3)_2$ (a soluble, strong electrolyte) solution to a patient for radiographic investigation. What treatment would you provide to prevent the absorption of soluble barium and subsequent barium poisoning?

Solution: An electrolyte is a compound that conducts electricity when dissolved in water. Electrolytes are able to do this by dissociating into ions in water. Because $\text{Ba}(\text{NO}_3)_2$ is a strong electrolyte, it will completely dissociate into ions according to the reaction



To prevent barium poisoning, the Ba^{2+} ions, must be removed from the solution as an insoluble compound so that they cannot be absorbed. One way to precipitate out the barium ion (Ba^{2+}) is by adding a dilute solution of Na_2SO_4 . The reason is that Na_2SO_4 reacts with Ba^{2+} and NO_3^{-} ions to produce BaSO_4 and NaNO_3 . The barium sulfate is extremely insoluble as indicated by a solubility product constant of $1.6 \times 10^{-9} \text{ mole}^2 \text{ liter}^{-2}$. (A solubility product constant is indicative of a salt's solubility in solution. If this value is low, the salt is insoluble or barely soluble.)

Therefore, by adding the Na_2SO_4 , the Ba^{2+} precipitates out and is not absorbed.

• PROBLEM 878

A household cleaning solution has a hydronium ion concentration of 10^{-11} M . What is the pH of the solution? Is the solution acidic or basic?

Solution: The pH of any solution by definition is

determined by using the following equation.

$$\text{pH} = -\log [\text{H}^+]$$

Thus, if hydronium concentration = 10^{-11} M,

$$\text{pH} = -\log 10^{-11} \quad \text{or} \quad \text{pH} = 11.$$

The solution is basic because it has a pH greater than 7.

• PROBLEM 879

10 cc of battery acid (density 1.21) is diluted to 100 cc. 30 cc of the dilute solution is neutralized by 45 cc of 0.5 N base. What is the normality of the diluted acid? What is its strength in grams per liter? What is the strength of the undiluted 1.21 density acid in g/l?

Solution: To solve this problem, use the following relationship for the neutralization of acids and bases

$$N_a V_a = N_b V_b$$

where N_a is the normality of the acid at volume V_a and N_b is the normality of the base at volume V_b . Thus, the normality of the diluted acid is

$$N_a = \frac{N_b V_b}{V_a} = \frac{(0.5 \text{ N})(0.045 \text{ l})}{(0.030 \text{ l})} = 0.75 \text{ N}.$$

Note: 1 cc = .001 l.

The undiluted acid has a density of 1.21 g/cc; thus 10 cc of acid, weighs 12.1 g. Thus, when 90 cc of H_2O is added to form 100 cc of the diluted acid the weight of the entire solution becomes $90 + 12.1 = 102.1$ g, but the weight of the acid is still 12.1 g. In a liter of diluted acid, there are

$$\frac{12.1 \text{ g}}{100 \text{ cc}} \times \frac{10}{10} = \frac{121.0 \text{ g}}{1000 \text{ cc}} = 121 \text{ g/l of acid}.$$

For the undiluted acid, the density is:

$$\frac{1.21 \text{ g}}{1 \text{ cc}} \times \frac{1000}{1000} = \frac{1210 \text{ g}}{1000 \text{ cc}} = 1210 \text{ g/l}.$$

THE CHEMISTRY PROBLEM SOLVER®

Year 2003 Printing

**Copyright © 2001, 1998, 1996, 1993, 1977 by
Research & Education Association. All rights
reserved. No part of this book may be
reproduced in any form without permission of
the publisher.**

Printed in the United States of America

Library of Congress Control Number 00-134401

International Standard Book Number 0-87891-509-5

**PROBLEM SOLVER is a registered trademark of
Research & Education Association, Piscataway, New Jersey 08854**

CONTENTS

Chapter No.	Page No.
UNITS CONVERSION FACTORS	xiv
1 UNITS OF MEASUREMENT	
Basic Attacks and Strategies for Solving Problems	1-A
Length	1
Area	2
Volume	3
Mass	6
Density	8
Temperature	10
2 GASES	
Basic Attacks and Strategies for Solving Problems	13-A
Pressure	13
Boyle's Law, Charles' Law, Law of Gay-Lussac	19
Combined Gas Laws	27
Avogadro's Law-The Mole Concept	35
The Ideal Gas Law	38
3 GAS MIXTURES AND OTHER PHYSICAL PROPERTIES OF GASES	
Basic Attacks and Strategies for Solving Problems	61-A
Mole Fraction	61
Dalton's Law of Partial Pressures	63
Graham's Law of Gaseous Diffusion	79
Kinetic Theory of Gases	84
4 AVOGADRO'S HYPOTHESIS; CHEMICAL COMPOUNDS AND FORMULAS	
Basic Attacks and Strategies for Solving Problems	95-A
Atomic and Molecular Weights	101
Equivalent Weights	114
Chemical Composition-Weight and Volume Percent	119

5 STOICHIOMETRY/WEIGHT AND VOLUME CALCULATIONS

Basic Attacks and Strategies for Solving Problems	145-A
Balancing Equations	145
Calculations Using Chemical Arithmetic	147
Weight-Weight Problems	150
Reactions with Limiting Reagents	169
Volume-Volume Problems	177
Weight-Volume Problems	179

6 SOLIDS

Basic Attacks and Strategies for Solving Problems	191-A
Crystal Structure	191
Lattice Structures and Unit Cells	194
Phase Diagrams	210

7 PROPERTIES OF LIQUIDS

Basic Attacks and Strategies for Solving Problems	213-A
Density	213
Freezing Point Depression and Boiling Point Elevation .	216
Raoult's Law and Vapor Pressure	231
Clausius-Clapeyron Equation of Vaporization	239
Osmotic Pressure	247
Surface Tension	249

8 SOLUTION CHEMISTRY

Basic Attacks and Strategies for Solving Problems	251-A
Density and Formality	251
Molality	253
Molarity	255
Normality	266
Neutralization	272

9 EQUILIBRIUM

Basic Attacks and Strategies for Solving Problems	277-A
The Equilibrium Constant	277
Equilibrium Calculations	281
The Shifting of Equilibrium-Le Chatelier's Principle	297

10 ACID-BASE EQUILIBRIA

Basic Attacks and Strategies for Solving Problems	305-A
Acids and Bases	305
The Autoionization of Water	308

Autoprotolysis	309
pH	312
The Ionization Constant	318
The Dissociation Constant	325
The Hydrolysis Constant	334
Neutralization	339
Buffers	349
Indicators	354
Complex Ions	357
Electrolytes	363
11 SOLUBILITY AND THE ION PRODUCT CONSTANT	
Basic Attacks and Strategies for Solving Problems	367-A
12 CALCULATIONS USING pH AND THE DISSOCIATION CONSTANT	
Basic Attacks and Strategies for Solving Problems	385-A
13 CHEMICAL KINETICS	
Basic Attacks and Strategies for Solving Problems	423-A
The Rate Law	423
The Order of Reactions	432
Half-life	439
The Arrhenius Equations; Relating Temperature and	
Reaction Rate	445
Chain Reactions	452
14 THERMODYNAMICS I	
Basic Attacks and Strategies for Solving Problems	459-A
Bond Energies	460
Heat Capacity	464
Enthalpy	469
Enthalpy Calculations Using the First Law of	
Thermodynamics	486
Heats of Fusion and Vaporization	494
15 THERMODYNAMICS II	
Basic Attacks and Strategies for Solving Problems	503-A
Entropy	503
Free Energy	508
Equilibrium Calculations	517

16 ELECTROCHEMISTRY

Basic Attacks and Strategies for Solving Problems	539-A
Conduction	539
Equivalent Weight	540
Redox Reactions	545
Faraday's Law of Electrolysis	552
Electrode Potential	559
Electrochemical Cell Reactions	564
Nernst Equation	567

17 ATOMIC THEORY

Basic Attacks and Strategies for Solving Problems	581-A
Atomic Weight	581
Valence and Electron Dot Diagrams	583
Ionic and Covalent Bonding	588
Electronegativity	592
Bond Length and Angles	601
Polarity of Bonds	606

18 QUANTUM CHEMISTRY

Basic Attacks and Strategies for Solving Problems	613-A
Pauli Exclusion Principle, Hund's Rule and	
Electronic Configuration	613
Molecular Orbital Theory	624
Early Quantum Chemistry	630
The Bohr Atom	637
The de Broglie Equation	640
Wave Functions	643
Atomic Spectroscopy	649

19 NUCLEAR CHEMISTRY

Basic Attacks and Strategies for Solving Problems	671-A
---	-------

20 ORGANIC CHEMISTRY I: NOMENCLATURE AND STRUCTURE

Basic Attacks and Strategies for Solving Problems	687-A
Alkanes	687
Alkenes and Alkynes	697
Alcohols	702
Other Functional Groups	706

21 ORGANIC CHEMISTRY II: REACTIONS

Basic Attacks and Strategies for Solving Problems	717-A
Alkanes	717
Alkenes and Alkynes	723
Alcohols	731
Other Functional Groups	736

22 BIOCHEMISTRY I

Basic Attacks and Strategies for Solving Problems	755-A
Cellular Construction and Dimensions	755
pH and pKa	759
Energy Conversion	766
Molecular Weights, Mole Ratios, Densities and Monomer Units	770
Physical Aspects of Biochemistry	780

23 BIOCHEMISTRY II

Basic Attacks and Strategies for Solving Problems	789-A
Proteins	789
Enzymes	795
Carbohydrates	800
Nucleic Acids	804

24 APPLIED POLLUTION/ENVIRONMENTAL PROBLEMS..... 809

25 APPLIED GAS AND GAS MIXTURE PROBLEMS 825

26 APPLIED LIQUID AND SOLUTION PROBLEMS 837

27 APPLIED STOICHIOMETRY PROBLEMS 847

28 APPLIED THERMOCHEMISTRY PROBLEMS 863

29 APPLIED ENERGY PROBLEMS 869

30 APPLIED WAVE PHENOMENA PROBLEMS 873

31 APPLIED ORGANIC AND POLYMER CHEMISTRY PROBLEMS..... 879

32 APPLIED BIOLOGICAL PROBLEMS 885

INDEX..... 908

UNITS CONVERSION FACTORS

This section includes a particularly useful and comprehensive table to aid students and teachers in converting between systems of units.

The problems and their solutions in this book use SI (**I**nternational **S**ystem) as well as English units. Both of these units are in extensive use throughout the world, and therefore students should develop a good facility to work with both sets of units until a single standard of units has been found acceptable internationally.

In working out or solving a problem in one system of units or the other, essentially only the numbers change. Also, the conversion from one unit system to another is easily achieved through the use of conversion factors that are given in the subsequent table. Accordingly, the units are one of the least important aspects of a problem. For these reasons, a student should not be concerned mainly with which units are used in any particular problem. Instead, a student should obtain from that problem and its solution an understanding of the underlying principles and solution techniques that are illustrated there.

To convert	To	Multiply by	For the reverse, multiply by
acres	square feet	4.356×10^4	2.296×10^{-5}
acres	square meters	4047	2.471×10^{-4}
ampere-hours	coulombs	3600	2.778×10^{-4}
ampere-turns	gilberts	1.257	0.7958
ampere-turns per cm. ..	ampere-turns per inch	2.54	0.3937
angstrom units	inches	3.937×10^{-8}	2.54×10^8
angstrom units	meters	10^{-10}	10^{10}
atmospheres	feet of water	33.90	0.02950
atmospheres	inch of mercury at 0°C	29.92	3.342×10^{-2}
atmospheres	kilogram per square meter	1.033×10^4	9.678×10^{-5}
atmospheres	millimeter of mercury at 0°C	760	1.316×10^{-3}
atmospheres	pascals	1.0133×10^5	0.9869×10^{-5}
atmospheres	pounds per square inch	14.70	0.06804
bars	atmospheres	9.870×10^{-7}	1.0133
bars	dynes per square cm.	10^6	10^{-6}
bars	pascals	10^5	10^{-5}
bars	pounds per square inch	14.504	6.8947×10^{-2}
Btu	ergs	1.0548×10^{10}	9.486×10^{-11}
Btu	foot-pounds	778.3	1.285×10^{-3}
Btu	joules	1054.8	9.480×10^{-4}
Btu	kilogram-calories	0.252	3.969
calories, gram	Btu	3.968×10^{-3}	252
calories, gram	foot-pounds	3.087	0.324
calories, gram	joules	4.185	0.2389
Celsius	Fahrenheit	$(^{\circ}\text{C} \times 9/5) + 32 = ^{\circ}\text{F}$	$(^{\circ}\text{F} - 32) \times 5/9 = ^{\circ}\text{C}$

To convert	To	Multiply	For the reverse, multiply by
Celsius	kelvin	$^{\circ}\text{C} + 273.1 = \text{K}$	$\text{K} - 273.1 = ^{\circ}\text{C}$
centimeters	angstrom units	1×10^8	1×10^{-8}
centimeters	feet	0.03281	30.479
centistokes	square meters per second	1×10^{-6}	1×10^6
circular mils	square centimeters	5.067×10^{-8}	1.973×10^5
circular mils	square mils	0.7854	1.273
cubic feet	gallons (liquid U.S.)	7.481	0.1337
cubic feet	liters	28.32	3.531×10^{-2}
cubic inches	cubic centimeters	16.39	6.102×10^{-2}
cubic inches	cubic feet	5.787×10^{-4}	1728
cubic inches	cubic meters	1.639×10^{-5}	6.102×10^4
cubic inches	gallons (liquid U.S.)	4.329×10^{-3}	231
cubic meters	cubic feet	35.31	2.832×10^{-2}
cubic meters	cubic yards	1.308	0.7646
curies	coulombs per minute	1.1×10^{12}	0.91×10^{-12}
cycles per second	hertz	1	1
degrees (angle)	mils	17.45	5.73×10^{-2}
degrees (angle)	radians	1.745×10^{-2}	57.3
dynes	pounds	2.248×10^{-6}	4.448×10^5
electron volts	joules	1.602×10^{-19}	0.624×10^{18}
ergs	foot-pounds	7.376×10^{-8}	1.356×10^7
ergs	joules	10^{-7}	10^7
ergs per second	watts	10^{-7}	10^7
ergs per square cm.	watts per square cm.	10^{-3}	10^3
Fahrenheit	kelvin	$(^{\circ}\text{F} + 459.67)/1.8$	$1.8\text{K} - 459.67$
Fahrenheit	Rankine	$^{\circ}\text{F} + 459.67 = ^{\circ}\text{R}$	$^{\circ}\text{R} - 459.67 = ^{\circ}\text{F}$
faradays	ampere-hours	26.8	3.731×10^{-2}
feet	centimeters	30.48	3.281×10^{-2}
feet	meters	0.3048	3.281
feet	mils	1.2×10^4	8.333×10^{-5}
fermis	meters	10^{-15}	10^{15}
foot candles	lux	10.764	0.0929
foot lamberts	candelas per square meter	3.4263	0.2918
foot-pounds	gram-centimeters	1.383×10^4	1.235×10^{-5}
foot-pounds	horsepower-hours	5.05×10^{-7}	1.98×10^6
foot-pounds	kilogram-meters	0.1383	7.233
foot-pounds	kilowatt-hours	3.766×10^{-7}	2.655×10^6
foot-pounds	ounce-inches	192	5.208×10^{-3}
gallons (liquid U.S.)	cubic meters	3.785×10^{-3}	264.2
gallons (liquid U.S.)	gallons (liquid British Imperial)	0.8327	1.201
gammas	teslas	10^{-9}	10^9
gausses	lines per square cm.	1.0	1.0
gausses	lines per square inch	6.452	0.155
gausses	teslas	10^{-4}	10^4
gausses	webers per square inch	6.452×10^{-8}	1.55×10^7
gilberts	amperes	0.7958	1.257
grads	radians	1.571×10^{-2}	63.65
grains	grams	0.06480	15.432
grains	pounds	$\frac{1}{7000}$	7000
grams	dynes	980.7	1.02×10^{-3}
grams	grains	15.43	6.481×10^{-2}

To convert	To	Multiply	For the reverse, multiply by
grams	ounces (avdp)	3.527×10^{-2}	28.35
grams	pounds	7.093×10^{-2}	14.1
hectares	acres	2.471	0.4047
horsepower	Btu per minute	42.418	2.357×10^{-2}
horsepower	foot-pounds per minute	3.3×10^4	3.03×10^{-5}
horsepower	foot-pounds per second	550	1.182×10^{-3}
horsepower	horsepower (metric)	1.014	0.9863
horsepower	kilowatts	0.746	1.341
inches	centimeters	2.54	0.3937
inches	feet	8.333×10^{-2}	12
inches	meters	2.54×10^{-2}	39.37
inches	miles	1.578×10^{-5}	6.336×10^4
inches	mils	10^3	10^{-3}
inches	yards	2.778×10^{-2}	36
joules	foot-pounds	0.7376	1.356
joules	watt-hours	2.778×10^{-4}	3600
kilograms	tons (long)	9.842×10^{-4}	1016
kilograms	tons (short)	1.102×10^{-3}	907.2
kilograms	pounds (avdp)	2.205	0.4536
kilometers	feet	3281	3.408×10^{-4}
kilometers	inches	3.937×10^4	2.54×10^{-5}
kilometers per hour	feet per minute	54.68	1.829×10^{-2}
kilowatt-hours	Btu	3413	2.93×10^{-4}
kilowatt-hours	foot-pounds	2.655×10^6	3.766×10^{-7}
kilowatt-hours	horsepower-hours	1.341	0.7457
kilowatt-hours	joules	3.6×10^6	2.778×10^{-7}
knots	feet per second	1.688	0.5925
knots	miles per hour	1.1508	0.869
lamberts	candles per square cm.	0.3183	3.142
lamberts	candles per square inch	2.054	0.4869
liters	cubic centimeters	10^3	10^{-3}
liters	cubic inches	61.02	1.639×10^{-2}
liters	gallons (liquid U.S.)	0.2642	3.785
liters	pints (liquid U.S.)	2.113	0.4732
lumens per square foot	foot-candles	1	1
lumens per square meter	foot-candles	0.0929	10.764
lux	foot-candles	0.0929	10.764
maxwells	kilolines	10^{-3}	10^3
maxwells	webers	10^{-8}	10^8
meters	feet	3.28	30.48×10^{-2}
meters	inches	39.37	2.54×10^{-2}
meters	miles	6.214×10^{-4}	1609.35
meters	yards	1.094	0.9144
miles (nautical)	feet	6076.1	1.646×10^{-4}
miles (nautical)	meters	1852	5.4×10^{-4}
miles (statute)	feet	5280	1.894×10^{-4}
miles (statute)	kilometers	1.609	0.6214
miles (statute)	miles (nautical)	0.869	1.1508
miles per hour	feet per second	1.467	0.6818
miles per hour	knots	0.8684	1.152
millimeters	microns	10^3	10^{-3}

To convert	To	Multiply	For the reverse, multiply by
mils	meters	2.54×10^{-5}	3.94×10^4
mils	minutes	3.438	0.2909
minutes (angle)	degrees	1.666×10^{-2}	60
minutes (angle)	radians	2.909×10^{-4}	3484
newtons	dynes	10^5	10^{-5}
newtons	kilograms	0.1020	9.807
newtons per sq. meter ..	pascals	1	1
newtons	pounds (avdp)	0.2248	4.448
oersteds	amperes per meter	7.9577×10	1.257×10^{-2}
ounces (fluid)	quarts	3.125×10^{-2}	32
ounces (avdp)	pounds	6.25×10^{-2}	16
pints	quarts (liquid U.S.)	0.50	2
poundals	dynes	1.383×10^4	7.233×10^{-5}
poundals	pounds (avdp)	3.108×10^{-2}	32.17
pounds	grams	453.6	2.205×10^{-3}
pounds (force)	newtons	4.4482	0.2288
pounds per square inch ..	dynes per square cm.	6.8946×10^4	1.450×10^{-5}
pounds per square inch ..	pascals	6.895×10^3	1.45×10^{-4}
quarts (U.S. liquid)	cubic centimeters	946.4	1.057×10^{-3}
radians	mils	10^3	10^{-3}
radians	minutes of arc	3.438×10^3	2.909×10^{-4}
radians	seconds of arc	2.06265×10^5	4.848×10^{-6}
revolutions per minute ..	radians per second	0.1047	9.549
roentgens	coulombs per kilogram	2.58×10^{-4}	3.876×10^3
slugs	kilograms	1.459	0.6854
slugs	pounds (avdp)	32.174	3.108×10^{-2}
square feet	square centimeters	929.034	1.076×10^{-3}
square feet	square inches	144	6.944×10^{-3}
square feet	square miles	3.587×10^{-8}	27.88×10^6
square inches	square centimeters	6.452	0.155
square kilometers	square miles	0.3861	2.59
stokes	square meter per second	10^{-4}	10^{-4}
tons (metric)	kilograms	10^3	10^{-3}
tons (short)	pounds	2000	5×10^{-4}
torrs	newtons per square meter	133.32	7.5×10^{-3}
watts	Btu per hour	3.413	0.293
watts	foot-pounds per minute	44.26	2.26×10^{-2}
watts	horsepower	1.341×10^{-3}	746
watt-seconds	joules	1	1
webers	maxwells	10^8	10^{-8}
webers per square meter ..	gausses	10^4	10^{-4}

CHAPTER 1

UNITS OF MEASUREMENT

Basic Attacks and Strategies for Solving Problems in this Chapter. See pages 1 to 12 for step-by-step solutions to problems.

Any measured or calculated quantity has two parts — a numerical value and a unit. Sometimes, when the unit is likely to be understood, it is not written. For example, if the United States Weather Service reports that the temperature in Phoenix is 108, it is understood to be 108°F. However, fewer errors in scientific and engineering calculations are made when the rule of always writing both the value and the unit for each quantity is strictly observed as illustrated below.

2 grams (2 g), 0.5 seconds (0.5 s), 6.28 meters (6.28 m), 98.6°F

There are only seven quantities — length, mass, time, thermodynamic (or absolute) temperature, electric current, amount of a substance, and luminous intensity — for which base units are defined and from which all other units can be derived. These quantities and their respective units and symbols are listed in the table below.

Table 1		
PHYSICAL QUANTITY	SI UNIT	SYMBOL
Length	meter	m
Mass	kilogram	kg
Time	second	s
Amt. of substance	mole	mol
Thermodynamic temp.	kelvin	K
Electric current	ampere	A
Luminous intensity	candela	cd

A unit of velocity is derived, for example, by dividing a unit of length by a unit of time to obtain miles per hour, feet per second, furlongs per fortnight, etc. In this chapter, only base units for length, mass, and absolute temperature are discussed, along with derived units of area (length squared), volume (length cubed), density (mass divided by volume), and the common Celsius ($^{\circ}\text{C}$) and Fahrenheit ($^{\circ}\text{F}$) temperature scales. If the rules illustrated with these units are learned and understood, they can be applied to any units that may be encountered.

Units can be treated like ordinary algebraic variables in equations when quantities are added, subtracted, multiplied, or divided. Two rules, borrowed from algebra, must be remembered:

1. The numerical values of two quantities can be added or subtracted **only** if the units are the same. This is true whether or not the units are the same quantity — e.g., the mixture of mass and length units in (b) or the two different length units in (c) below.

a) $5 \text{ feet} - 2 \text{ feet} = 3 \text{ feet}$ ($5x - 2x = 3x$)

b) $5 \text{ grams} - 2 \text{ feet} = ?$ ($5y - 2x = ?$)

c) $5 \text{ meters} - 2 \text{ feet} = ?$ ($5z - 2x = ?$)

2. Numerical values and their associated units can **always** be multiplied or divided.

a) $15 \text{ grams} \div 5 \text{ cm}^3 = 3 \text{ g/cm}^3$ ($15y \div 5x = 3y/x$)

b) $7 \text{ cm} \times 5 \text{ cm} = 35 \text{ cm}^2$ ($7y \times 5y = 35y^2$)

Conversion of Units

Most new scientific work is now performed and reported in the International System of Units (SI units) but, for practical reasons, many other systems of units are still in use and will remain so for many years. It is essential, therefore, that rules for converting from one system or set of units to another be understood. There is one simple rule that covers most cases for converting from one unit or set of units to another:

Multiply or divide by the appropriate conversion factor or factors including the units — i.e., be certain to multiply and/or divide the units of each conversion factor as well as the numerical values.

The exception to this rule occurs when the zero points of two units, such as exists with the Celsius and Fahrenheit temperature scales, are different. The conversion then requires a linear transformation — i.e., $y = mx + b$, where m is the ratio of the magnitude of the unit y to the unit x , and b is the value of y when x is zero. Except for the example of temperature scales, such cases are rare, indeed. In virtually all other cases, the zero point is independent of the unit; a velocity of zero, for example, is zero in any velocity unit — ft/sec, miles/hr, m/sec, etc.

Conversion factors are tabulated in handbooks but, in simple cases, can be drawn from memory. Conversion factors for units formed by adding the common prefixes — centi (c), milli (m), kilo (k), etc. — are simply the factors associated with those prefixes — 10^{-2} , 10^{-3} , 10^3 , etc., — respectively. Let us convert, as an example, a length measurement from feet to inches. To do this we multiply or divide by the appropriate conversion factor including its units. In this case, most people know the conversion factor(s) from memory — 12 inches/ft or 1 ft/12 inches.

$$15 \text{ ft} \times 12 \text{ inches/ft} = 180 \text{ inches} \quad 1-1$$

Note that the numerical values ($15 \times 12 = 180$) and the units (ft \times inches/ft = inches) are each multiplied separately to obtain the correct answer which includes both a value and a unit. The reciprocal conversion factor, 1 ft/12 inches, can also be used, but it is necessary, then, to divide rather than to multiply.

$$15 \text{ ft} \div 1 \text{ ft/12 inches} = (15 \div 1/12)[\text{ft} \div (\text{ft/in})] = 180 \text{ in} \quad 1-2$$

Observe how careful multiplication or division of the units as well as the numerical values can help to avoid errors in the conversion of units. In this case, suppose that the conversion factor in the first equation, 12 in/ft, was used but division rather than multiplication was employed mistakenly.

$$15 \text{ ft} \div 12 \text{ inches/ft} = (15/12)[\text{ft} \div \text{in/ft}] = 1.25 \text{ ft}^2/\text{in} \quad 1-3$$

The unexpected unit in the answer signals that an error has been made.

Several successive multiplication and/or division steps can be used to convert one or more units. If careful multiplication and/or division of the units, as well as the numerical values is observed, the correct unit will result. If the expected unit is not obtained, the probability of an error in the calculation is signaled. Let us convert, as an example, the density of water,

62.4 lb/ft³, to the appropriate value in grams per cubic centimeter. The answer, in this case, is known in advance to be 1.0 g/cm³ or 1.0 g/cc. The needed conversion factors (454 g/lb, 2.54 cm/in, 12 in/ft) can be obtained from a handbook with a table of conversion factors.

$$62.4 \text{ lb/ft}^3 \times 454 \text{ g/lb} + [12 \text{ in/ft} \times 2.54 \text{ cm/in}]^3 =$$

$$62.4 \times 454 + (12 \times 2.54)^3 [\text{lb/ft}^3 \times \text{g/lb} + (\text{in/ft} \times \text{cm/in})^3] = 1.0 \text{ g/cm}^3$$

1-4

Many tables of conversion factors contain a factor for conversion directly from centimeters to feet (30.48 cm/ft) and from cubic centimeters to cubic feet (28,317 cm³/ft³). These factors can also be used and the calculation is simplified as follows.

$$62.4 \text{ lb/ft}^3 \times 454 \text{ g/lb} + (30.48 \text{ cm/ft})^3 = 1.0 \text{ g/cm}^3$$

1-5

$$62.4 \text{ lb/ft}^3 \times 454 \text{ g/lb} + 28,317 \text{ cm}^3/\text{ft}^3 = 1.0 \text{ g/cm}^3$$

1-6

As shown, more than one procedure can lead to the correct answer, and some may result in fewer or simpler calculations than others. If careful attention is paid to the multiplication and division of units as well as numerical values, the chance of making an error is diminished.

From the pioneer in Study and Solution Guides...

REA's

Problem Solvers®

Complete Solution Guides to All Textbooks

**Why Students & Instructors Choose the PROBLEM SOLVERS
and What They Say About Them:**

"They are more helpful and instructive than the usual textbook solution manuals."

REA's Problem Solvers solve not only the simple problems, but also those difficult problems not found in study/solution manuals. It's the difficult ones that you encounter on tests.

"REA's Problem Solvers don't just promise. They deliver."

The Problem Solvers cover all assigned topics in the textbook. They will help no matter what textbook is used.

"They produce top grades for students."

You can expect to be well-prepared for any exam on any topic in any subject.

"...Thank you for the superb work you have done in publishing the Problem Solvers. These books are the best review books on the market ..." – Student, New Orleans, Louisiana

"... We are especially pleased with the Chemistry Problem Solver. We have never seen a supplementary book as valuable as this one ..." – Science Teacher, La Habra, California

"...I am very pleased with the content and presentation of your Chemistry Problem Solver ..." – Student, Wardensville, West Virginia

AVAILABLE IN THE FOLLOWING SUBJECTS:

Accounting	Economics	Heat Transfer	Pre-Calculus
Advanced Calculus	Electric Circuits	Linear Algebra	Probability
Algebra &	Electrical	Machine Design	Psychology
Trigonometry	Machines	Mathematics for	Statistics
Automatic Control	Electromagnetics	Engineers	Strength of Materials &
Systems/Robotics	Electronic	Mechanics	Mechanics of Solids
Biology	Communications	Numerical Analysis	Technical Design
Business, Accounting, & Finance	Electronics	Operations	Graphics
Calculus	Finite & Discrete	Research	Thermodynamics
Chemistry	Math	Optics	Topology
Complex Variables	Fluid Mechanics	Organic Chemistry	Transport
Differential Equations	Genetics	Physical Chemistry	Phenomena
	Geometry	Physics	Vector Analysis

We have made every effort to make this the best book possible. Our paper is opaque, with minimal show-through; it will not discolor or become brittle with age. Pages will not drop out. The binding will not crack or split. This is a permanent book.

Visit our website at www.rea.com



Research & Education Association

61 Ethel Road West • Piscataway, New Jersey 08854

ISBN 0-87891-509-5



9 780878 915095